

B B

TRI-PP-87-46
May 1987

19 OCT. 1987



- 2 -

C |

Gamow-Teller Strength Deduced from $^{54}\text{Fe}(\text{n},\text{p})^{54}\text{Mn}$ Cross-Sections at 298 MeV.

M.C. Vetterli¹, O. Häusser^{1,2}, W.P. Alford³,
D. Frekers^{1,4}, R. Helmer³, R. Henderson^{1,5},
K. Hicks¹, K.P. Jackson¹, R.G. Jeppesen^{1,2},
C.A. Miller¹, M.A. Moinester^{1,6}, K. Raywood⁵,
S. Yen¹.

- 1) TRIUMF, 4004 Wesbrook Mall, Vancouver, British Columbia, Canada, V6T 2A3.
- 2) Simon Fraser University, Burnaby, British Columbia, Canada, V5A 1S6.
- 3) University of Western Ontario, London, Ontario, Canada, N6A 3K7.
- 4) University of Toronto, Toronto, Ontario, Canada, M5S 1A7.
- 5) University of Melbourne, Melbourne, Australia.
- 6) Tel Aviv University, 69978 Ramat Aviv, Israel.

ABSTRACT

Gamow-Teller (GT) strength has been extracted from cross-sections for the $^{54}\text{Fe}(\text{n},\text{p})^{54}\text{Mn}$ reaction at 0° , 2.5° , 5° , 7° , and 12° , and for an incident energy of 298 MeV. This determination of the β^+ GT strength, S_+ , allows for the first time, a full test of the Ikeda sum rule. These data also show that the best available shell model calculations overestimate this strength by a factor of 2. This result is significant in models of supernovae since electron capture in (fp) shell nuclei, which is proportional to S_+ , depletes the relativistic electron gas in pre-collapse stars.

NUCLEAR REACTION: $^{54}\text{Fe}(\text{n},\text{p})^{54}\text{Mn}$, $E= 298$ MeV, angular distribution.
PACS nos.: 24.30.Cz, 25.40.Fq, 27.40.+z

Charge exchange reactions at small momentum transfer have been very useful for the study of β decay strength distributions, especially in states for which β decay is not energetically possible ^{1,2}. In particular, V_{sr} , the spin-isospin flip component of the nucleon-nucleus interaction, dominates over V_r , the non spin flip component, suppressing Fermi transitions in (p,n) reactions above about 100 MeV incident energy. Furthermore, because of the neutron excess in nuclei, isospin selection rules do not allow Fermi transitions in (n,p) reactions. Therefore, intermediate energy nucleon charge exchange reactions are excellent filters of Gamow-Teller (GT) strength ($\Delta T=1$, $\Delta S=1$, $\Delta L=0$, $0\hbar\omega$ excitations) since their 0° cross-section is proportional to $B(\text{GT})$, the β decay strength.

These reactions can be used to test the sum rule ³:

$$S_- - S_+ = 3(N-Z),$$

where S_- is the integrated GT strength in β^- decay which is related to the 0° (p,n) cross-section, S_+ is the integrated GT strength in β^+ decay, related to 0° (n,p) cross-sections, and $N-Z$ is the neutron excess of the target. Previous (p,n) results have suggested that as much as 40% of the sum rule strength is missing. This factor is common to nuclei spanning the periodic table ⁴. The value of 40% assumed $S_+=0$ due to Pauli blocking in neutron rich nuclei, since there was essentially no experimental information on S_+ until now.

In this letter, we present results for the $^{54}\text{Fe}(\text{n},\text{p})^{54}\text{Mn}$ reaction which, when combined with the $^{54}\text{Fe}(\text{p},\text{n})^{54}\text{Co}$ results of Rapaport *et al*⁵, will allow the first full test of the sum rule. GT excitations are not blocked in the (n,p) ($\Delta T_z=+1$) case because of the $\pi(f_{7/2}) \rightarrow \nu(f_{5/2})$ transition. Indeed, GT strength has been observed in the $^{54}\text{Fe}(t,^3\text{He})^{54}\text{Mn}$ reaction at 25 MeV ⁶. We will also show that shell model calculations by Bloom and Fuller ⁷ and by Muto ⁸ overestimate S_+ by at least a factor of two. This result is of interest in astrophysics since S_+ is related to electron capture rates in (fp) shell nuclei which influence the electron to baryon ratio, Y_e , stars prior to gravitational collapse. In turn, Y_e influences the mass of the core which will collapse ⁷.

Cross-sections for $^{54}\text{Fe}(\text{n},\text{p})^{54}\text{Mn}$ were measured at five angles between 0° and

(Submitted to Physical Review Letter)

CERN LIBRARIES, GENEVA



CM-P00068247

PLEASE
MAKE A
PHOTOCOPY
or check out as
NORMAL
LOAN

12° using the charge exchange (CHARGELEX) facility of the TRIUMF medium resolution spectrometer (MRS). Nearly monoenergetic neutrons with an energy of 298 MeV were produced by the $^7\text{Li}(\text{p},\text{n})^7\text{Be}$ reaction. The proton beam was swept away by a compact dipole magnet into a shielded beam dump. The secondary neutron beam then struck targets located at the pivot of the MRS. Protons from the (n,p) reaction were analysed by the spectrometer in the usual way⁹ with the addition of a second front end wire chamber to provide better ray tracing back to the target. The CHARGELEX facility is described in more detail elsewhere¹⁰. The target consisted of six layers separated by wire planes. By considering the pattern of hit wires in these planes, it was possible to determine in which layer the (n,p) reaction occurred. Corrections were made for the energy loss in subsequent target layers in order to recover good resolution (≈ 1 MeV). This technique allows the use of a thick segmented target or the simultaneous acquisition of data from different target types. A full description of the target box can be found elsewhere¹¹. The configuration used for this experiment was: empty, ^{12}C (46 mg/cm²), empty, ^{54}Fe (140 mg/cm²), ^{54}Fe (140 mg/cm²), CH_2 (47.2 mg/cm²). The empty target locations were used to determine the contribution to the spectra from the wire chamber gas and windows. The isolated peak from the $^1\text{H}(\text{n},\text{p})$ reaction in CH_2 provided the means for normalization, simultaneously collected with the data from the primary targets. A value of 53.6 mb/sr in the lab frame was assumed for this cross-section at 0°; this was obtained from the Arndt 1986 phase shifts¹². The carbon contribution was subtracted from the CH_2 spectrum to obtain the energy profile of the incident neutron beam. The latter exhibited a low energy tail, of intensity $\approx 1\%/\text{MeV}$ of the nearly monoenergetic neutron peak, which was unfolded from the data.

Results for the $^{54}\text{Fe}(\text{n},\text{p})^{54}\text{Mn}$ cross-section at 298 MeV are shown in figure 1. Of note is the strongly forward peaked feature below 10 MeV excitation. This is the Gamow-Teller resonance and its location and shape are reasonably well reproduced by the calculation of Bloom and Fuller⁷ (see the 0° spectrum). Although the theory agrees qualitatively with the data, it must be renormalized to fit the magnitude of the peak.

The $\Delta L=0$, GT component of the cross-section was determined by a multipole decomposition using angular momentum transfers up to $\Delta L=2$. The data were summed in 1 MeV bins and the resulting angular distributions were fitted to the theoretical curves generated by DWIA calculations¹³ with the Love-Franey interaction at 270 MeV¹⁴. The contribution of each multipolarity was thus determined for each bin. For more details see reference¹⁵. The angular distributions were calculated for simple 1 particle-3 hole configurations: $[\pi(f_{7/2})^{-3}\nu(f_{5/2})]_{1+}$ for $\Delta L=0$, $[\pi(f_{7/2})^{-3}\nu(g_{9/2})]_{1-}$ for $\Delta L=1$, and $[\pi(f_{7/2})^{-3}\nu(f_{5/2})]_{3+}$ for $\Delta L=2$. These configurations are analogous to those used in the analysis of $^{58}\text{Ni}(\text{p},\text{n})^{58}\text{Cu}$ by Rapaport et al⁵. The Q-value dependence of the shapes was accounted for by calculating the angular distributions in 10 MeV intervals from 0 MeV to 40 MeV and interpolating for excitation energies in between. The optical potentials used for the distortions in DW81 were of the Woods-Saxon form and were obtained from $^{54}\text{Fe}(\text{p},\text{p})$ elastic data taken at TRIUMF¹⁶. The results of the decomposition are plotted for 0° in figure 2. The low energy peak is well fitted. However, the decomposition produces a tail at high energy for the $\Delta L=0$ component. The amount of $\Delta L=0$ strength above 10 MeV is uncertain and depends on the assumed shape of the $\Delta L=1$ angular distribution. Since the $\Delta L=1$ contribution dominates between 10 MeV and 30 MeV, it is possible that a small change in the $\Delta L=1$ shape could produce spurious $\Delta L=0$ strength in this region of excitation energy. The integrated cross-section below 10 MeV is 16.9 ± 1.1 mb/sr excluding the uncertainty on the $^1\text{H}(\text{n},\text{p})$ normalization. Of this, 12.9 mb/sr, or 76%, is $\Delta L=0$ as determined by the multipole decomposition. Ascribing an error of ± 5 to this percentage because of uncertainties in the shapes of the angular distributions, we get 12.9 ± 1.2 mb/sr for the 0°, $\Delta L=0$ cross-section below 10 MeV. We note that the $\Delta L=0$ cross-section between 10 MeV and 40 MeV is 8.5 mb/sr. However, since the $\Delta L=0$ component is uncertain above 10 MeV, we will not consider strength above this energy in the present analysis.

Due to the absence of a strong, known β^- decay in the (fp) shell with which to calibrate the (n,p) reaction, the ratio $\sigma(0^\circ)/B(GT^+)$ of the (n,p) cross-section to the β^+ decay strength had to be calculated theoretically. This was done using the com-

puter program OXBASH¹⁷ with the simple $\pi(f_{7/2})^{-3}\nu(f_{5/2})$ configuration to obtain $B(GT^+)$ and the one body transition densities needed for input to DW81 which was used to calculate $\sigma(0^\circ)$. As a check, we obtained $\sigma(0^\circ, q=0)/B(GT^+) \approx 7 \text{ mb/sr}$ for ^{54}Fe at 160 MeV, in agreement with the calculation shown in fig. 6 of reference². A further test of the calculation was done at 300 MeV using our data for the $^{12}\text{C}(\text{n},\text{p})^{12}\text{B}_{g.s.}$ transition. The ratio was found to be 7.1 experimentally, as predicted by DWIA. Hence, we can use this method for obtaining $\sigma/B(GT)$ with some confidence. A value of 3.4 was calculated for the ratio $\sigma(0^\circ)/B(GT)$ in ^{54}Fe at 300 MeV. Note that this is not extrapolated to $q=0$ but is simply the ratio at 0° . Using this result, we obtain: $S_+ = 3.8 \pm 0.4$. Bloom and Fuller have predicted $S_+ = 9.1$ for a 2 particle-4 hole ($2p-4h$) model⁷. This model includes (1p-1h) excitations from $f_{7/2}$ to $p_{3/2}, p_{1/2}$, and $f_{5/2}$ in both the parent and daughter nuclei. Muto has done a similar calculation and obtains comparable results ($S_+ = 9.4$)⁸. Thus, these calculations overestimate GT strength by a factor of 2.4. A likely cause of the large discrepancy is the severe truncation of the model space used in the calculations. This truncation has more drastic consequences for (n,p) reactions because they are more sensitive to the details of the Fermi surface than either (p,n) or (p,p) reactions. Indeed, in reference⁸, Muto extends the model space to include 2p-2h excitations in the parent nucleus and calculates $S_+ = 5.1$. However, since 2p-2h excitations are not included in the daughter nucleus, this model does not satisfy the sum rule since $\pi(f_{7/2}) \rightarrow \nu(f_{5/2})$ transitions are not allowed from 2p-2h parent configurations. Using a sum rule technique which eliminates the daughter wave function from the evaluation of the total GT strength, Muto estimates $S_+ = 7.4$, 20% smaller than the simpler 1p-1h model. However, this still overestimates the experimental value by a factor of almost 2, ($S_+^{(\text{exp})}/S_+^{(\text{th})} = 0.51$). Similar truncation effects have been seen in the (sd) shell by Wildenthal and others¹⁸. In other studies, Auerbach *et al.*¹⁹ and Cha²⁰ have shown that the use of RPA wave functions instead of the simplest shell model reduces the calculated GT⁺ strength by a factor of about 2. Thus, it is crucial to consider the largest possible configuration space in the calculation of GT strength for (fp) shell nuclei.

As mentioned above, these results are important for models of supernovae because the relativistic electron gas which resists the gravitational collapse of the star, is depleted by electron capture. Thus, the rate of this capture influences the mass of the core which will collapse and eventually create an outward going shock wave which may lead to a supernova explosion.

Our result of $S_+ = 3.8 \pm 0.4$ can be used in conjunction with the $^{54}\text{Fe}(\text{p},\text{n})^{54}\text{Co}$ result⁵, $S_- = 7.8 \pm 1.9$, to test the sum rule. We get: $S_- - S_+ = 4.0 \pm 2.0$. This is 67% of the sum rule value of 6. However, the large uncertainty also makes $(S_- - S_+)$ consistent with the sum rule. Angular distributions for the $^{54}\text{Fe}(\text{p},\text{n})^{54}\text{Co}$ reaction at 300 MeV have been measured at TRIUMF²¹. It is hoped that the $\Delta L=0$ component can be extracted reliably from these data and that the uncertainty in S_- can be reduced. More definite conclusions on the sum rule must await this analysis. Furthermore, we recall that the possible $\Delta L=0$ (n,p) strength above 10 MeV excitation has been ignored in this analysis. A complete study of the sum rule will also have to take this into account.

The authors wish to thank Drs. G. Fuller and S. Bloom for providing their calculations and for useful discussions on the theory related to this paper. This work is supported by a grant from the Natural Sciences and Engineering Research Council of Canada.

REFERENCES

- ¹ C.D. Goodman *et al.*, Phys. Rev. Lett. 44, 1755 (1980).
- ² C. Garde, Nucl. Phys. A396, 127c (1983).
- ³ I. Ikeda, Progress in Theoretical Physics 31, 258 (1964).
- ⁴ G.T. Bertsch and H. Esbensen, Argonne Nat. Lab., PHY-4803-TH-86, (1986).
- ⁵ J. Rapaport *et al.*, Nucl. Phys. A410, 371 (1983).
- ⁶ F. Ajzenberg-Selove *et al.*, Phys. Rev. C30, 1850 (1984).
- ⁷ S.D. Bloom and G.M. Fuller, Nucl. Phys. A440, 511 (1985).
- ⁸ K. Muto, Nucl. Phys. A451, 481 (1986).
- ⁹ MRS manual, TRIUMF.
- ¹⁰ R. Helmer, in "Isovector Excitations in Nuclei", TRIUMF, Vancouver, to be published in Can. J. Phys.
- ¹¹ R.S. Henderson *et al.*, Accepted by Nucl. Inst. and Methods.
- ¹² R.A. Arndt and L.D. Soper, Scattering Analysis Interactive Dial-in (SAID) program, unpublished.
- ¹³ program DWBA70, R. Schaeffer and J. Raynal (unpublished) extended version DW81 by J.R. Comfort (unpublished).
- ¹⁴ M.A. Franey and W.G. Love, Phys. Rev. C31, 488 (1985).
- ¹⁵ M.A. Moinester, in "Isovector Excitations in Nuclei", TRIUMF, Vancouver, to be published in Can. J. Phys.
- ¹⁶ O. Häusser, R.G. Jeppesen, *et al.*, unpublished.
- ¹⁷ OXBASH, (Nov. 1985 MSU version), B.A. Brown, A. Etchegoyen, W.D.M. Rae, MSUCL Report No. 524.
- ¹⁸ H. Wildenthal, Prog. in Part. and Nucl. Phys., 11, 5 (1984).
- ¹⁹ N. Auerbach *et al.*, Phys. Lett. 118B, 256 (1982).
- ²⁰ D. Cha, Phys. Rev. C27, 2269 (1983).
- ²¹ M.C. Vetterli *et al.*, to be published.

FIGURE CAPTIONS

Fig.1. Cross-sections for the $^{54}\text{Fe}(\text{n},\text{p})^{54}\text{Mn}$ reaction at 298 Mev for five angles between 0° and 12°. The strongly forward peaked feature below 10 MeV in excitation energy is the Gamow-Teller resonance. The dashed curve in the 0° spectrum is the calculation of Bloom and Fuller⁷ normalized by a factor of about 0.45.

Fig.2. Multipole decomposition of the 0°, $^{54}\text{Fe}(\text{n},\text{p})^{54}\text{Mn}$ spectrum. The top graph is a plot of the data with the total fitted curve, as well as the $\Delta L=0$ component shown as the hatched region. The bottom graph shows the $\Delta L=1$ component (cross-hatched region) and the $\Delta L=2$ component (vertically hatched region).




Fig. 2

Fig. 1