

EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH

Proposal to the ISOLDE and Neutron Time-of-Flight Committee

Investigating the key *rp* process reaction $^{61}\text{Ga}(p,\gamma)^{62}\text{Ge}$ reaction via $^{61}\text{Zn}(d,p)^{62}\text{Zn}$ transfer

January 11th 2017

G. Lotay¹, D.T. Doherty¹, W.N. Catford¹, Zs. Podolyak¹, P.A. Butler², R.D. Page², D.K. Sharp³, S.J. Freeman³, M. Labiche⁴, B.P. Kay⁵, C.R. Hoffman⁵, R.V.F. Janssens⁵, D.G. Jenkins⁶, N. Orr⁷, A. Matta⁷

¹*Department of Physics, University of Surrey, Guildford, Surrey, GU2 7XH. UK.*

²*Oliver Lodge Laboratory, University of Liverpool, Liverpool, L69 7ZE, UK.*

³*School of Physics and Astronomy, University of Manchester, Manchester, M13 9PL. UK.*

⁴*STFC Daresbury Laboratory, Daresbury, Warrington, WA4 4AD. UK.*

⁵*Physics Division, Argonne National Laboratory, Argonne, Illinois, 60439. USA.*

⁶*Department of Physics, University of York, Heslington, York, YO10 5DD. UK.*

⁷*LPC-ENSICAEN, IN2P3/CNRS et Universite de Caen, 1405 Caen, FRANCE.*

Spokesperson: G. Lotay (g.lotay@surrey.ac.uk)

Co-spokesperson: D.T. Doherty (d.t.doherty@surrey.ac.uk)

ISOLDE contact: Karl Johnston (karl.johnston@cern.ch)

Abstract: We propose to study the $^{61}\text{Zn}(d,p)^{62}\text{Zn}$ reaction in inverse kinematics for the first time, using the ISOL Solenoidal Spectrometer currently being installed at ISOLDE. This measurement represents the mirror analog of the astrophysically important $^{61}\text{Ga}(p,\gamma)^{62}\text{Ge}$ process (a reaction that cannot be presently studied with conventional means) and will allow for the first ever constraints to be placed on the stellar reaction rate. In particular, the energies and spectroscopic factors obtained for excited states in ^{62}Zn will be used to determine the resonant properties of proton-unbound levels in the nucleus ^{62}Ge , which are expected to dominate the $^{61}\text{Ga}(p,\gamma)^{62}\text{Ge}$ reaction in X-ray bursts. This study is very timely as the $^{61}\text{Ga}(p,\gamma)^{62}\text{Ge}$ reaction directly affects astronomical observables that are currently being obtained by the latest generation of space-based telescopes with unprecedented precision. Moreover, it complements the already approved proposals for the ISOL Solenoidal Spectrometer and extends the programme to studies relevant for nuclear astrophysics.

Requested shifts: 7 days (14 shifts)

Installation: ISOL Solenoidal Spectrometer



1 Physics Case

Type-I X-ray bursts are interpreted as thermonuclear explosions in the atmospheres of accreting neutron stars in close binary systems [1]. These astronomical scenarios exhibit brief recurrent bursts of intense X-ray emission and represent some of the most frequent and violent stellar events to occur in our Galaxy. Recently, space-borne satellites such as BeppoSAX and Chandra have produced a wealth of observational data on Type-I X-ray bursts, marking a new era in X-ray astronomy [2]. However, despite the vast quantities of observational data now available, many key questions about the exact nature of X-ray bursts remain, particularly with regards to the shape and structure of the observed light curves [3]. Consequently, in order to fully exploit the remarkable achievements of X-ray astronomy, similar advances in our understanding of the underlying nuclear physics processes governing nucleosynthesis and energy generation are required. In fact, as neutron stars represent some of the most extreme states of nuclear matter, a complete understanding of Type-I X-ray bursts is likely to have significant implications beyond the field of nuclear astrophysics.

In between bursts, energy is generated at a constant rate by the β -limited hot CNO cycles. However, during the burst, sufficiently high temperatures are achieved ($T_{\text{peak}} \sim 0.8 - 1.5$ GK) such that it is possible to “breakout” from the hot CNO cycles into a whole new set of thermonuclear reactions, known as the rp -process [4]. This process involves a series of rapid proton captures resulting in the synthesis of very proton-rich nuclei up to the Sn – Te mass region and could be a candidate for the production of the astrophysically important p -nuclei such as ^{92}Mo and ^{96}Ru , although the exact escape mechanism is not clear. Recently, significant increases in computing power have allowed for detailed theoretical models of X-ray burst nucleosynthesis to be used to estimate the impact of individual reaction rate uncertainties on both final isotopic abundance yields and overall energy output [5,6]. In particular, a study by Cyburt *et al.* [6] investigated the dependence of X-ray bursts on uncertainties in (p,γ) , (α,γ) , and (α,p) nuclear reaction rates using fully self-consistent models that account for feedbacks between changes in nuclear energy generation and changes in astrophysical conditions. Rather surprisingly, despite varying 1931 different nuclear processes through their associated uncertainties, only a handful of reactions were found to have a significant affect on both the burst light curve and final isotopic compositions [6]. Of these, the $^{61}\text{Ga}(p,\gamma)^{62}\text{Ge}$ reaction was highlighted as being particularly significant (Figs. 1 and 2) and at present, its rate over the temperature range of X-ray bursts is effectively unknown.




Figure 1 - Effect of varying the $^{61}\text{Ga}(p,\gamma)$ reaction rate on the observed light curve [6].




Figure 2 - Effect of varying the $^{61}\text{Ga}(p,\gamma)$ reaction rate on final isotopic abundances [6].

The astrophysical ${}^{61}\text{Ga}(p,\gamma){}^{62}\text{Ge}$ reaction, like many reactions that occur in the rp process, is expected to be dominated by resonant capture to excited states in ${}^{62}\text{Ge}$ that exist above the proton-emission threshold. However, studying such resonances in systems that lie far from stability is extremely difficult. In fact, almost no experimental information exists for excited states in the nucleus ${}^{62}\text{Ge}$.

A direct investigation of resonances in the ${}^{61}\text{Ga}(p,\gamma){}^{62}\text{Ge}$ reaction is presently unfeasible as sufficiently intense radioactive beams of ${}^{61}\text{Ga}$ are not available. That being said, it has recently been shown that a possible solution to this problem may be achieved by obtaining the nuclear properties of analog states in the mirror system, which in turn may be used to determine the critical unknown properties of resonances in the proton-rich system through isospin symmetry [7,8]. Consequently, we propose to utilise the intense radioactive beams of ${}^{61}\text{Zn}$ available at CERN as well as the newly installed ISOL Solenoid Spectrometer to obtain spectroscopic information on excited states in the astrophysically important mirror nucleus ${}^{62}\text{Zn}$, via ${}^{61}\text{Zn}(d,p)$ transfer. This information will then be used to place the first ever constraints on the key ${}^{61}\text{Ga}(p,\gamma){}^{62}\text{Ge}$ reaction rate in X-ray burster environments, thereby allowing a detailed comparison between the latest theoretical models and current astronomical data.

2 Experimental details

We propose to measure the ${}^{61}\text{Zn}(d,p)$ reaction in inverse kinematics using a radioactive beam of ${}^{61}\text{Zn}$ at 7.5 MeV/u to bombard a $\sim 100 \mu\text{g}/\text{cm}^2$ thick CD_2 target, in order to probe excited states in the nucleus ${}^{62}\text{Zn}$. At these energies, the angular distributions for transfer to final states of differing l are more pronounced and forward peaked, compared to lower energy measurements, such that assignments of the transferred angular momentum are more distinct. In particular, we aim to measure the neutron spectroscopic factors of low- l transfer levels in the excitation energy region of 2 – 3.5 MeV. Such states, for which the cross section peaks at forward centre-of-mass angles, represent analogs of resonances in the ${}^{61}\text{Ga}(p,\gamma){}^{62}\text{Ge}$ reaction and as such, by determining their properties, it is possible to evaluate the stellar reaction rate.

The resulting protons following ${}^{61}\text{Zn}(d,p)$ transfer will be detected using the new ISOL Solenoidal Spectrometer (ISS). In analogy with two recently approved proposals [9,10], the ISS will be operated in a manner similar to the HELIOS Spectrometer at Argonne National Laboratory (ANL) [11,12]. Specifically, the same position-sensitive silicon array and associated electronics from ANL will be used. Members of this collaboration have extensive experience using these detectors. The extracted cross sections and angular distributions will be compared to calculations using the ADWA code TWOFNR [13], to obtain information on the l of the final states and spectroscopic factors. Based on recent simulations for both the ${}^{28}\text{Mg}(d,p)$ and ${}^{206}\text{Hg}(d,p)$ reactions using the ISS [9,10], we estimate an excitation energy resolution of <100 keV. If we consider the ${}^{61}\text{Ni}(d,p)$ reaction [14] as a guide (Fig. 3), we find that the level density up to 3.5 MeV excitation is relatively low. Consequently, it is expected that an excitation energy resolution of ~ 100 keV will be more than sufficient to easily separate the strong spectroscopic factor states of most astrophysical importance.




Figure 3 - Proton spectrum obtained from a $^{61}\text{Ni}(d,p)^{62}\text{Ni}$ transfer reaction study [14].

The experimental configuration for the $^{61}\text{Zn}(d,p)$ measurement is shown in Fig. 4. The silicon array will be positioned -45 cm from the target as measured to the nearest detector edge, covering a range in z from the target of -45 to -80 cm. The solenoid field will be set at 2.5 T. With these settings, protons emitted at $10^\circ < \theta_{\text{cm}} < 30^\circ$ will be incident on the array for all states up to 3.5 MeV. This range of angles covers the maxima for astrophysically important low- l transfers. Elastically-scattered deuterons will be detected in an annular silicon detector positioned at $z = +17$ cm. The proposed measurement does not require recoil detection, in analogy with Kay *et al.* [10], and as such, a smooth background of protons, and to a lesser extent alpha particles, from fusion-evaporation reactions will be present at backwards angles.




Figure 4 - Proposed experimental setup within the solenoidal spectrometer. Distances are relative to the target position. For the (d,p) reaction, the black lines represent population of the ground state, while the red lines indicate population of an excited state at 3.5 MeV.

3 Beam time request

Beam intensities of up to 4×10^6 pps have been reported for ^{61}Zn on the ISOLDE yield database. Consequently, we assume that a beam intensity of $\sim 4 \times 10^5$ pps of ^{61}Zn should be reasonably achievable at HIE-ISOLDE. From discussions with beam experts at CERN, it is expected that RILIS ionized neutron-deficient Zn beams will be largely free from contamination. In particular, Ga (which has previously been observed as a contaminant in running neutron-rich Zn beams at CERN) is expected to be suppressed by more than 7 orders of magnitude in comparison to Zn for the proposed study. That being said, we would still intend to run with and without RILIS to establish any potential background contamination peaks.

The array has an efficiency of 50% in the azimuthal angle and 85% in the theta angle. Protons in the angular range $10^\circ < \theta_{\text{cm}} < 30^\circ$ will be incident on the array. Using cross sections estimated using the ADWA code TWOFNR, a CD_2 target $\sim 100 \mu\text{g}/\text{cm}^2$ thick and assuming a spectroscopic factor of ~ 0.5 for excited states in the energy region 2 – 3.5 MeV, we estimate that ~ 220 counts per day will be observed in the whole array for the states of interest. On average, there will be ~ 44 counts in each ring of detectors per day, where each ring essentially corresponds to an angular bin. To obtain a spectroscopic factor of the level 0.5 to a precision of $\sim 20\%$, we would require 10 shifts of beam on target. In order to establish background, we would intend to run without RILIS for 3 shifts.

The present proposal links exceptionally well with the two currently approved proposals for the ISOL Solenoidal Spectrometer and highlights the capability of the device for measurements of astrophysical importance.

Summary of requested shifts:

14 shifts of protons are requested for this measurement. This will be split in three, with 10 shifts running with RILIS, 3 shifts running without RILIS and 1 shift for beam optimisation.

References:

- [1] H. Schatz and K.E. Rehm, Nucl. Phys. A **777**, 601 (2006).
- [2] A. Paizis *et al.*, Astrophys. J. **755**, 52 (2012).
- [3] J. L. Fisker *et al.*, Astrophys. J. **608**, L61-L64 (2004).
- [4] H. Schatz *et al.*, Phys. Rev. Lett. **86**, 3471 (2001).
- [5] A. Parikh, J. Jose, F. Moreno and C. Iliadis, Astrophys. J. Suppl. Ser. **178**, 110 (2008).
- [6] R.H. Cyburt *et al.*, Astrophys. J. **830**, 2 (2016).
- [7] V. Margerin, G. Lotay *et al.*, Phys. Rev. Lett. **115**, 062701 (2015).
- [8] S.D. Pain *et al.*, Phys. Rev. Lett. **114**, 212501 (2015).
- [9] D.K. Sharp *et al.*, INTC P-470 (2016).
- [10] B.P. Kay *et al.*, INTC P-467 (2016).
- [11] A. H. Wuosmaa, et al., Nucl. Instrum. Meth. A, 580, 1290 (2007).
- [12] J. C. Lighthall, et al., Nucl. Instrum. Meth. A, 622, 97 (2010).
- [13] J. A. Tostevin, University of Surrey version of the code TWOFNR (of M. Toyama, M. Igarashi and N. Kishida) and code FRONT (private communication).
- [14] O. Karban *et al.*, Nucl. Phys. **A366**, 68 (1981).

Appendix

DESCRIPTION OF THE PROPOSED EXPERIMENT

The experimental setup comprises: *(name the fixed-ISOLDE installations, as well as flexible elements of the experiment)*

Part of the Choose an item.	Availability	Design and manufacturing
[if relevant, name fixed ISOLDE installation: MINIBALL + only CD, MINIBALL + T-REX]	<input type="checkbox"/> Existing	<input type="checkbox"/> To be used without any modification
ISOL Solenoidal Spectrometer	<input type="checkbox"/> Existing <input checked="" type="checkbox"/> New	<input type="checkbox"/> To be used without any modification <input type="checkbox"/> To be modified <input type="checkbox"/> Standard equipment supplied by a manufacturer <input checked="" type="checkbox"/> CERN/collaboration responsible for the design and/or manufacturing

HAZARDS GENERATED BY THE EXPERIMENT

(if using fixed installation) Hazards named in the document relevant for the fixed [MINIBALL + only CD, MINIBALL + T-REX] installation.

Additional hazards:

Hazards			
Thermodynamic and fluidic			
Pressure			
Vacuum			
Temperature	4 K		
Heat transfer			
Thermal properties of materials			
Cryogenic fluid	LHe, ~1650 l, LN ₂ , ~200 l, 1.0 Bar		
Electrical and electromagnetic			
Electricity	0 V, 300 A		
Static electricity			
Magnetic field	2.5 T		
Batteries	<input type="checkbox"/>		
Capacitors	<input type="checkbox"/>		
Ionizing radiation			
Target material	Deuterated Polyethylene [CD ₂]		
Beam particle type (e, p, ions, etc)	⁶¹ Zn		
Beam intensity	4 x 10 ⁵ pps		
Beam energy	7.5 MeV/u		
Cooling liquids			
Gases			
Calibration sources:	<input checked="" type="checkbox"/>		
• Open source	<input checked="" type="checkbox"/> (alpha calibrations source)		

• Sealed source	<input type="checkbox"/> [ISO standard]		
• Isotope			
• Activity			
Use of activated material:			
• Description	<input type="checkbox"/>		
• Dose rate on contact and in 10 cm distance	[dose][mSV]		
• Isotope			
• Activity			
Non-ionizing radiation			
Laser			
UV light			
Microwaves (300MHz-30 GHz)			
Radiofrequency (1-300MHz)			
Chemical			
Toxic			
Harmful			
CMR (carcinogens, mutagens and substances toxic to reproduction)			
Corrosive			
Irritant			
Flammable			
Oxidizing			
Explosiveness			
Asphyxiant	Helium		
Dangerous for the environment			
Mechanical			
Physical impact or mechanical energy (moving parts)			
Mechanical properties (Sharp, rough, slippery)			
Vibration			
Vehicles and Means of Transport			
Noise			
Frequency			
Intensity			
Physical			
Confined spaces			
High workplaces			
Access to high workplaces			
Obstructions in passageways			
Manual handling			
Poor ergonomics			

0.1 Hazard identification

Average electrical power requirements (excluding fixed ISOLDE-installation mentioned above): 5 kW.