

Mixing and CP violation results in b -hadron decays at LHCb

XUESONG LIU

*On behalf of the LHCb Collaboration,
Center for High Energy Physics,
Tsinghua University, Beijing, 100084, China*

ABSTRACT

Measurements of mixing and CP violation in b -hadron decays are great probes to search for physics beyond the Standard Model. A selection of recent results based upon 3.0 fb^{-1} of LHCb data are presented. These are the first measurement of the CP violating phase ϕ_s using $B_s^0 \rightarrow J/\psi K^+ K^-$ in the mass region above the $\phi(1020)$ resonance and $B_s^0 \rightarrow \psi(2S)\phi$ decays, the determination of B_s^0 lifetime using $B_s^0 \rightarrow D^{(*)-} \mu^+ \nu_\mu$ decays, the updated measurements of the CKM angle γ with $B^\pm \rightarrow DK^{*\pm}$ and $B_s^0 \rightarrow D_s^\mp K^\pm$ decay modes and the CP violation searches in $\Lambda_b^0 \rightarrow p\pi^-\pi^+\pi^-$ and $\Lambda_b^0 \rightarrow pK^-\mu^+\mu^-$ decays.

PRESENTED AT

The Fifth Annual Conference
on Large Hadron Collider Physics
Shanghai Jiao Tong University, Shanghai, China
May 15-20, 2017

1 Measurements of the CP violating phase ϕ_s

The interference of the decay amplitudes and the $B_s^0\text{-}\bar{B}_s^0$ mixing amplitudes can give rise to a CP violating phase

$$\phi_s = \phi_M - 2\phi_D, \quad (1)$$

where ϕ_M is the $B_s^0\text{-}\bar{B}_s^0$ mixing phase and ϕ_D is the weak phase between the B_s^0 and \bar{B}_s^0 decay amplitudes. Measurement of the ϕ_s is obtained through the time-dependent CP asymmetry

$$A_{CP}(t) = \frac{\Gamma(\bar{B}_s^0(t) - f) - \Gamma(B_s^0(t) - f)}{\Gamma(\bar{B}_s^0(t) - f) + \Gamma(B_s^0(t) - f)} = \frac{C_f \cos(\Delta mt) - S_f \sin(\Delta mt)}{\cosh(\Delta\Gamma t/2) + A^{\Delta\Gamma} \sinh(\Gamma t/2)}, \quad (2)$$

where C_f is related to CP violation in the decay and S_f represents CP violation in the interference. Neglecting penguin pollution and assuming no Beyond Standard Model (BSM) contribution, the phase ϕ_s is predicted to be $\phi_s = -2\beta_s = 0.0365_{-0.0012}^{+0.0013}$ rad [1]. The world average result by HFLAV, $\phi_s = -0.021 \pm 0.031$ rad [2], is dominated by LHCb measurements and is compatible with Standard Model (SM) prediction.

Measurement of ϕ_s with $B_s^0 \rightarrow J/\psi K^+ K^-$ in the mass region above the $\phi(1020)$

A study performed by LHCb, using a data sample corresponding to 1 fb^{-1} shows a rich resonance spectrum in the $K^+ K^-$ mass distribution [3], indicating that the significant contribution from the $f_2'(1525)$ resonance and nonresonant S-wave amplitudes can be used for further studies of CP violation [4]. In this analysis, the CP observables are determined as functions of the B_s^0 decay time in a four-dimensional phase space including the three helicity angles characterizing the decay and the invariant mass of the two kaons, m_{KK} . With respect to the previous publication focusing on the $J/\psi \phi(1020)$ decays [5], the $B_s^0 \rightarrow J/\psi K^+ K^-$ sample is separated into two regions, having m_{KK} above or below the $\phi(1020)$ resonance. A simultaneous fit is performed for the two samples.

All the resonances found in the previous analysis are considered as contribution to the entire fit, except for the unconfirmed $f_2(1640)$ state. The fit projection of m_{KK} is shown in Figure 1 [6]. The results are




Figure 1: Fit projection of m_{KK} . The resonance $\phi(1020)$, $f_2'(1525)$, $\phi(1680)$ and S-wave components are shown by magenta, brown, cyan and green curves; the other f_2 resonances are described by black solid curves.

determined as

$$\begin{aligned} \phi_s &= 0.119 \pm 0.107 \pm 0.034 \text{ rad}, \\ \Delta\Gamma_s &= 0.066 \pm 0.018 \pm 0.010 \text{ ps}^{-1}, \\ m_{f'(1525)} &= 1522.2 \pm 1.3 \pm 1.1, \text{ MeV}, \\ \Gamma_{f'(1525)} &= 78.0 \pm 3.0 \pm 3.7, \text{ MeV}, \end{aligned}$$

Parameter	Value
Γ_s [ps ⁻¹]	$0.668 \pm 0.011 \pm 0.006$
$\Delta\Gamma_s$ [ps ⁻¹]	$0.066^{+0.041}_{-0.044} \pm 0.007$
$ A_\perp ^2$	$0.264^{+0.024}_{-0.023} \pm 0.002$
$ A_0 ^2$	$0.422 \pm 0.014 \pm 0.003$
δ_\parallel [rad]	$3.67^{+0.13}_{-0.18} \pm 0.03$
δ_\perp [rad]	$3.29^{+0.43}_{-0.39} \pm 0.04$
ϕ_s [rad]	$0.23^{+0.29}_{-0.28} \pm 0.02$
$ \lambda $	$1.045^{+0.069}_{-0.050} \pm 0.007$
F_S	$0.061^{+0.026}_{-0.025} \pm 0.007$
δ_S [rad]	$0.03 \pm 0.14 \pm 0.02$

Table 1: Results of the maximum likelihood fit to the selected $B_s^0 \rightarrow \psi(2S)\phi$ candidates. The parameters are defined in Ref. [8]. The first uncertainty is statistical and the second is systematic.

where $\Delta\Gamma_s$ is the decay width difference between the light and the heavy mass eigenstates, $m_{f'(1525)}$ and $\Gamma_{f'(1525)}$ are the mass and width of $f'(1525)$, the first uncertainty is statistical and the second is systematic.

The combination with previous results from B_s^0 decays in the $\phi(1020)$ region [5] and $J/\psi\pi^+\pi^-$ final states [7] gives $\phi_s = 0.001 \pm 0.037$ rad [6].

Measurement of ϕ_s with $B_s^0 \rightarrow \psi(2S)\phi$

The $\psi(2S)$ resonance state is for the first time used in the ϕ_s measurement with $B_s^0 \rightarrow \phi(2S)(\mu^+\mu^-)\phi(K^+K^-)$ decay mode. This decay has a similar topology compared with the ‘golden mode’ $B_s^0 \rightarrow J/\psi(\mu^+\mu^-)\phi(K^+K^-)$ [5]. Compared to the latter analysis, a multivariate selection is added to optimise the statistical sensitivity and the control mode $B_s^0 \rightarrow \phi(2S)K^*$ is used to calibrate the decay time acceptance. In total, 4697 ± 71 signal candidates are found in this channel. An unbinned maximum likelihood fit to the four-dimensional distributions of the $B_s^0 \rightarrow \psi(2S)\phi$ decay time and helicity angles is done to estimate the CP violation phase ϕ_s . The results, listed in Table 1 [8], are compatible with previous LHCb measurements [5, 7].

2 Flavour-specific B_s^0 lifetime measurement with $B_s^0 \rightarrow D_s^{(*)-}\mu^+\nu_\mu$

The flavour-specific lifetime is defined as

$$\tau_s^{\text{fs}} = \frac{1}{\Gamma_s} \left[\frac{1 + (\Delta\Gamma_s/2\Gamma_s)^2}{1 - (\Delta\Gamma_s/2\Gamma_s)^2} \right], \quad (3)$$

where $\Delta\Gamma_s = \Gamma_{s,H} - \Gamma_{s,L}$ is the decay width difference between the light and the heavy mass eigenstates and $\Gamma_s = (\Gamma_{s,H} + \Gamma_{s,L})/2$ is the average width.

LHCb has reported a measurement of $\tau_s^{\text{fs}} = 1.535 \pm 0.015 \pm 0.014$ ps [10] using $B_s^0 \rightarrow D_s^- \pi^+$ decays. Semileptonic B_s^0 decays have a larger signal yield than in hadronic decays, such that can offer richer potential for precise measurement. In this analysis, the B_s^0 candidates are reconstructed with $D_s^{(*)-} \rightarrow K^+K^-\pi^-$ and muon candidates. The B_s^0 yields are determined with the ‘corrected-mass’, defined as $m_{\text{corr}} = p_{\perp,D\mu} + \sqrt{m_{D\mu}^2 + p_{\perp,D\mu}^2}$. The lifetime is determined from the variation in the B_s^0 signal yield as a function of decay time, relative to that of B^0 decays that are reconstructed in the same final state. The result is $\tau_s^{\text{fs}} = 1.547 \pm 0.013(\text{stat}) \pm 0.010(\text{syst}) \pm 0.004(\tau_B)$ ps [9] and is consistent with and more precise than the current measured values [10, 11].

3 Measurements of the CKM angle γ

The angle γ is the least well known angle in the CKM unitarity triangle. It can be measured using tree-level processes based on the interference between $b \rightarrow c\bar{u}s$ and $b \rightarrow u\bar{c}s$ transitions and has negligible theoretical uncertainty [12] in SM predictions. Disagreement between such direct measurement of γ and the value inferred from global CKM fits would indicate new physics beyond the SM. The world-average value of the γ measurements with $(B \rightarrow DK)$ -like decays is $\gamma = (74.0^{+5.8}_{-6.4})^\circ$ [2]. The CKMFitter group quoted $\gamma = (66.9^{+0.9}_{-3.4})^\circ$ [13] from a global fit of CKM parameters. And the latest LHCb combination yields $\gamma = (76.8^{+5.1}_{-5.7})^\circ$ [14, 15]. Exploiting the Run-II dataset with already used or even new decay modes will allow to reduce the uncertainty on the measured value of γ .

Measurement of the CP observables using $B^\pm \rightarrow DK^{*\pm}$

The CP violation in $B^\pm \rightarrow DK^{*\pm}$ decays is sensitive to γ at tree level. Decays of the D mesons to $K^-\pi^+$, K^+K^- , $\pi^+\pi^-$, $K^+\pi^-$, $K^-\pi^+\pi^-\pi^+$, $\pi^+\pi^-\pi^+\pi^-$, $K^+\pi^-\pi^+\pi^-$ and the $K^*(892)^\pm$ with the $K_s^0\pi^\pm$ final states are used in the reconstruction. LHCb has reported the measurement of γ with $B^\pm \rightarrow DK^\pm$ decays modes [16]. Compared with these channels, the branching fractions of $B^\pm \rightarrow DK^{*\pm}$ is of a similar magnitude, however the reconstruction efficiencies for the $K^{*\pm}$ decay are much lower.

Several CP asymmetries and ratios, corresponding to different final state of the D^0 , were measured as the γ sensitive observables. The mass spectrum of the $B^\pm \rightarrow D(K^+K^-)K^{*\mp}$ decay modes are shown in Figure 2 [17]. A simultaneous fit is performed to all D final state.

The measured CP observable in the favoured mode, $A_{K\pi} = -0.004 \pm 0.023(\text{stat}) \pm 0.008(\text{syst})$ [17], is consistent with zero. The combined results from the CP -eigenstate K^+K^- and $\pi^+\pi^-$ decay modes, $R_{CP+} = 1.18 \pm 0.08(\text{stat}) \pm 0.01(\text{syst})$, $A_{CP+} = 0.08 \pm 0.06(\text{stat}) \pm 0.01(\text{syst})$ [17], are consistent with and more precise than the previous measurements from BaBar [18].




Figure 2: Mass distribution of the $B^\pm \rightarrow D(K^+K^-)K^{*\mp}$ modes using Run 1 and Run 2 data. The signal component is represented by the red shaded area, the combinatorial background by the dotted blue line and the partially reconstructed background by the solid green line.

Measurement of the CKM angle γ using $B_s^0 \rightarrow D_s^\mp K^\pm$

Another approach to extract γ is through the neutral B meson decays and perform a time-dependent measurement of the CP observables. In $B_s^0 \rightarrow D_s^\mp K^\pm$ decays, the sensitivity to the weak phase $\gamma - 2\beta_s$ arises from the interference of B_s^0 and \bar{B}_s^0 decaying into the same final state: $D_s^+K^-$ or $D_s^-K^+$. The CP observables can be interpreted in terms of either γ or β_s by using an independent measurement of the other parameters as input and are given by

$$C_f = \frac{1-r_{D_s K}^2}{1+r_{D_s K}^2}, \quad A_f^{\Delta\Gamma} = \frac{-2r_{D_s K} \cos(\delta - (\gamma - 2\beta_s))}{1+r_{D_s K}^2}, \quad \bar{A}_f^{\Delta\Gamma} = \frac{-2r_{D_s K} \cos(\delta + (\gamma - 2\beta_s))}{1+r_{D_s K}^2}, \quad (4)$$

$$S_f = \frac{2r_{D_s K} \sin(\delta - (\gamma - 2\beta_s))}{1+r_{D_s K}^2}, \quad \bar{S}_f = \frac{-2r_{D_s K} \sin(\delta + (\gamma - 2\beta_s))}{1+r_{D_s K}^2},$$

Parameter	Value
C_f	$0.735 \pm 0.142 \pm 0.048$
$A_f^{\Delta\Gamma}$	$0.395 \pm 0.277 \pm 0.122$
$A_{\bar{f}}^{\Delta\Gamma}$	$0.314 \pm 0.274 \pm 0.107$
S_f	$-0.518 \pm 0.202 \pm 0.073$
$S_{\bar{f}}$	$-0.496 \pm 0.197 \pm 0.071$

Table 2: Results of the maximum likelihood fit to the selected $B_s^0 \rightarrow D_s^\mp K^\pm$ candidates. The first uncertainty is statistical and the second is systematic.

where $r_{D_s K} \equiv |\lambda_{D_s K}| = |A(\overline{B}_s^0 \rightarrow D_s^- K^+)/A(B_s^0 \rightarrow D_s^- K^+)|$ is the magnitude of the amplitude ratio, δ is the strong phase difference between the favoured and suppressed amplitudes. The initial flavour of the B_s^0 mesons is determined with ‘flavour tagging’ process. The kinematically similar mode $B_s^0 \rightarrow D_s^- \pi^+$ is used as a control channel for the determination of the decay time efficiency and flavour tagging performance. The fitted CP observables are listed in Table 2. With the value of β_s from previous LHCb measurement in $B_s^0 \rightarrow J/\psi K^+ K^-$ [5] and $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$ [7], the angle γ is determined as $\gamma = (127_{-22}^{+17})^\circ$.

4 CP violation searches in the b -baryons decays

In the SM, decays of the Λ_b^0 baryon are predicted to have non-negligible CP asymmetries, as large as 20% for certain three-body decay modes [19]. Large numbers of beauty baryons are produced at the LHC, the production ratio of $B^0 : \Lambda_b^0 : B_s^0$ particles is approximately 4 : 2 : 1 within the LHCb detector acceptance [20], which allows to perform sensitive studies of CP violation in b -baryon sector.

CP violation searches in $\Lambda_b^0 \rightarrow p\pi^-\pi^+\pi^-$

The $\Lambda_b^0 \rightarrow p\pi^-\pi^+\pi^-$ decay is mediated by the weak interaction, CP violation in decay can arise from the interference of a tree amplitude and a penguin amplitude.

The scalar triple products of final-state particle momenta in the Λ_b^0 centre-of-mass frame are used for P and CP violating searches, defined as $C_{\hat{T}} = \vec{p}_p \cdot (\vec{p}_{\pi_{\text{slow}}} \times \vec{p}_{\pi^+}) \propto \sin\Phi$ for Λ_b^0 , and $\overline{C}_{\hat{T}} = \vec{p}_p \cdot (\vec{p}_{\pi_{\text{slow}}} \times \vec{p}_{\pi^-}) \propto \sin\overline{\Phi}$ for $\overline{\Lambda}_b^0$, where Φ is the angle between the decay plane of the proton and higher-momentum π^- and that of the π^+ and the lower-momentum π^- . Two \hat{T} -odd asymmetries are built as

$$A_{\hat{T}} = \frac{N_{\Lambda_b^0}(C_{\hat{T}} > 0) - N_{\Lambda_b^0}(C_{\hat{T}} < 0)}{N_{\Lambda_b^0}(C_{\hat{T}} > 0) + N_{\Lambda_b^0}(C_{\hat{T}} < 0)}, \quad \overline{A}_{\hat{T}} = \frac{N_{\overline{\Lambda}_b^0}(-\overline{C}_{\hat{T}} > 0) - N_{\overline{\Lambda}_b^0}(-\overline{C}_{\hat{T}} < 0)}{N_{\overline{\Lambda}_b^0}(-\overline{C}_{\hat{T}} > 0) + N_{\overline{\Lambda}_b^0}(-\overline{C}_{\hat{T}} < 0)}. \quad (5)$$

The P and CP violating observables can be constructed as

$$a_P^{\hat{T}\text{-odd}} = \frac{1}{2}(A_{\hat{T}} + \overline{A}_{\hat{T}}), \quad a_{CP}^{\hat{T}\text{-odd}} = \frac{1}{2}(A_{\hat{T}} - \overline{A}_{\hat{T}}). \quad (6)$$

The analysis was performed in ten bins of $|\Phi|$ and in twelve bins of the phase space. The distribution of the asymmetries are shown in Figure 3. Combining the results of both schemes gives evidence for CP violation in $\Lambda_b^0 \rightarrow p\pi^-\pi^+\pi^-$ at the level of 3.3σ [21].

CP violation searches in $\Lambda_b^0 \rightarrow pK^-\mu^+\mu^-$

The $\Lambda_b^0 \rightarrow pK^-\mu^+\mu^-$ is mediated by flavour-changing neutral-current transitions in the SM. The theoretical prediction of CP violation in this decay is very small [22, 23] which makes it particularly sensitive to CP




Figure 3: The asymmetries $a_P^{\hat{T}-odd}$ (upper panels) and $a_{CP}^{\hat{T}-odd}$ (lower panels) for $\Lambda_b^0 \rightarrow p\pi^-\pi^+\pi^-$ decays in bins of phase space and Φ .

violation contributions from BSM physics. The signal yield is determined to be 600 ± 44 . The asymmetry A_{CP} is related to the raw asymmetry A_{raw} via

$$A_{raw} \equiv \frac{N(\Lambda_b^0 \rightarrow pK^-\mu^+\mu^-) - N(\bar{\Lambda}_b^0 \rightarrow \bar{p}K^+\mu^-\mu^+)}{N(\Lambda_b^0 \rightarrow pK^-\mu^+\mu^-) + N(\bar{\Lambda}_b^0 \rightarrow \bar{p}K^+\mu^-\mu^+)} \approx A_{CP}(\Lambda_b^0 \rightarrow pK^-\mu^+\mu^-) + A_{\text{prob}}(\Lambda_b^0) - A_{\text{reco}}(K^+) + A_{\text{reco}}(p), \quad (7)$$

where $A_{\text{prob}}(\Lambda_b^0)$ is the Λ_b^0 production asymmetry, $A_{\text{reco}}(K^+)$ and $A_{\text{reco}}(p)$ are the reconstruction asymmetries for kaons and protons. Measurement of the difference of raw asymmetries between the signal and $\Lambda_b^0 \rightarrow pK^-J/\psi$ control mode, can cancel the effects from the production and reconstruction asymmetries. Thus two different CP violating observables that are sensitive to different manifestations of CP violation, ΔA_{CP} and $a_{CP}^{\hat{T}-odd}$, are measured. The CP -odd observable is defined following the same method in the above $\Lambda_b^0 \rightarrow p\pi^-\pi^+\pi^-$ analysis. The results, $\Delta A_{CP} = (-3.5 \pm 5.0(\text{stat}) \pm 0.2(\text{syst})) \times 10^{-2}$, $a_{CP}^{\hat{T}-odd} = (1.2 \pm 5.0(\text{stat}) \pm 0.7(\text{syst})) \times 10^{-2}$ [24], are compatible with CP and parity conservation and agree with SM predictions.

5 Conclusions

Many new measurements of CP violation searches in b -hadron decays are performed by the LHCb experiment. These results improve our knowledge on the CKM parameters. All the measurements are still statistically limited and can be improved with LHC Run-II data.

References

- [1] J. Charles *et al.*, Phys. Rev. **D91** (2015) 7 073007, [arXiv:1501.05013[hep-ex]].
- [2] Y. Amhis *et al.* [Heavy Flavor Averaging Group], [arXiv:1612.07233[hep-ex]].
- [3] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. **D87** (2013) 7 072004, [arXiv:1302.1213[hep-ex]].
- [4] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. Lett. **108** (2012) 151801, [arXiv:1112.4695[hep-ex]].
- [5] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. Lett. **114** (2015) 4 041801, [arXiv:1411.3104[hep-ex]].

- [6] R. Aaij *et al.* [LHCb Collaboration], JHEP **08** (2017) 037, [arXiv:1704.08217[hep-ex]].
- [7] R. Aaij *et al.* [LHCb Collaboration], Phys. Lett. **B736** (2014) 186-195, [arXiv:1405.4140[hep-ex]].
- [8] R. Aaij *et al.* [LHCb Collaboration], Phys. Lett., **B762** (2016) 253-262, [arXiv:1608.04855[hep-ex]].
- [9] R. Aaij *et al.* [LHCb Collaboration], submitted to Phys. Rev. Lett. [arXiv:1705.03475[hep-ex]].
- [10] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. Lett. **113** (2014) 17 172001, [arXiv:1407.5873[hep-ex]].
- [11] V. Abazov *et al.* [D0 Collaboration], Phys. Rev. Lett. **114** (2015) 6 062001, [arXiv:1410.1568[hep-ex]].
- [12] B. Joachim and Z. Jure, JHEP **01** (2014) 051, [arXiv:1308.5663 [hep-ph]].
- [13] J. Charles *et al.* [CKMfitter Group], Eur. Phys. J. C **41** (2005) 1 1-131, [arXiv:hep-ph/0406184], updated results and plots available at: <http://ckmfitter.in2p3.fr>.
- [14] R. Aaij *et al.* [LHCb Collaboration], JHEP **12** (2016) 087, [arXiv:1611.03076[hep-ex]].
- [15] R. Aaij *et al.* [LHCb Collaboration], LHCb-CONF-2017-004.
- [16] R. Aaij *et al.* [LHCb Collaboration], Phys. Lett. **B760** (2016) 117-131, [arXiv:1603.08993[hep-ex]].
- [17] R. Aaij *et al.* [LHCb Collaboration], LHCb-PAPER-2017-030.
- [18] B. Aubert *et al.* [BaBar Collaboration], Phys. Rev. **D80** (2009) 092001.
- [19] Y.K. Hsiao and C.Q. Geng, Phys. Rev. **D91** (2015) 11 116007, [arXiv:1412.1899[hep-ex]].
- [20] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. **D85** (2012) 032008, [arXiv:1111.2357[hep-ex]].
- [21] R. Aaij *et al.* [LHCb Collaboration], Nature Phys. **13** (2017) 391-396, [arXiv:1609.05216[hep-ex]].
- [22] M.A. Paracha *et al.*, PTEP **2015** (2015) 3 033B04, [arXiv:1408.4318[hep-ph]].
- [23] A.K. Alok *et al.*, JHEP **11** (2011) 122, [arXiv:1103.5344[hep-ph]].
- [24] R. Aaij *et al.* [LHCb Collaboration], JHEP **06** (2017) 108, [arXiv:1703.00256[hep-ex]].