

CC

CBPF - NF 93-065
su 9419



CBPF-CENTRO BRASILEIRO DE PESQUISAS FÍSICAS

Notas de Física

CBPF-NF-065/93

*Highly Deformed q -Oscillator
Systems*

by

*M.R-Monteiro, I. Roditi and
L.M.C.S. Rodrigues*

*Rio de Janeiro
1993*

NOTAS DE FÍSICA é uma pré-publicação de trabalho original em Física

NOTAS DE FÍSICA is a preprint of original works unpublished in Physics

Pedidos de cópias desta publicação devem ser enviados aos autores ou ã:

Requests for copies of these reports should be addressed to:

Centro Brasileiro de Pesquisas Físicas
Área de Publicações
Rua Dr. Xavier Sigaud, 150 - 4º andar
22.290 - Rio de Janeiro, RJ
BRASIL

CBPF-NF-065/93

*Highly Deformed q -Oscillator
Systems*

by

*M.R-Monteiro, I. Roditi and
L.M.C.S. Rodrigues*

Centro Brasileiro de Pesquisas Físicas — CBPF/CNPq
Rua Dr. Xavier Sigaud, 150
22290-180 – Rio de Janeiro, RJ – Brasil

Abstract

We consider the large q limit of systems made of deformed Heisenberg operators. When the deformation parameter is infinite the Fock space and the statistical properties have a fermionic behaviour. We also investigate the ideal q -gas and find the virial expansion of its equation of state.

Key-words: Quantum algebras; Statistical mechanics.

1 Introduction

There has been a great interest in Quantum Groups [1-4] in the last years, both from physicists and mathematicians. This mathematical structure, also called Quasitriangular Hopf algebras, has emerged as an appealing non-trivial generalization of Lie algebras and groups which are recovered when the deformation parameter (or a set of parameters) goes to one.

Quantum Groups have left their trace in several areas of physics [5-8] and deformed Heisenberg algebras [9] have been attracting increasing interest mainly due to the role played by Heisenberg algebras in a wide range of problems. Recently, the connection of q -oscillators with quantum algebras was investigated [10] thus permitting the discussion of the thermal properties of q -oscillators systems [11-15] and the analysis of possible applications of Quantum Groups to physical phenomena [10-12].

Due to the nature of deformed Hamiltonians, made of bosonic q -oscillators, it is quite difficult to obtain exact expressions when studying their statistical properties. Most of the works done in this area have considered approximations around q (the deformation parameter) equal to one. In this paper we are going to analyse the large q limit of deformed systems. In section 2 we consider the canonical ensemble for the bosonic q -oscillators and we find that for infinite deformation the statistical properties are those of fermions. In addition, it is shown that for large q the system behaves like a deformation of fermions. Section 3 is devoted to the study of a deformed ideal system with large q , and we find the virial expansion for its equation of state. Final remarks are given in section 4.

2 Bosonic q -Oscillators in the large q limit

One calls bosonic q -oscillators the associative algebra generated by the elements α, α^+ and N satisfying the relations [10-16]

$$\begin{aligned} [N, \alpha^+] &= \alpha^+ , \quad [N, \alpha] = -\alpha \\ [\alpha, \alpha^+]_{\alpha} &= f_{\alpha}(N). \end{aligned} \quad (2.1)$$

We are going to consider here the following forms of the above algebra (2.1):

$$[a, a^+]_a \equiv aa^+ - qa^+a = q^{-N} \quad (2.2.a)$$

$$[A, A^+]_A \equiv AA^+ - q^2A^+A = 1. \quad (2.2.b)$$

The above two algebras can be related to each other via

$$A = q^{N/2}a \quad , \quad A^+ = a^+q^{N/2}$$

with q a real parameter.

It is possible to construct representations of the relations (2.2) in the Fock space \mathcal{F} spanned by the normalized eigenstates $|n\rangle$ of the number operator N as

$$\begin{aligned} \alpha|0\rangle &= 0 \quad , \quad N|n\rangle = n|n\rangle \quad n = 0, 1, 2, \dots \\ |n\rangle &= \frac{1}{\sqrt{[n]_{\alpha}!}} (\alpha^+)^n |0\rangle \end{aligned} \quad (2.3)$$

where $[n]_\alpha! \equiv [n]_\alpha \cdots [1]_\alpha$, $[n]_\alpha = (q^n - q^{-n})/(q - q^{-1})$ and $[n]_A = (q^{2n} - 1)/(q^2 - 1)$.

In the Fock space \mathcal{F} it is possible to express the deformed oscillators in terms of the standard bosonic ones b, b^+ as [16-17]

$$\alpha = \left(\frac{[N+1]_\alpha}{N+1} \right)^{1/2} b, \quad \alpha^+ = b^+ \left(\frac{[N+1]_\alpha}{N+1} \right)^{1/2}; \quad (2.4)$$

it can easily be shown in \mathcal{F} that

$$\alpha\alpha^+ = [N+1]_\alpha, \quad \alpha^+\alpha = [N]_\alpha, \quad (2.5)$$

and as expected the standard bosonic algebra is obtained in the $q \rightarrow 1$ limit.

We are now going to investigate highly deformed q -bosons ($q \rightarrow \infty$ limit). In this limit for $n \geq 2$, $[n]_\alpha \rightarrow \infty$ and as a result when $q = \infty$ Fock space (2.3) is reduced to a fermionic one since the eigenstates $|n\rangle$ vanish for $n \geq 2$. Consequently, the statistical properties of q -deformed oscillators (2.2) become those of fermions.

In order to exhibit the statistical properties close to this fermionic limit let us consider the canonical partition function for the Hamiltonian

$$H = \omega A^+ A = \omega [N]_A, \quad (2.6)$$

which is given by

$$Z = 1 + e^{-\beta\omega} + e^{-\beta\omega(1+q^2)} + \dots + e^{-\beta\omega[1+q^2+\dots+q^{2(n-1)}]} + \dots \quad (2.7)$$

where $\beta = (k_B T)^{-1}$, with k_B the Boltzmann constant. The above expression has often been taken as the starting point in the analysis of q -bosons at finite temperature [11-15], which is indeed the case when q is close to one. On the other hand for infinite deformations, (2.7) is clearly the partition function of fermions. Therefore, in the canonical ensemble, when $q \gg 1$ expression (2.7) can be understood as a deformation around fermions.

The average of N to first order in the large q limit is given by

$$\langle N \rangle \cong \frac{1 + 2e^{-\beta\omega q^2}}{1 + e^{\beta\omega} + e^{-\beta\omega q^2}}. \quad (2.8)$$

As expected in the $q = \infty$ limit the Fermi-Dirac distribution is recovered.

3 Highly Deformed Ideal q -Gas

Following the standard lore [14] we define the Hamiltonian of an ideal deformed system as:

$$H = \sum_i \omega_i A_i^+ A_i = \sum_i \omega_i [N_i]_A, \quad (3.1)$$

where A_i, A_i^+ and N_i are interpreted respectively as annihilation, creation and occupation number operators of particles in level i , with energy ω_i . These operators satisfy the algebra (2.2b) and commute for different levels.

The grand canonical partition function is given by:

$$Z = \text{Tr} \exp[-\beta(H - \mu N)] = \exp(-\beta\Omega) \quad (3.2)$$

where N is the total number operator

$$N = \sum_i N_i, \quad (3.3)$$

μ is the chemical potential and Ω is the grand canonical potential. For the above system Z factorizes and the grand canonical potential is given by a sum over single level partition functions

$$\Omega = -\frac{1}{\beta} \sum_i \log Z_i^0(\omega_i, \beta, \mu), \quad (3.4)$$

where

$$Z_i^0(\omega_i, \beta, \mu) = \sum_{n=0}^{\infty} e^{-\beta(\omega_i[n]\Lambda - \mu n)}. \quad (3.5)$$

The energy of the non-relativistic q -boson is

$$\omega_i = \vec{p}^2/2m, \quad (3.6)$$

and the usual approach [18] is to enclose the system in a large volume V , which allows the sum over levels to be replaced by an integral over the p space:

$$\sum_i \rightarrow \frac{V}{(2\pi h)^3} \int d^3 p. \quad (3.7)$$

If we keep only the first correction in the large q limit, the grand canonical potential is found to be

$$-\beta\Omega \cong \frac{1}{2\pi^2} V h^{-3} \int_0^{\infty} dp p^2 \ln[1 + z e^{-\beta p^2/2m} + z^2 e^{-\beta(q^2+1)p^2/2m}], \quad (3.8)$$

where z is the fugacity, $z = e^{\beta\mu}$; later we shall discuss the region of validity of the above approximation. After integrating by parts and defining the new variable $\eta = \beta p^2/2m$ (3.8) reduces to

$$-\beta\Omega \cong \frac{V}{6\pi^{7/2}} \Lambda^{-3} \int_0^{\infty} d\eta \eta^{3/2} \frac{z e^{-\eta} + z^2(q^2+1)e^{-(q^2+1)\eta}}{1 + z e^{-\eta} + z^2 e^{-(q^2+1)\eta}}, \quad (3.9)$$

where $\Lambda = (h^2\beta/2\pi m)^{1/2}$, also called thermal wavelength, is the relevant expansion parameter in the thermodynamic functions.

Finally, assuming that the fugacity z is small compared to one, expanding Ω and keeping terms up to third order in z the pressure $P = -\Omega/V$ is given by

$$P = \frac{\beta^{-1}\Lambda^{-3}}{(2\pi)^3} z \{1 + z[-2^{-5/2} + (q^2+1)^{-3/2}] + z^2[3^{-5/2} - (q^2+2)^{-3/2}] + 0(z^3)\}. \quad (3.10)$$

The q -boson density $n = \frac{\partial P}{\partial \mu}|_{T,V}$ is easily found to be

$$n = \frac{\Lambda^{-3}}{(2\pi)^3} z \{1 + 2z[-2^{-5/2} + (q^2+1)^{-3/2}] + 3z^2[3^{-5/2} - (q^2+2)^{-3/2}] + 0(z^3)\}. \quad (3.11)$$

Inverting the power series above we obtain

$$z = n\Lambda^3 - 2Q_1(n\Lambda^3)^2 + (8Q_1^2 - 3Q_2)(n\Lambda^3)^3 + \dots, \quad (3.12)$$

where

$$Q_1 = -2^{-5/2} + (q^2 + 1)^{-3/2}, \quad Q_2 = 3^{-5/2} - (q^2 + 2)^{-3/2}. \quad (3.13)$$

Substituting (3.12-13) in (3.10) and expanding in powers of n , we obtain the virial expansion of the equation of state

$$P = \frac{n}{(2\pi)^3\beta} \left\{ 1 + \left[\frac{1}{2^{5/2}} - \frac{1}{(q^2 + 1)^{3/2}} \right] n\Lambda^3 + \left[\frac{1}{8} - \frac{2}{3^{5/2}} + \frac{4}{(q^2 + 1)^3} - \frac{4}{(2q^2 + 2)^{3/2}} + \frac{2}{(q^2 + 2)^{3/2}} \right] n^2\Lambda^6 + \dots \right\}. \quad (3.14)$$

Looking at eq. (3.14) one immediately sees that in the $q = \infty$ limit our q -gas behaves exactly like a non-relativistic Fermi-gas [18]. For finite large values of q the pressure is reduced with respect to the Fermi-gas. We would like to stress that analogously to non-deformed Bose or Fermi gases, the approximations done here are valid for large $V, z \ll 1$ and $n\Lambda^3 \ll 1$, implying that for a given density n , we have a high-temperature approximation or, for a given temperature, a low-density approximation.

A similar procedure is employed in the case of ultrarelativistic q -bosons whose energy is given by $\omega_i = c\vec{p}$. The virial expansion of the equation of state in this regime is

$$P = \frac{n}{(2\pi)^3\beta} \left\{ 1 + \left[\frac{1}{2^4} - \frac{1}{(q^2 + 1)^3} \right] \Delta^3 n + \left[\frac{1}{2^6} - \frac{2}{3^4} - \frac{1}{2(q^2 + 1)^3} - \frac{2}{(q^2 + 2)^3} + \frac{4}{(q^2 + 1)^6} \right] \Delta^6 n^2 + \dots \right\}, \quad (3.15)$$

where now the relevant parameter of the expansion is $\Delta^3 n$ with $\Delta = ch\beta/2\pi^{1/3}$, the so called optical wavelength. The comments in the last paragraph about the approximations performed remain valid for (3.15). In the $q = \infty$ limit (3.15) is the ultra-relativistic equation of state and also here the effect of finite q is to reduce the pressure as compared with the infinitely deformed gas.

4 Final Comments

In this paper we have analysed some statistical properties of q -oscillators in the large q limit. It is quite remarkable that infinitely deformed q -bosons acquire a fermionic behaviour. This point deserves further investigation at the level of quantum algebras.

Due to the relevance of Fermi-gases in Condensed Matter and Nuclear Physics we expect that our results can find an application in these fields.

ACKNOWLEDGEMENTS

The authors thank C. Tsallis for enlightening discussions.

References

- [1] V.G. Drinfeld, *Sov. Math. Dokl.* **32** (1985) 254;
- [2] M. Jimbo, *Lett. Math. Phys.* **10** (1985) 63; **11** (1986) 247;
- [3] L.D. Faddeev, N. Yu. Reshetikhin and L.A. Takhtadzhyan, *Algebra and Analysis* **1** (1987) 178;
- [4] For reviews see for instance: S. Majid, *Int. J. Mod. Phys.* **A45** (1990) 1;
P. Aschieri and L. Castellani, *Int. J. Mod. Phys.* **A8** (1993) 1667;
M.R.-Monteiro, "Introduction to Quantum Groups", preprint CBPF-NF-061/93, to appear in the Proceedings of XIV ENFPC, Caxambú, Brazil;
- [5] C. Zachos, *Contemporary Mathematics* **134** (1992) 351 (and references therein);
- [6] J.L. Matheus-Valle and M.R.-Monteiro, *Mod. Phys. Lett.* **A7** (1992) 3032; *Phys. Lett.* **B66** (1993) 330;
- [7] L. Castellani and M.R.-Monteiro, *Phys. Lett.* **B314** (1993) 25;
- [8] A. Lerda and S. Sciuto, *Nucl. Phys.* **B401** (1993) 613;
R. Caracciolo and M.R.-Monteiro, *Phys. Lett.* **B308** (1993) 58;
M. Frau, M.R.-Monteiro and S. Sciuto, " q -Deformed Lie Algebras and Their Anyonic Realization", preprint DFTT 16/93, CBPF-NF-26-93, to appear in *J. Phys. A*; J.L. Matheus-Valle and M.R.-Monteiro, "Anyonic construction of the $sl_{q,s}(2)$ algebra", preprint CBPF-NF-062/93;
- [9] V.V. Kuryshkin, *Ann. Found. L. de Broglie* **5** (1980) 111;
- [10] A.J. Macfarlane, *J. Phys.* **A22** (1989) 4581; L.C. Biedenharn, *J. Phys.* **A22** (1989) L873; M. Chaichian and P. Kulish, *Phys. Lett.* **B234** (1990) 72;
- [11] M. Martin-Delgado, *J. Phys.* **A24** (1991) 1285;
P. Neškovic and B. Urošević, *Int. J. of Mod. Phys.* **A7** (1992) 3379;
- [12] V. Man'ko, G. Marmo, S. Solimeno and F. Zaccaria, *Int. J. of Mod. Phys.* **A8** (1993) 3577; S. Vokos and C. Zachos, "Thermodynamic q -Distributions that Aren't", preprint UW/PT-93-05, ANL-HEP-CP-93-39;
- [13] V. Man'ko, G. Marmo, S. Solimeno and F. Zaccaria, *Phys. Lett.* **A176** (1993) 173;
- [14] M. Chaichian, R. Gonzalez Felipe and C. Montonen, *J. Phys.* **A26** (1993) 4025;
- [15] M.R.-Monteiro and I. Roditi, "Thermo-Field Dynamics of Deformed Systems", preprint CBPF-NF-037/93, to appear in *Mod. Phys. Lett. B*; "Deformed Systems at Finite Temperature", preprint CBPF-NF-060/93;
- [16] P. Kulish and E. Damaskinsky, *J. Phys.* **A23** (1990) L415;
- [17] A. Polychronakos, *Mod. Phys. Lett.* **A5** (1990) 2325;

- [18] See for instance: "Equilibrium and Non-Equilibrium Statistical Mechanics", Radu Balescu, John Wiley & Sons, New York, 1975; K. Huang, Statistical Mechanics, John Wiley & Sons, New York, 1963.