



Instituto de Física Teórica
Universidade Estadual Paulista

July/94

IFT-P.025/94

**Many-body problems with composite particles and
 q -Heisenberg algebras**


S. S. Avancini

*Departamento de Física-CFM
Universidade Federal de Santa Catarina
Cx. Postal 476
88040-900 - Florianópolis, S.C.
Brazil*

and

G. Krein

*Instituto de Física Teórica
Universidade Estadual Paulista
Rua Pamplona 145
01405-900 - São Paulo, S.P.
Brazil*



**Instituto de Física Teórica
Universidade Estadual Paulista
Rua Pamplona, 145
01405-900 – São Paulo, S.P.
Brazil**

Telephone: 55 (11) 251.5155

Telefax: 55 (11) 288.8224

Telex: 55 (11) 31870 UJMFBR

Electronic Address: LIBRARY@IFT.UESP.ANSP.BR
47553::LIBRARY

Many-body problems with composite particles and q -Heisenberg algebras

S.S. Avancini

Departamento de Física-CFM, Universidade Federal de Santa Catarina

Cxa. Postal 476 - 88040-900 Florianópolis-SC - Brazil

G. Krein

Instituto de Física Teórica - Universidade Estadual Paulista

Rua Pamplona, 145 - 01405-900 São Paulo-SP - Brazil

Abstract

We propose to employ deformed commutation relations to treat many-body problems of composite particles. The deformation parameter is interpreted as a measure of the effects of the statistics of the internal degrees of freedom of the composite particles. A simple application of the method is made for the case of a gas of composite bosons.

PACS:02.00.03,65.Fd,02.10.N

In recent years there has been a great deal of interest in the subject of deformations of the basic commutation relations of Fermi and Bose fields. One focus of interest is on possible (small) violations of the Fermi and Bose statistics by particles (“quons”) whose annihilation and creation operators obey deformed commutation relations which interpolate between bosons and fermions[1]. Another focus of interest closely related, not only in physics but also in mathematics, is the subject of quantum algebras[2], whose origins are in the study of the Yang-Baxter equations connected with the quantum inverse scattering problem. The discovery by Macfarlane[3] and Biedenharn[4] of a new realization of the quantum algebra $su(2)_q$ in terms of q -analogues of the harmonic oscillator has given rise to speculations on possible applications in real physical problems. In addition to speculations on small violations of Fermi or Bose statistics[1] and on possible generalizations of quantum mechanics at higher energies (e.g., in the early universe)[5], quantum algebras have been used with relative success in phenomenological studies of deformed nuclei[6], diatomic molecules[7], spin chains[8], and anyonic oscillations with fractional statistics[9]. Despite of the successful phenomenological applications, a clear physical meaning of the deformation parameter q is lacking.

In this letter we propose to employ q -Heisenberg algebra[10] as a convenient tool to describe many-body problems involving composite particles (i.e. not point-like). We argue that the physical meaning of the deformation parameter is that it is a measure of the effects of the statistics of the internal degrees of freedom of the composite particles, and its value depends on the “degree of overlap” of the extended structure of the particles in the medium. The interpretation of deformed algebras as describing composite particles is not new and can be found in several places in the literature (see e.g. Ref. [1]); here we explicitly demonstrate in a simple example the realization of this. Many-body problems involving composite particles have complications in addition to the usual ones involving point particles due to the simultaneous presence of “macroscopic” (composites) and “microscopic” (constituents) degrees of freedom. Due to the internal degrees of freedom, the algebra of

the creation and annihilation operators of the composite particles deviates from the usual canonical ones for point particles; it becomes deformed. It is therefore natural to speculate on the possibility that the deformation of the algebra of the creation and annihilation operators would provide a convenient way to effectively take into account the microscopic degrees of freedom of the composites. The deformation of the algebra is the price one pays for not taking into account explicitly the microscopic degrees of freedom. Of course, one can hope for the success of such a program for systems where the degree of overlap of the internal structures of the particles is not very large, in the sense that the system is not dissolved into its constituents. A related work [11] describes the possibility of using deformed commutation relations to treat correlated fermion pairs in a single-j nuclear shell.

We start with a heuristic discussion on the relation of the deformation parameter with the composite nature of bosons. We consider a composite boson state with quantum number α as a bound state of two distinct fermions

$$A_\alpha^\dagger|0\rangle = \sum_{\mu\nu} \Phi_\alpha^{\mu\nu} a_\mu^\dagger b_\nu^\dagger|0\rangle, \quad (1)$$

where $\Phi_\alpha^{\mu\nu}$ is the bound-state wave-function, a_μ^\dagger and b_μ^\dagger are the fermion creation operators, and $|0\rangle$ is the vacuum state. The quantum number α stands for the center of mass momentum, the internal energy, the spin, and other internal degrees of freedom of the composite boson. The μ and ν stand for the space and internal quantum numbers of the constituent fermions. The sum over μ and ν is to be understood as a sum over discrete quantum numbers and an integral over continuous variables.

The fermion creation and annihilation operators satisfy canonical anti-commutation relations:

$$\begin{aligned} \{a_\mu, a_\nu\} = \{a_\mu^\dagger, a_\nu^\dagger\} &= 0 & \{b_\mu, b_\nu\} = \{b_\mu^\dagger, b_\nu^\dagger\} &= 0, \\ \{a_\mu, a_\nu^\dagger\} = \delta_{\mu\nu} & & \{b_\mu, b_\nu^\dagger\} &= \delta_{\mu\nu}. \end{aligned} \quad (2)$$

It is convenient to work with normalized wave-functions $\Phi_\alpha^{\mu\nu}$, such that

$$\langle\alpha|\beta\rangle = \delta_{\alpha,\beta}, \quad (3)$$

and therefore

$$\sum_{\mu\nu} \Phi_\alpha^{\mu\nu*} \Phi_\beta^{\mu\nu} = \delta_{\alpha,\beta}. \quad (4)$$

Using the fermion anticommutation relations of Eq. (2) and the wave-function normalization Eq. (4), one can easily show that the composite boson operators satisfy the following commutation relations:

$$\begin{aligned} [A_\alpha, A_\beta] &= [A_\alpha^\dagger, A_\beta^\dagger] = 0, \\ [A_\alpha, A_\beta^\dagger] &= \delta_{\alpha,\beta} - \Delta_{\alpha,\beta}, \end{aligned} \quad (5)$$

where $\Delta_{\alpha,\beta}$ is given by

$$\Delta_{\alpha,\beta} = \sum_{\mu\nu} \Phi_\alpha^{\mu\nu*} \left(\sum_{\mu'} \Phi_\beta^{\mu'\nu} a_\mu^\dagger a_{\mu'} + \sum_{\nu'} \Phi_\beta^{\mu\nu*} b_\nu^\dagger b_{\nu'} \right). \quad (6)$$

One can also easily show the following commutation relations:

$$[a_\mu, A_\alpha^\dagger] = \sum_{\mu'\nu} \delta_{\mu\mu'} \Phi_\alpha^{\mu'\nu} b_\nu^\dagger, \quad [b_\nu, A_\alpha^\dagger] = - \sum_{\mu'\nu'} \delta_{\nu\nu'} \Phi_\alpha^{\mu'\nu'} a_\mu^\dagger. \quad (7)$$

The composite nature of the bosons is evident from the presence of $\Delta_{\alpha,\beta}$, it is a sort of a “deformation” of the canonical boson algebra. The effect of this term becomes unimportant in the infinite tight binding limit, i.e. in the limit of point-like bosons. Eq. (7) shows the lack of kinematical independence of the microscopic operators a_μ and b_ν from the macroscopic ones A_α 's.

Now let us consider a system of N composite bosons in a box of volume V at zero temperature. If the bosons were ideal point-like particles, the ground state of the system would be the one where all bosons condense in the zero momentum state. In the case of composite bosons, the closest analog of the ideal gas ground state is

$$|N\rangle = \frac{1}{\sqrt{N!}} (A_0^\dagger)^N |0\rangle, \quad (8)$$

where A_0^\dagger is the creation operator of a composite boson in its ground state (ground state Φ) and with zero center of mass momentum. Due to the composite nature of the bosons, this state incorporates kinematical correlations implied by the Pauli exclusion principle which operates on the constituent fermions. Among other effects, the Pauli principle forbids the macroscopic occupation of the zero momentum state. The closest analog to the boson occupation number in the state (8) is

$$N_0 = \frac{\langle N | A_0^\dagger A_0 | N \rangle}{\langle N | N \rangle}. \quad (9)$$

In order to evaluate (9), we consider a spin zero boson and use for the spatial part of Φ a simple Gaussian form such that the r.m.s radius of the boson is r_0 . To lowest order in the density of the system $n = N/V$, N_0 is given by

$$N_0 = N \left(1 - \gamma n r_0^3 \right), \quad (10)$$

where $\gamma = 4\pi^{-3/2} \simeq 1$ is a numerical factor that comes from the functional form of the wave function. If we had taken another functional form for Φ , we still would have obtained for N_0 the result of Eq. (10), but γ would have taken a different value.

It is apparent from Eq. (10) that in the limit of infinite tight binding, $r_0 \rightarrow 0$, one has the familiar Bose-Einstein condensation. For finite values of the size of the bosons, the effects of the Pauli principle become important and one has a depletion on the amount of condensed bosons. Moreover, from (10) one has that if the size of the bound state is of the order of the mean separation of the bosons in the medium, $d \sim n^{-1/3}$, the depletion is almost total. The depletion of the condensation is a direct consequence of the deformation of the boson algebra by the term $\Delta_{\alpha\beta}$.

Next we show that the effect of the composite nature of the boson can be effectively taken into account employing a deformed boson algebra. The deformed boson commutation relations are given by

$$A_\alpha A_\beta^\dagger - q^2 A_\beta^\dagger A_\alpha = \delta_{\alpha\beta}. \quad (11)$$

where q^2 is the deformation parameter of the algebra; A_α annihilates the vacuum

$$A_\alpha |0\rangle = 0. \quad (12)$$

Note that no commutation relation can be imposed on $A_\alpha A_\beta^\dagger$ and $A_\alpha A_\alpha$. However, as remarked by Greenberg[1], similarly to the case of normal Bose commutation relations, no such rule is needed for practical evaluation of expectation values of polynomials in A_α and A_α^\dagger when (12) holds. Such matrix elements can be evaluated with the repeated use of Eq. (11) solely; annihilation operators are moved to the right using (11) until they annihilate the vacuum or creation operators are moved to the left using the adjoint of (11) until they annihilate the vacuum.

If one writes $q^2 = 1 - x$, the deformed commutator can be written as

$$[A_\alpha, A_\beta^\dagger] = \delta_{\alpha\beta} - x A_\alpha^\dagger A_\beta. \quad (13)$$

The similarity of this with the second equation of (5) is evident. In some sense, the weighted (by the Φ 's) fermion operators $a_\mu^\dagger a_\mu$ and $b_\mu^\dagger b_\mu$ in (5) are effectively modeled by the term $x A_0^\dagger A_0$.

In the same spirit as in the previous case, we take as the closest analog of the ideal boson gas ground state the N q-boson state

$$|N\rangle = \frac{1}{\sqrt{|N|!}} (A_0^\dagger)^N |0\rangle, \quad (14)$$

where $|N|! = [N][N-1][N-2]\dots 1$ is the q -factorial, and

$$[N] = \frac{1 - q^{2N}}{1 - q^2}. \quad (15)$$

As before, the operator $A_0^\dagger A_0$ is the number operator in the zero deformation limit only. The effect of the deformation can be evaluated taking the expectation value of the $A_0^\dagger A_0$ in the state $|N\rangle$ of equation (14). To evaluate the expectation value, we make use of the result[3]

$$A_0^\dagger A_0 = [\hat{N}_0] = \frac{1 - q^{2N_0}}{1 - q^2}, \quad (16)$$

where \hat{N} is the number operator in the deformed algebra

$$\hat{N}_0|N\rangle = N|N\rangle. \quad (17)$$

Using the result of (16), one obtains to lowest order in x ,

$$N_0 = \frac{1}{x} [1 - (1 - x)^N] \simeq N \left(1 - \frac{1}{2}Nx\right). \quad (18)$$

Comparing this result with the one of Eq. (10) it is clear that the effect of the deformation is such that

$$N_x \sim \frac{Nx_0^3}{V}, \quad (19)$$

that is, the effect of the deformation parameter is proportional to the ratio of the volume occupied by the bosons to the volume of the system. It is clear that when this ratio is small, meaning that there is no considerable overlap of the boson internal wave functions, the system behaves as a normal boson gas.

This completes our heuristic discussion on the plausibility of using q-Heisenberg algebra as a means of treating effectively the internal degrees of freedom of composite particles in a many-body system. Next, we consider the thermodynamics of an ideal q-bose gas[12, 13]. This example is taken to illustrate the effects of the constituent fermions on the thermodynamic properties of a gas of composite bosons. The Hamiltonian of the deformed bose gas is defined[12] as:

$$H = \sum_i \epsilon_i A_i^\dagger A_i, \quad (20)$$

where A_i^\dagger , A_i are respectively the creation and annihilation operator of a q - boson in the state of energy ϵ_i . The energy ϵ_i is the kinetic energy plus the internal ground state energy of the boson.

It follows from Eq. (16) that the energy eigenvalues of H are given by

$$E(n) = \sum_i \epsilon_i [n_i] = \sum_i E_i(n_i), \quad (21)$$

where $n = \sum_i n_i$ and n_i are the occupation numbers associated to the state of energy ϵ_i . In order to study the thermodynamics of the deformed system we evaluate the grand canonical partition function:

$$Z = Tr e^{-\beta(H - \mu N)} = e^{-\beta\Omega} \quad (22)$$

where $\beta = 1/kT$, μ is the chemical potential and \hat{N} is the number operator. It follows from Eqs. (20) and (22) that in the representation where the Hamiltonian is diagonal:

$$Z = \prod_i Z_1(i, \beta, \mu) \quad (23)$$

with

$$Z_1(i, \beta, \mu) = \sum_{n_i=0}^{\infty} e^{-\beta(\epsilon_i(n_i) - \mu n_i)}$$

As the exact calculation of Z_1 is not possible, we consider its expansion in terms of the parameter x , where we recall that $q^2 = 1 - x$. (Note that this corresponds to an expansion in the density of the q - gas). In order to carry out such expansion we note that:

$$[n] = \frac{1 - q^{2n}}{1 - q^2} = \sum_{k=1}^n \binom{n}{k} x^{k-1}.$$

From the equation above and Eq. (23) we obtain for the expansion of Z_1 up to second order in x the following:

$$\begin{aligned} Z_1(i, \beta, \mu) = & Z_0(z, y_i) \left[1 + y_i (z e^{-\beta y_i})^2 Z_0^2(z, y_i) x \right. \\ & + \left. \left(\frac{y_i^2}{2} + (3y_i - 1)y_i z e^{-\beta y_i} Z_0(z, y_i) + 3y_i^2 (z e^{-\beta y_i})^2 Z_0^2(z, y_i) \right) \right. \\ & \left. \cdot (z e^{-\beta y_i})^2 Z_0^2(z, y_i) x^2 + \dots \right]. \end{aligned} \quad (24)$$

In the above the usual definition of fugacity $z = e^{\beta\mu}$ and $y_i = \beta\epsilon_i$ have been used and

$$Z_0(z, y_i) = \sum_{n=0}^{\infty} (z e^{-\beta y_i})^n = \frac{1}{1 - z e^{-\beta y_i}}.$$

From Eqs. (22, 23) we obtain for $\beta\Omega$, up to second order in x :

$$\begin{aligned} \beta\Omega = & -\sum_i \log Z_{0i}(z, y_i) - \sum_i y_i (ze^{-y_i})^2 Z_{0i}^2(z, y_i) x \\ & - \sum_i \left[\frac{y_i^2}{2} (ze^{-y_i})^2 Z_{0i}^2(z, y_i) + (3y_i - 1) y_i (ze^{-y_i})^3 Z_{0i}^3(z, y_i) \right. \\ & \left. + \frac{5}{2} y_i^2 (ze^{-y_i})^4 Z_{0i}^4(z, y_i) \right] x^2 + O(x^3) \end{aligned} \quad (25)$$

Proceeding as usual, we consider the system contained in a large container of volume V , and replace the summations by an integral as:

$$\sum_i f(\epsilon_i) \rightarrow \frac{V}{2\pi^3} \int d^3k f(k).$$

We take for the kinetic energy of the boson the nonrelativistic expression $\epsilon_i = \epsilon_i(k) = \gamma \bar{k}^2$, where $\gamma = 1/2m$. In the case of a composite boson one could have, in addition to the kinetic energy, internal excitation energies. Our choice of ϵ_i represents the energy spectrum of composite bosons in their internal ground states. Since the constant internal ground state energy of the bosons do not play any role in the thermodynamic properties of the system, one needs to consider only the kinetic energy spectrum. We note that all the integrals that appear in the evaluation of $\beta\Omega$, through the procedure just outlined, can be obtained by successive z derivatives of the equation

$$\int_0^\infty dy y^{a-1} \log(1 - ze^{-y}) = -\Gamma(a) g_{a+1}(z),$$

where a is a real number and $g_l(z) = \sum_{k=1}^\infty z^k / k^l$.

Thus the pressure per unit volume can be straightforwardly obtained:

$$\begin{aligned} \beta p \equiv -\beta\Omega = & a \left\{ g_{5/2}(z) + \frac{3}{2} [g_{3/2}(z) - g_{5/2}(z)] x \right. \\ & \left. + \left[\frac{13}{16} g_{1/2}(z) - \frac{3}{2} g_{3/2}(z) + \frac{11}{16} g_{5/2}(z) \right] x^2 + O(x^3) \right\}, \end{aligned} \quad (26)$$

where $a = (m/2\pi\beta)^{3/2}$.

With this, one can obtain the virial expansion of the equation of state. The density, ($n = N/V$), is obtained by the usual expression:

$$n = \frac{N}{V} = \left(\frac{\partial p}{\partial \mu} \right)_{T,V} = z \left(\frac{\partial \beta p}{\partial z} \right)_{\beta}.$$

Therefore, using Eq. (26), one obtains

$$\begin{aligned} n = & a \left\{ g_{3/2}(z) + \frac{3}{2} [g_{1/2}(z) - g_{3/2}(z)] x \right. \\ & \left. + \left[\frac{13}{16} g_{-1/2}(z) - \frac{3}{2} g_{1/2}(z) + \frac{11}{16} g_{3/2}(z) \right] x^2 + \dots \right\}, \end{aligned} \quad (27)$$

where we have used the relation

$$z \frac{d g_l(z)}{dz} = g_{l-1}(z).$$

In the low density regime one may write:

$$z = a_1 \left(\frac{n}{a} \right) + a_2 \left(\frac{n}{a} \right)^2, \quad (28)$$

Using this in Eq. (27), and keeping only terms up to second order in x one obtains:

$$\begin{aligned} a_1 = & 1 \\ a_2 = & -\frac{1}{2^{3/2}} \left(1 + \frac{3}{2} x - \frac{3}{16} x^2 \right). \end{aligned} \quad (29)$$

Through the substitution of z given in Eq. (28) in Eq. (25) the virial expansion is obtained,

$$\beta p = n(1 + B n + \dots), \quad (30)$$

where

$$B = -\frac{1}{2^{3/2}} \frac{1}{a} \left(\frac{1}{2} + \frac{3}{4} x - \frac{21}{32} x^2 \right).$$

Note that in the limit of zero deformation, $x = 0$, one obtains the usual result [14]. For finite, small x , one sees that B becomes more negative than in the limit of zero deformation. Thus the deformation has the effect of an attractive potential. This result is easily understood in terms of the interpretation of the deformation parameter as related to the internal fermion degrees of freedom of the composite boson: the Fermi statistics of the constituents cause a depletion of the boson occupation numbers, and in the sum of Eq. (21) this depletion has the effect of an attractive potential, $\sum_i \epsilon_i |n_i| \leq \sum_i \epsilon_i n_i$.

Concluding, we have discussed the possibility of studying many-body problems with composite particles in terms of deformed algebras for the creation and annihilation operators (q -Heisenberg algebras) of the composites. We have considered the example of a gas of bosons and examined the role of the deformation of the algebra of the boson operators. The interpretation of the deformation is that it models the effects of the Pauli principle operating on the constituent fermions. Among other effects, the deformation of the algebra causes the depletion of the single particle occupation of the bosons.

Although we have discussed the example of a system of composite bosons, it is clear that the same methods is applicable to composite fermions. We plan to study such case in a future publication. Another interesting subject of future work is the study of interacting composite bosons (or fermions). In particular, it would be very interesting to explore the role of the deformation on phase transitions. Such a study might be of interest to superconductivity/superfluidity and quark-gluon plasma phase transitions.

Acknowledgments: Work partially supported by CNPq and FAPESP.

References

- [1] O.W. Greenberg, Phys. Rev. D **43**, 4111(1991).
- [2] *Quantum Groups Workshop*, Proceedings, Argonne, Illinois, edited by T. Curtright, D. Fairlie, and C. Zachos (World Scientific, Singapore, 1991).
- [3] A.J. Macfarlane, J. Phys. A **22**, 4581(1989).
- [4] L.C. Biedenharn, J. Phys. A **22**, L873(1989).
- [5] Y.J. Ng, J. Phys. A **23**, 1023(1990); S.V. Shabanov, J. Phys. A **26**, 2583(1993).
- [6] P.P. Raychev, R.P. Roussev, and Yu. F. Smirnov, J.Phys. G **16**, L137(1990); D. Bonatsos, E.N. Argyres, S.B. Drenska, P.P. Raychev, R.P. Roussev, and Yu.F. Smirnov, Phys. Lett. B **251**, 477(1990); D. Bonatsos, S.B. Drenska, P.P. Raychev, R.P. Roussev, and Yu.F. Smirnov, J. Phys. G. **17**, L67(1991).
- [7] D. Bonatsos, S.B. Drenska, P.P. Raychev, R.P. Roussev, and Yu.F. Smirnov, Chem. Phys. Lett. **175**, 300(1990).
- [8] M.T. Batchelor, L. Mezincescu, R.I. Nepomechie, and V. Rittenberg, J. Phys. A **23**, L141(1990); P.P. Kulish and E.K. Sklyanin, J. Phys. A **24**, L435(1991).
- [9] A. Lerda and S. Sciuto, Nucl Phys. B **401**, 613(1993).
- [10] D. B. Fairlie and C. Zachos, proceedings of the NATO ARW on Quantum Field Theory, Statistical Mechanics, Quantum Groups, and Topology, ed. T. Curtright, Miami, 1991
- [11] D. Bonatsos, J. Phys. A:Math. Gen. 25(1992)L101; D. Bonatsos, C. Daskaloyannis and A. Faessler, J. Phys. A:Math. Gen. 27(1994)1299.
- [12] M. Chaichian, R. Gonzalez Felipe and C. Montonen, J. Phys. A:Math. Gen. 26(1993)4025

- [13] M. Martin-Delgado, J. Phys. A: Math. Gen. 24(1991)11285
- [14] L.D. Landau and E.M. Lifshitz, Statistical Physics (Pergamon Press, New York, 1966),(2nd. edition), Eq. (55.15) page 151.

