

BB

E1-95-263

CERN LIBRARIES, GENEVA

SW 9543



SCAN-9510172

FIRST MEASUREMENT
OF THE TENSOR ANALYSING POWER, T_{20}
IN INELASTIC $(d, d')X$ SCATTERING
AT 0° ON 1H AND ^{12}C AT 4.5 AND 5.5 GeV/c

Submitted to «Physics Letters B»

1995

L.S.Azhgirey, E.V.Chernykh, A.P.Kobushkin¹, P.P.Korovin², B.Kuehn³,
V.P.Ladygin, S.Nedev, C.F.Perdrisat⁴, N.M.Piskunov, V.Punjabi⁵, I.M.Sitnik,
G.D.Stoletov, E.A.Strokovsky⁶, A.I.Syamtomov¹, S.A.Zaporozhets

¹ITP, 252130, Metrologicheskaya 14, Kiev, Ukraine

²Samara Air-Space University, 443086, Samara, Moskovskoe Shosse 34, Russia
³01326 Dresden, Dresdner Str. 57, Germany

⁴The College of William and Mary, Williamsburg, VA 23185 USA

⁵Norfolk State University, Norfolk, VA 23504 USA

⁶E-mail address: strok@sunhe.jinr.dubna.su

Characteristics of broad hadronic resonances excited in nuclei and the related nuclear medium response on high energy excitations have been discussed intensively during last decade. Still, a wide gap remains between experimental results and theory[1]; apart from a few exceptions, most of the well established experimental observations don't have a *quantitative* theoretical interpretation.

The main interest and at the same time the main difficulties of these studies, is related to the fact that the behaviour of nuclear matter at high excitation energies is governed not only by its nucleonic degrees of freedom, but also by the internal degrees of freedom of the constituent nucleons. When the energy "pumped" into the nuclear medium is close to the characteristic energy of excitation of the internal degrees of freedom of a nucleon, these can no longer be treated independently. Such excitations reveal themselves as nuclear $N \rightarrow N^*$, transitions followed by radiation of particles.

The non-trivial difference between resonance excitation off a free proton and off nuclei was first observed in experiments measuring inelastic charge exchange cross sections; it was shown that the properties of Δ -excitations of nuclei cannot be described in the picture of quasifree production[2]. Non-quasifree mechanisms observed in inclusive experiments were confirmed in exclusive experiments[3] but a number of theoretical uncertainties still persist[4].

The relative contributions of these new mechanisms can be studied[5, 4, 1] by changing the initial energy and the quantum numbers in the initial state by choosing different projectiles (α , ^{3}He , t , d , p). In this respect inelastic (d, d') scattering[6] is as promising as the (α, α') reaction with coherent pion production[5], and excitation of $N^*(1440)$ resonance[7]. For example, excitation of Δ "in the target" is forbidden by isospin conservation both in the (d, d') and (α, α') reactions: Δ can be excited only "in the projectile"[5, 1].

New valuable information may come from studies of polarization effects when broad hadronic resonances are excited in the nuclear medium[1]. A difference between proton and nuclear targets could be expected[8] for some polarization observables, but there are only a few experiments exploring the polarization observables of these reactions[9]. An example of the use of the (d, d') reaction with polarized deuterons as a filter of specific reaction mechanisms was demonstrated in ref.[10].

The experiment was performed at the Laboratory for High Energies at the Joint Institute for Nuclear Research (JINR) using tensorially polarized deuteron beam of the Synchrophasotron. The ALPHA setup (shown in Fig.1) was used in the same configuration as in a concurrent investigation of deuteron breakup and the $p(d, p)d$

backward elastic scattering [1]. The (d, d') data were taken before and after the data for T_{20} in $\mu(d, p)d$ at $\theta_{d'}$ = 180°; the systematic uncertainty of the final results was estimated using this circumstance.

The sign of the beam polarization was changed in a cyclic fashion, "burst-after burst", as (0, -+, +), where "0" means absence of polarization, "-+" and "+" correspond to the sign of the $P_{zz} \approx \sqrt{2} p_{20}$; the quantization axis was perpendicular to the plane containing the mean beam orbit in the accelerator. The polarization of the beam was measured with the ALPHA polarimeter[12] before and after data taking; averaged values of the polarization are: $P_{zz}^{(-)} = -0.822 \pm 0.007 \pm 0.008$ and $P_{zz}^{(+)} = +0.800 \pm 0.011 \pm 0.021$ where both statistical and systematical uncertainties are shown. Other beam parameters (positions and widths of the beam spot at control points) together with parameters of the machine, were monitored by the beam control system of the accelerator. They were transmitted to our on-line computer after each burst and were used in the off-line analysis.




Fig.1 ALPHA spectrometer at the VPI beam line. PC: multi-wire proportional chambers; S01-02, S11-12, S31-32, S41-42: scintillation counters of the TOF system; S1, S2, SAB1-2: trigger scintillation counters; M2: the analysing magnet. The TOF-base between S11-12 and S41-42 was about 50 m.

The 30 cm liquid hydrogen (${}_1^1H_2$) or 5.7 cm carbon (${}^{12}C$) targets (T) were placed at the focus F3 of the slowly extracted beam. Two beam intensity monitors were placed upstream from the target: an ionization chamber and a scintillation counter operated in the counting mode but at lowered high voltage in order to avoid overloading at intensities (10^7 deuterons per burst or higher).

The deuterons scattered at 0° and the unscattered part of the primary beam entered the beam line VP1; dipole magnets placed between the foci F3 and F4 (not shown in Fig.1) removed the unscattered part of the beam while the scattered deuterons were transported further to the ALPHA-spectrometer (Fig.1). The momentum acceptance

$\Delta p/p \approx 4.5\%$ of the setup was mostly determined by the sizes of the trigger scintillation counters S1, S2, S3 and SAB1,2. The measurements were performed in several steps by changing the spectrometer and the VPI beam line tuning for a continuous coverage of the momentum spectrum. At every setting the intensity of the primary beam was optimized for a reasonable counting rates in the counters at the F1 point; the typical intensity varied from 10^7 to $5 \cdot 10^8$ per burst with a spill duration of ≈ 400 msec. At each setting data were taken with both the $LH2$ and C target; at most settings data were also taken without target, the "no target" background was not more than 10% at $Q > 0.6$ GeV and less than 50% at $Q \approx 0.1$ GeV. Here $Q := E_d - E_p$ is the transferred energy, E_d is the energy of the projectile and E_p the energy of the scattered deuteron.

The momentum of the detected particles was measured with accuracy $\sigma_p/p \approx 0.25\%$. The final resolution of the TOF system after all corrections was $\sigma_{TOF} \approx 0.21$ nsec. Particle identification was performed using momentum and TOF information; this resolution provided a complete separation between protons, pions and deuterons.

The spectrometer accepted a symmetrical over azimuthal angle for deuterons detected at 90° . When the spin quantization axis is perpendicular to the particles momenta, the cross section expressed [13] in terms of spherical tensor analyzing powers contains only the T_2 -term, as the contribution with $T_{1,3}$ is zero for reasons of symmetry. Therefore T_{20} can be calculated directly from the numbers n_\pm of "good" events detected for the "+" and "-" modes of the beam polarization, normalized to the corresponding monitor numbers (the last part of the Eq.(1) is valid when $|\rho_{20}^{(t)}| \approx |\rho_{20}^{(s)}|$):

$$T_{20} = \frac{2(n_+ + n_-)}{\rho_{20}^{(t)} n_+ + \rho_{20}^{(s)} n_-} = \frac{1}{\rho_{20}^{(t)} + \rho_{20}^{(s)}} \cdot \frac{n_+ + n_-}{n_+ + n_-} \quad (1)$$

The values of T_{20} were obtained separately for both parts of the run as mentioned above. Because several overlapped setup settings were available at each initial energy and for each target, further checks of systematics could be made. We estimate the overall systematical uncertainty in T_{20} as $\sigma_{T_{20},syst} \approx 0.05$ which is about the same size as the typical statistical uncertainty.


The final data for T_{20} are presented in Fig.2 as a function of Q ; the data points shown are weighted averages (with weights $1/\sigma_{T_{20}}$) from both parts of the run.

The systematic uncertainty of the "zero point" of the Q scale was $\sigma_{Q,syst} \approx 20$ MeV; as a consequence the data shown in Fig.2 might be shifted as a whole within this corridor. The resolution on Q was $\sigma_Q \approx 15$ MeV.

The main features of the data are the following:

- 1) T_{20} is negative and increases (almost linearly) in absolute value when Q increases.

Fig.2 Lense-matrix elements of the $p(d,d';\Delta)$ (open squares) and $p(d,d';\bar{\Lambda})$ (full circles) inelastic scattering at 0° versus the energy transfer Q defined in the text.



- 2) T_{20} is small and compatible with zero in the region of coherent pion production where the 4-momentum transfer squared is small, $|t| \lesssim 0.05 \text{ GeV}^2/c^2$. It significantly differs from zero in the region of the $N(1440)$ resonance excitation and above.
- 3) $|T_{20}|$ becomes relatively big at large Q (about 0.4 to 0.6 at $Q \simeq 1 \text{ GeV}$).
- 4) There is no visible difference between the $p(d, d')X$ and $C(d, d')X$ data at 0° .

These features indicate that T_{20} in the inelastic $(d, d')X$ scattering at 0° is not highly sensitive to nuclear medium effects in excitation of broad nucleonic resonances. Perhaps the deuteron formfactor determines most of the overall Q -dependence of T_{20} , as it appears to be the case for the general trend of the cross sections[6]. For example, the $p(d, d')$ scattering at low Q (i.e. low $|t|$) should be determined by the coherent pion production via the Δ excitation in the projectile[5]. Interpreting this process as a coherent scattering of a virtual pion[14] on the deuteron, one could explain the smallness of T_{20} at low Q as a result of the dominance of the S wave part in the deuteron formfactor at small $|t|$.

Independently of any interpretation, the data demonstrate that this reaction can be used for the polarimetry of tensorially polarized deuterons at high energies.

We are grateful to the JHE accelerator staff, the POLARIS team and A.D. Kirillov, P.A. Rykoyatkin, A.I. Svetov, L.B. Golovanov, Yu.T. Borzynov, A.P. Tsvinev for providing good conditions for the experiment, L.N. Strunov, A.A. Nomofilov and V.I. Sharov for measurements of the polarization of the primary deuteron beam; L.V. Budkin, A.I. Obukhovsky, Z.P. Motina and O.N. Piskunov for their help during the experiment and preparation the manuscript.

The research described in this publication was supported in part by grants: № 93-02-3961 of the Russian Foundation for Fundamental Research, № RFU000 from the International Science Foundation, № RFU300 from the International Science Foundation and Russian Government, PHY-94-11620 from the US National Science Foundation and DE-FG05 89ER40525 from the US Department of Energy.

References

- [1] E.A.Strokovsky, F.A.Gareev, JINR preprint E1-94-241 (1994) and references therein; Yad. Fiz. 58 (1995), in print.
- [2] V.G.Ableev et al, JETP Lett., 40 (1984) 763; JINR preprint E1-83-486 (1983); Phys. Lett. B264 (1991) 264; C.Ellegaard et al, Phys. Rev. Lett. 50 (1983) 1745;

E.A.Strokovsky et al, Phys. Part. & Nucl. 24 (1993) 255 and references therein.

- [3] S.A.Avramenko et al, JINR preprint P1-91-240 (1991); JINR Rapid Comm. 3[54]-92 (1992) 13; *ibid* 6[63]-93 (1993) 5; T.Nagae et al, Phys. Lett. B191 (1987) 31; J.Chiba et al, Nucl. Phys. A478 (1988) 491c; Phys. Rev. Lett. 67 (1991) 1982; M.Roy-Stephan et al, Nucl. Phys. A553 (1993) 209c; S.A.Avramenko et al, in: Proc. of the Int. Conf. Mesons & Nuclei at Intermediate Energies (Dubna, May 3-7, 1994) (in print); B.Rainstein et al, *ibid* and preprint IPNO-DRE.94-10 (1994); R.Gilman et al, Nucl. Phys. A577 (1994) 227c.
- [4] F.A.Gareev et al, Yad. Fiz. 57 (1994) 1020; *ibid* 1401.
- [5] P.Fernandez de Cordoba, E.Oset, Nucl. Phys. A544 (1992) 793; P.Fernandez de Cordoba et al, Phys. Lett. B319 (1993) 416; Nucl. Phys. A586 (1995) 586.
- [6] V.G.Ableev et al, Yad. Fiz. 37 (1983) 348 (carbon target); see also J.Banaigs et al, Phys. Lett. A53B (1973) 535 (proton target).
- [7] H.P.Morsch et al, Phys. Rev. Lett. 69 (1992) 1436; Nucl. Phys. A553 (1993) 645c.
- [8] T.Udagawa et al, Phys. Rev. C49 (1994) 3162; B.Koerfgen et al, Phys. Rev. C50 (1994) 1637; V.Dmitriev, Nucl. Phys. A577 (1994) 249c.
- [9] G.Glass et al, Phys. Lett. B129 (1983) 365; A.B.Wicklund et al, Phys. Rev. D34 (1986) 49; C.Ellegaard et al, Phys. Lett. 231 (1989) 365; D.Prout et al, Nucl. Phys. A577 (1994) 233c.
- [10] E.Tomas Gustafsson et al, Saclay preprint LNS/P/92-05 (1992); M.Morlet et al, Phys. Rev. C46 (1992) 1008; C.Furget et al., Phys.Rev. C51 (1995) 1562.
- [11] L.S.Azhigirey et al, in: Proc. of the 14-th IUPAP Conf. on Few Body Probl. in Phys. (Williamsburg, May 26-31, 1991) ed. F.Gross (The Coll. of William and Mary & CEBAF, 1991) p.18 and p.22; JINR preprints E1-94-155, E1-94-156 (1994).
- [12] V.G.Ableev et al, NIM A306 (1991) 73.
- [13] Proceedings of the 3-d International Symposium (Madison, 1970); ed. H.H.Barschall, W.Haeberli, (Univ. of Wisconsin Press, Madison, 1970).
- [14] T.E.O.Ericson, Nucl. Phys. A560 (1993) 458; 1075E.

Received by Publishing Department
on June 21, 1995.

Принимается подписка на препринты, сообщения Объединенного института ядерных исследований и «Краткие сообщения ОИЯИ».

Установлена следующая стоимость подписки на 12 месяцев на издания ОИЯИ, включая пересылку, по отдельным тематическим категориям:

Индекс	Тематика	Цена подписки на год
1.	Экспериментальная физика высоких энергий	22600 р.
2.	Теоретическая физика высоких энергий	59200 р.
3.	Экспериментальная нейтронная физика	7800 р.
4.	Теоретическая физика низких энергий	23400 р.
5.	Математика	14800 р.
6.	Ядерная спектроскопия и радиохимия	12000 р.
7.	Физика тяжелых ионов	2200 р.
8.	Криогеника	1400 р.
9.	Ускорители	12200 р.
10.	Автоматизация обработки экспериментальных данных	12200 р.
11.	Вычислительная математика и техника	14300 р.
12.	Химия	1200 р.
13.	Техника физического эксперимента	21300 р.
14.	Исследования твердых тел и жидкостей ядерными методами	7200 р.
15.	Экспериментальная физика ядерных реакций при низких энергиях	2600 р.
16.	Дозиметрия и физика защиты	2200 р.
17.	Теория конденсированного состояния	12200 р.
18.	Использование результатов и методов фундаментальных физических исследований в смежных областях науки и техники	1800 р.
19.	Биофизика	1800 р.
	«Краткие сообщения ОИЯИ» (5—6 выпусков)	15000 р.

Подписка может быть оформлена с любого месяца года.

Организациям и лицам, заинтересованным в получении изданий ОИЯИ, следует перевести (или отправить по почте) необходимую сумму на расчетный счет 000608905 Дубненского филиала ММКБ, г.Дубна Московской области, п/инд. 141980 МФО 211844, указав: «За подписку на издания ОИЯИ».

Во избежание недоразумений необходимо уведомить издательский отдел о произведенной оплате и вернуть «Карточку подписчика», отметив в ней номера и названия тематических категорий, на которые оформляется подписка, по адресу:

141980 г. Дубна Московской обл.
ул. Жолио Кюри, 6
ОИЯИ, издательский отдел

SUBJECT CATEGORIES OF THE JINR PUBLICATIONS

Index	Subject
1.	High energy experimental physics
2.	High energy theoretical physics
3.	Low energy experimental physics
4.	Low energy theoretical physics
5.	Mathematics
6.	Nuclear spectroscopy and radiochemistry
7.	Heavy ion physics
8.	Cryogenics
9.	Accelerators
10.	Automatization of data processing
11.	Computing mathematics and technique
12.	Chemistry
13.	Experimental techniques and methods
14.	Solid state physics. Liquids
15.	Experimental physics of nuclear reactions at low energies
16.	Health physics. Shieldings
17.	Theory of condensed matter
18.	Applied researches
19.	Biophysics

Ажгирей Л.С. и др.

E1-95-263

Первые измерения тензорной анализирующей способности T_{20} неупругого $(d, d')X$ -рассеяния под 0° на 1H и ^{12}C при 4,5 и 5,5 ГэВ/с

Представлены первые результаты измерения T_{20} для неупругой реакции $(d, d')X$ выше порога Δ изобары. Обнаружено, что T_{20} отрицательна, ее абсолютная величина невелика в области Δ изобары и растет почти линейно к $|T_{20}| \simeq (0,4 - 0,6)$ при передаче энергии $Q \simeq 1$ ГэВ. Значимой зависимости от типа мишени и начальной энергии не замечено.

Работа выполнена в Лаборатории высоких энергий ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна, 1995

Azhgirey L.S. et al.

E1-95-263

First Measurement of the Tensor Analysing Power T_{20}
in Inelastic $(d, d')X$ -Scattering at 0° on 1H and ^{12}C
at 4.5 and 5.5 GeV/c

First results for T_{20} of the inelastic $(d, d')X$ reaction above the Δ -threshold are presented. It is observed that T_{20} is negative, small in absolute value in the Δ region and rises almost linearly to $|T_{20}| \simeq (0.4 - 0.6)$ at energy transfers $Q \simeq 1$ GeV. No significant dependence on the type of the target and on the initial energy was observed.

The investigation has been performed at the Laboratory of High Energies, JINR.

Preprint of the Joint Institute for Nuclear Research. Dubna, 1995

Макет Т.Е.Попеко

Подписано в печать 11.07.95
Формат 60×90/16. Офсетная печать. Уч.-изд.листов 0,8
Тираж 410. Заказ 48391. Цена 480 р.

Издательский отдел Объединенного института ядерных исследований
Дубна Московской области