

CERN-EP-2024-268
2024/11/19

CMS-BPH-21-002

Angular analysis of the $B^0 \rightarrow K^*(892)^0 \mu^+ \mu^-$ decay in proton-proton collisions at $\sqrt{s} = 13$ TeV

The CMS Collaboration*

Abstract

A full set of optimized observables is measured in an angular analysis of the decay $B^0 \rightarrow K^*(892)^0 \mu^+ \mu^-$ using a sample of proton-proton collisions at $\sqrt{s} = 13$ TeV, collected with the CMS detector at the LHC, corresponding to an integrated luminosity of 140 fb^{-1} . The analysis is performed in six bins of the squared invariant mass of the dimuon system, q^2 , over the range $1.1 < q^2 < 16 \text{ GeV}^2$. The results are among the most precise experimental measurements of the angular observables for this decay and are compared to a variety of predictions based on the standard model.

Submitted to Physics Letters B

1 Introduction

The $b \rightarrow s\ell^+\ell^-$ flavor-changing neutral-current processes are strongly suppressed according to the standard model (SM) of particle physics, since they only proceed through electroweak loop-level diagrams. Since virtual contributions of physics beyond the SM could occur with comparable amplitudes and alter the decay properties, these transitions are an excellent environment for indirect new physics searches. Among these processes, the $B^0 \rightarrow K^{*0}(K^+\pi^-)\mu^+\mu^-$ decay, where K^{*0} indicates the $K^*(892)^0$ meson, has long been considered a promising channel, thanks to its rich kinematical structure defined by the angular momenta of the final state particles. Compared to other well known decay modes, as for example $B_s^0 \rightarrow \mu^+\mu^-$, where only the branching ratio and mean life are measured [1–4], it provides access to a larger set of observables with high sensitivity to deviations from SM expectations. An extensive program of measuring angular observables from this decay has been carried out by the BaBar [5], Belle [6, 7], and CDF [8] Collaborations. The data collected during Run 1 of the CERN LHC (2011–2012) were used by the LHCb [9–11], CMS [12–14], and ATLAS [15] Collaborations to investigate this decay mode. Since the outcomes of those measurements were statistically limited, the additional data collected during Run 2 (2015–2018) can be used to significantly reduce the experimental uncertainties. The most recent model-independent result from the LHCb Collaboration [16], which includes data from Run 1 and 2016, confirms their previous measurement and appears to exhibit tension with some of the predictions based on the SM. These predictions, while all based on the SM, yield different values with varying uncertainties. The largest differences in the calculations arise from the estimation of the non-local hadronic matrix elements and the evaluation of the corresponding uncertainty [17–19].

This Letter contains the first CMS measurement of a complete set of optimized observables averaged over charge-conjugation and parity (CP) of the $B^0 \rightarrow K^{*0}\mu^+\mu^-$ decay. The results are obtained from the Run 2 data set collected in 2016–2018 at a center-of-mass energy of 13 TeV, corresponding to an integrated luminosity of 140 fb^{-1} . The analysis is performed in bins of the invariant mass squared of the dimuon system, q^2 , ranging from 1.1 to 16 GeV^2 . Tabulated results are provided in the HEPData record for this analysis [20]. Throughout this Letter, charge-conjugate states are implied unless explicitly stated otherwise.

2 Decay rate and angular analysis observables

The decay $B^0 \rightarrow K^{*0}\mu^+\mu^-$, where the K^{*0} meson decays to $K^+\pi^-$, can be described by q^2 and three decay angles: θ_l , θ_K , and ϕ . The definition of the angles is consistent with the one used in experimental publications and detailed in Ref. [9]. A sketch representing the definition of the angles is shown in Fig. 1. The angle θ_l is defined as the angle between the positive (negative)




Figure 1: Sketch representing the definition of the angles θ_l (left), θ_K (center), and ϕ (right).

muon momentum and the direction opposite to the B^0 (\bar{B}^0) momentum in the dimuon rest frame; the angle θ_K is defined as the angle between the kaon momentum and the direction opposite to the B^0 (\bar{B}^0) momentum in the K^{*0} (\bar{K}^{*0}) rest frame; the angle ϕ is the angle between the plane containing the μ^+ and μ^- and the plane containing the kaon and the pion from the K^{*0} decay in the B^0 rest frame.

Following the notation of Ref. [21] and the approach further proposed in Refs. [22, 23], the CP -averaged angular decay rate of the $B^0 \rightarrow K^{*0} \mu^+ \mu^-$ decay can be written as:

$$\begin{aligned} \Gamma_P \equiv \frac{1}{d\Gamma/dq^2} \frac{d^4\Gamma}{dq^2 d\cos\theta_l d\cos\theta_K d\phi} &= \frac{9}{32\pi} \left[\frac{3}{4}(1 - F_L) \sin^2 \theta_K + F_L \cos^2 \theta_K \right. \\ &+ \left. \left(\frac{1}{4}(1 - F_L) \sin^2 \theta_K - F_L \cos^2 \theta_K \right) \cos 2\theta_l \right. \\ &+ \frac{1}{2} P_1 (1 - F_L) \sin^2 \theta_K \sin^2 \theta_l \cos 2\phi \\ &+ \sqrt{(1 - F_L) F_L} \left(\frac{1}{2} P'_4 \sin 2\theta_K \sin 2\theta_l \cos \phi + P'_5 \sin 2\theta_K \sin \theta_l \cos \phi \right) \\ &- \sqrt{(1 - F_L) F_L} \left(P'_6 \sin 2\theta_K \sin \theta_l \sin \phi - \frac{1}{2} P'_8 \sin 2\theta_K \sin 2\theta_l \sin \phi \right) \\ &\left. + 2P_2 (1 - F_L) \sin^2 \theta_K \cos \theta_l - P_3 (1 - F_L) \sin^2 \theta_K \sin^2 \theta_l \sin 2\phi \right], \end{aligned} \quad (1)$$

where F_L is the fraction of longitudinal polarization of the K^{*0} meson, and $P_{1,2,3}$ and $P'_{4,5,6,8}$ are observables that have been optimized to reduce their dependence on the values and uncertainties of the form factors. In the publications of the LHCb Collaboration [11, 16] the definition of some of the $P_i^{(\prime)}$ observables is different, namely $P_4^{\text{CMS}} = 2P_4^{\text{LHCb}}$, $P_6^{\text{CMS}} = -P_6^{\text{LHCb}}$, and $P_8^{\text{CMS}} = -2P_8^{\text{LHCb}}$.

In order to also describe $B^0 \rightarrow K^+ \pi^- \mu^+ \mu^-$ events in which the $K^+ \pi^-$ system can be in an S -wave instead of P -wave configuration, the following extension of the angular decay rate (as in Eq. (44) of Ref. [23]) is used:

$$\begin{aligned} \frac{1}{d\Gamma/dq^2} \frac{d^4\Gamma}{dq^2 d\cos\theta_l d\cos\theta_K d\phi} &= (1 - F_S) \Gamma_P + \frac{3}{16\pi} (F_S \sin^2 \theta_l + A_S \sin^2 \theta_l \cos \theta_K \\ &+ A_S^4 \sin \theta_K \sin 2\theta_l \cos \phi + A_S^5 \sin \theta_K \sin \theta_l \cos \phi \\ &+ A_S^7 \sin \theta_K \sin \theta_l \sin \phi + A_S^8 \sin \theta_K \sin 2\theta_l \sin \phi), \end{aligned} \quad (2)$$

where the term proportional to F_S is the S -wave component, and the terms proportional to the A_S and $A_S^{4,5,7,8}$ coefficients describe the interference between the S - and P -wave terms.

3 The CMS detector and event samples

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter, and a brass and scintillator hadron calorimeter, each composed of a barrel and two endcap sections. Forward calorimeters extend the pseudorapidity (η) coverage provided by the barrel and endcap detectors. Muons are measured in the range $|\eta| < 2.4$, with detection planes made using three technologies: drift tubes, cathode strip chambers, and resistive-plate chambers. The single-muon trigger efficiency exceeds 90% over the full η range, and the efficiency to reconstruct and identify muons

is greater than 96%. Matching muons to tracks measured in the silicon tracker results in a relative transverse momentum (p_T) resolution, for muons with p_T up to 100 GeV, of 1% in the barrel and 3% in the endcaps [24]. The silicon tracker used in 2016 measured charged particles within the range $|\eta| < 2.5$. For nonisolated particles of $1 < p_T < 10$ GeV and $|\eta| < 1.4$, the track resolutions were typically 1.5% in p_T and 25–90 (45–150) μm in the transverse (longitudinal) impact parameter [25]. At the start of 2017, a new pixel detector was installed [26]; the upgraded tracker measured particles up to $|\eta| = 3.0$ with typical resolutions of 1.5% in p_T and 20–75 μm in the transverse impact parameter [27] for nonisolated particles of $1 < p_T < 10$ GeV. A more detailed description of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, can be found in Ref. [28].

Events of interest are selected using a two-tiered trigger system. The first level, composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz within a fixed latency of about 4 μs [29]. The second level, known as the high-level trigger, consists of a farm of processors running a version of the full event reconstruction software optimized for fast processing, and reduces the event rate to around 1 kHz before data storage [30]. All events used in this analysis are selected by a set of triggers demanding two identified muons of opposite charges and an additional track to form a common vertex, displaced from the region of the proton-proton interactions (beamspot). The trigger required each muon to have $p_T > 4$ GeV, $|\eta| < 2.5$, and to pass within 2 cm of the beam axis. The dimuon system was required to have $p_T > 6.9$ GeV, a vertex fit χ^2 probability larger than 10%, a separation of the vertex relative to the beamspot in the transverse plane larger than 3 times the uncertainty of the separation, and the cosine of the angle in the transverse plane between the dimuon momentum vector and the vector from the beamspot to the dimuon vertex greater than 0.9. The additional track was required to have $p_T > 0.8$ (1.2) GeV and a minimum distance of closest approach from the beamspot greater than 0 (2) times the uncertainty for data collected in 2016 (2017–2018). As the trigger configuration was restricted in the q^2 range, the analysis is limited to events with $q^2 < 16$ GeV².

Monte Carlo (MC) simulations are used to determine the impact of the detector geometry, trigger, reconstruction, and selection on the angular distribution of the signal, and to validate the analysis procedure. Simulated events are generated with PYTHIA 8.230 [31], decays of particles containing b or c quarks are simulated with EVTGEN 1.6.0 [32], and final-state radiation is included via PHOTOS 3.61 [33, 34]. Events are passed through the CMS detector simulation based on GEANT4 [35] and simulated minimum bias events are added to mimic the number of multiple vertices per event as seen in the data.

4 Event reconstruction, selection, and efficiency modeling

The offline reconstruction requires two oppositely charged muons and two oppositely charged tracks. Muons must satisfy general identification (soft muon) criteria [24] and match the muons used in the trigger. The p_T and η requirements applied by the trigger must be satisfied by the offline muons as well. The two tracks are required to have $p_T > 0.8$ GeV, $|\eta| < 2.4$, a distance of closest approach to the beamspot greater than 0.8 times the uncertainty, and to pass high-quality [25] identification criteria. One of the two tracks must match the track used in the trigger. In the 2017 and 2018 data sets, the trigger-matched track is required to have $p_T > 1.2$ GeV and the distance of the closest approach to the beamspot greater than twice the uncertainty. A veto is applied to reject tracks that are also reconstructed and identified as muons. The two tracks are fitted to a common vertex to form the K^{*0} candidate. The K^{*0} meson invariant mass is computed for both the $K^+\pi^-$ and $K^-\pi^+$ mass hypotheses, and at least one

of the two combinations is required to lie within 150 MeV of the nominal K^{*0} meson mass [36]. A $\phi(1020)$ veto is applied, demanding that the invariant mass of the two-track system, when assigning the kaon mass to both, is larger than 1.035 GeV.

The B^0 candidates are formed by combining the two oppositely charged muons with a K^{*0} candidate. The four daughter tracks of the B^0 candidate must originate from a common vertex, with vertex fit χ^2 probability greater than 1%. The momenta of the four tracks are then refitted using the common vertex as a constraint to improve the resolution of the track parameters. The flavor of the B^0 and K^{*0} candidates, and hence the mass assigned to the two hadronic tracks, is determined by the mass assignment that gives a K^{*0} meson mass closest to its nominal value. Based on simulation studies, this criterion results in incorrect flavor assignment for 12–14% of the candidates, depending on the data-taking year and the q^2 region.

A multivariate analysis is used to optimize the background rejection. A boosted decision tree is trained on background data events from the mass sidebands and simulated signal events, separately for each year of data-taking. The input variables of the discriminator are: the χ^2 probability of the B^0 candidate decay vertex; the transverse distance between the B^0 candidate decay vertex and the beamspot divided by its uncertainty; the cosine of the angle between the B^0 candidate momentum and the distance vector from the beamspot to the B^0 candidate decay vertex, in the transverse plane; the transverse B^0 candidate impact parameter with respect to the beamspot divided by its uncertainty; the transverse impact parameter of each hadronic track with respect to the beamspot divided by its uncertainty; the K^{*0} candidate invariant mass; and the sum of the final-state particles relative isolation, defined as the sum of the p_T of all the high-purity tracks having $p_T > 0.8$ GeV within a cone of $\Delta R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} < 0.4$ centered on the particle, divided by the p_T of the particle. Candidates with a decision tree score above a threshold are retained for analysis, with the threshold set to optimize the expected signal significance.

To reduce the contamination from $B^+ \rightarrow K^+ \mu^+ \mu^-$, including the dimuon decays of J/ψ and $\psi(2S)$ mesons, a veto is imposed on the invariant mass of the two possible $K^+ \mu^+ \mu^-$ triplets, where K^+ is, in turn, one of the two hadronic tracks from the $K^+ \pi^- \mu^+ \mu^-$ candidate, with the charged kaon mass assigned. The event is discarded if the mass of either $K^+ \mu^+ \mu^-$ combination is within a region corresponding to three times the $K^+ \mu^+ \mu^-$ mass resolution around the B^+ meson mass. This selection suppresses the contamination from these decays to a negligible level.

The J/ψ and $\psi(2S)$ meson control samples are defined by the ranges of the dedicated q^2 bins, 8.68–10.09 GeV² and 12.86–14.18 GeV², and the requirement $|q - m_{J/\psi}| < 3\sigma_q$ and $|q - m_{\psi(2S)}| < 3\sigma_q$, respectively. Here σ_q is the per-event uncertainty in the dimuon invariant mass q , and m represents the nominal mass of the charmonium state. Events satisfying these requirements are excluded from the signal q^2 bins. However, a fraction of events from the resonant channels still appears in the adjoining q^2 bins, mainly due to an unreconstructed photon from the charmonium decay. This contamination is reduced by applying combined requirements on q and the B^0 candidate invariant mass, which are tuned independently for the dimuon mass regions below the J/ψ meson mass, between the J/ψ and the $\psi(2S)$ meson masses, and above the $\psi(2S)$ meson mass. These requirements are set so that less than 10% of the background events originate from the resonant channels, by using simulated events to measure the remaining resonant contamination. For $q < m_{J/\psi}$ ($q > m_{J/\psi}$), candidates must satisfy the condition $|(m - m_{B^0}) - (q - m_{J/\psi})| > 190$ (90) MeV, where m is the invariant mass of the B^0 candidate. For $q < m_{\psi(2S)}$ ($q > m_{\psi(2S)}$), candidates must satisfy the condition $|(m - m_{B^0}) - (q - m_{\psi(2S)})| > 70$ (80) MeV. In the 2016 data sample, these requirements are

tightened to 200, 100, 80, and 110 MeV. In addition, B^0 candidates formed by the final-state particles from the $B^0 \rightarrow J/\psi K^{*0}$ decay, where the identification of a muon and a hadron track is inverted, are rejected with dedicated requirements obtained from simulation. Two alternative B^0 candidates, denoted as “alt”, are formed by reversing the identification of the same-sign muon and hadron track pairs, and the one giving a K^{*0} mass closest to its nominal value is selected (B_{alt}^0). Requirements on $m(K_{\text{alt}}^{*0})$, q_{alt} , $m(B_{\text{alt}}^0)$, $\Delta R(\mu, \text{track})$, and $\Delta p_{\text{T}}(\mu, \text{track})$ are placed so that 90% of the reversed candidates are rejected, with less than 1% impact on the signal efficiency. Peaking backgrounds from $B_s^0 \rightarrow \mu^+ \mu^- \phi(1020)$, $B_s^0 \rightarrow \mu^+ \mu^- K^{*0}$ and $B_s^0 \rightarrow \mu^+ \mu^- K^+ K^-$ decays are studied in simulation, using their equivalent decays with intermediate J/ψ and $\psi(2S)$ resonances. After the full reconstruction and selection chain, their contribution is negligible compared to the J/ψ resonant channel decay.

The contamination from $\Lambda_b^0 \rightarrow p K^- \mu^+ \mu^-$ decays is investigated directly with data by assigning the proton and kaon masses to the hadronic tracks of the $K^+ \pi^- \mu^+ \mu^-$ candidates, and searching for a peak at the nominal Λ_b^0 mass in the candidate invariant mass distribution. No evidence of such contamination is found.

Finally, partially reconstructed decays $B^+ \rightarrow K^+ \psi(2S) (\rightarrow J/\psi \pi^+ \pi^-)$ contribute to the low-mass sideband of the J/ψ control sample and present a narrow peak in the $\cos \theta_K$ distribution in the region $0.6 < \cos \theta_K < 0.8$. To reject this background, a selection is applied on $m(\mu \mu K)$, $m(\mu \mu \pi)$, $m(\mu \mu K_{\pi} \pi_K)$, and $m(K_{\pi} \pi_K)$ invariant masses in the events with $p_{\text{T}}(K) > p_{\text{T}}(\pi)$, where K_{π} and π_K are given the kaon and pion flavor assignment, respectively, based on which combination is closest to the K^{*0} mass, but are assigned the charged pion and kaon mass, respectively, in the invariant mass calculations.

The acceptance, as well as the efficiency to trigger, reconstruct, and select the signal and control channel events, are modeled with a function of the three angular variables, using simulated event samples, individually for each year of data-taking and q^2 bin. These functions are obtained by applying the nonparametric kernel density estimator method to the unbinned distributions of simulated events before and after imposing the acceptance and selection criteria previously described, and computing their ratio. This method is able to reproduce important features of the efficiency and has also proven to be robust even in regions of the phase space poorly populated by simulated events. Efficiencies are determined for both correctly identified (the kaon and pion have the correct charge) and misidentified (the kaon and pion charges are reversed) candidates, with both contributing as signal in the fit described below. Corrections derived from the data are applied to the simulated events to improve the modeling of the additional proton-proton interactions within the same or nearby bunch crossings and of those variables correlated with the decay angles. The latter are evaluated by training a classifier on the difference between simulation and data from the $B^0 \rightarrow J/\psi K^{*0}$ control sample. The background-subtracted data distributions have been obtained via the \mathcal{P} lot technique [37], where the reconstructed B^0 candidate invariant mass is used as the discriminating variable. The inputs to the classifier are the transverse momenta of the final-state particles, the B^0 candidate η , and the topological variables that are correlated to the angular ones. The output score of the classifier is used to reweight the simulation. The effectiveness of the method has been validated assessing the compatibility of the distributions of the input features in data and in the corrected simulation, on an independent sample of $B^0 \rightarrow J/\psi K^{*0}$ events, as well as on a sample of $B^0 \rightarrow \psi(2S) K^{*0}$ events.

5 Fit procedure

The angular observables are measured in each q^2 bin by performing an unbinned maximum-likelihood fit to the four-dimensional distribution of the invariant mass of the B^0 candidate, m , and the three angular variables: $\cos \theta_K$, $\cos \theta_l$, and ϕ . The bin definition is the same as in Ref. [14], except that the first bin starts from 1.1 GeV^2 rather than 1.0 GeV^2 to exclude the contamination from the $\phi(1020)$ resonance. All B^0 candidates with the invariant mass in the range $5 < m < 5.6 \text{ GeV}$ are included and a simultaneous fit is performed to data sets collected in 2016, 2017, and 2018.

The probability density function (PDF) used to fit each of these three data sets is:

$$\begin{aligned}
 P(m, \cos \theta_K, \cos \theta_l, \phi) = & Y_S \left[S^C(m) S^a(\cos \theta_K, \cos \theta_l, \phi) \epsilon^C(\cos \theta_K, \cos \theta_l, \phi) \right. \\
 & \left. + R S^M(m) S^a(-\cos \theta_K, -\cos \theta_l, -\phi) \epsilon^M(\cos \theta_K, \cos \theta_l, \phi) \right] \\
 & + Y_B B^m(m) B^a(\cos \theta_K, \cos \theta_l, \phi),
 \end{aligned} \tag{3}$$

where the first and second terms represent the contributions of the signal events for which the flavor has been correctly and wrongly identified, respectively, while the third term describes the background contribution. The Y_S and Y_B parameters represent the yield of signal and background events, respectively. The functions S^C and S^M describe the mass distribution of correctly identified and misidentified signal candidates, respectively, and they are parametrized, depending on the q^2 bin, by a Crystal Ball function [38] plus a Gaussian or a double Crystal Ball function [39] (with a single or double Gaussian kernel and two power-law tails in opposite directions). The function S^a is the angular decay rate as written in Eq. (2). The functions ϵ^C and ϵ^M represent the efficiency for correctly and misidentified events, respectively, as a function of the angular variables. The parameter R represents the ratio of the relative amount of misidentified events in data and the same value as computed on MC samples; the information of the latter value is contained in the relative value of ϵ^M with respect to ϵ^C , while R is introduced to account for differences between data and simulation. The function B^m represents the mass distribution of the combinatorial background events and consists of an exponential function; in the ranges $6 < q^2 < 8.68 \text{ GeV}^2$ and $10.09 < q^2 < 12.86 \text{ GeV}^2$, the exponential is multiplied by a function describing the reduction of background events from the requirements used to reduce the J/ψ and $\psi(2S)$ meson contamination. The function B^a represents the angular distribution of the background, which is modeled by a three-dimensional Bernstein polynomial determined from the data sidebands. The sidebands are regions of the $K^+ \pi^- \mu^+ \mu^-$ mass spectrum that do not contain the signal peak, and they are defined in such a way that the number of signal events contained in the sideband region is no more than 6% of the total number of events in that region. The Bernstein polynomial degree is chosen as the lowest value that produces a p -value larger than 5%: the degree varies between 2 and 6 for $\cos \theta_K$, between 1 and 5 for $\cos \theta_l$, and between 2 and 5 for ϕ , depending on the data-taking year and q^2 bin. The functions ϵ^C , ϵ^M , and B^a are constrained to have the same value at $\phi = \pm\pi$.

In the simultaneous fit, the parameters of the PDF corresponding to the angular observables are shared among the three data sets, while other parameters are independent to account for the effects of different data-taking conditions on the efficiency and resolution. The coefficients of the S -wave and its interference with the P -wave are treated as nuisance parameters and are shared among the three data sets. All parameters are free, except those of the background angular shape, which are fixed to the values obtained from the data sidebands as described above. The parameters of the signal mass shapes are constrained to the results of the fit to the

$K^+\pi^-\mu^+\mu^-$ mass distribution of simulated signal events through Gaussian likelihood factors. An additional Gaussian constraint is applied to the parameter R , with a mean equal to one and a width corresponding to the expected statistical fluctuation of R in the data, which varies from 5 to 20% depending on the data-taking year and the q^2 bin. The fit strategy comprises a procedure relying on the penalization of the likelihood function to ensure that the resulting angular observables lie in the physical region of the parameter space, i.e. provide a non-negative decay rate.

The best fit value of the parameters and approximate 68% CL confidence intervals are extracted following the procedure described in Section 3.2 of Ref. [40]. The validity of this method and its robustness to the presence of nonphysical regions of the parameter space have been proven by an extensive study of its statistical coverage. The fit procedure and results are validated through fits to pseudo-experiments, MC simulation samples, and control samples.

6 Systematic uncertainties

Several sources of systematic uncertainties are evaluated to account for possible biases introduced by the analysis procedure, differences between the signal features in data and simulation, and any unaccounted minor background sources.

The possible bias introduced by mismodeling the efficiency shape is evaluated using simulated events as the difference between the results from fitting the generated angular variable distribution of generated events and fitting the reconstructed angular variable distributions of reconstructed events incorporating the efficiency. Additionally, a systematic uncertainty is estimated to account for possible biases in the fit results when the fit procedure is performed in realistic conditions (presence of background, limited number of available events). A set of 100 independent samples is created with the signal obtained from MC simulation and the background generated from the corresponding PDF, with signal and background yields matched to the respective values estimated from the data. Each of the samples is fit with the standard procedure and the average value over all 100 samples is calculated for each parameter. The systematic uncertainty is obtained by subtracting the average value from the value measured in the fit to the full signal-only MC simulation sample.

To evaluate the uncertainty due to the misidentification fraction being measured using simulated samples, the Gaussian constraint on the parameter R is removed in the fit to the $B^0 \rightarrow J/\psi K^{*0}$ control sample, yielding an unconstrained measurement of R . Then, a set of alternative fits is performed, using the measured R values (1.07, 1.02, and 1.02 for 2016, 2017, and 2018, respectively) as the means for the Gaussian constraints in the signal fits.

The systematic uncertainty associated with possible differences in the mass resolution between data and simulation is evaluated by first measuring a resolution scale factor with the $B^0 \rightarrow J/\psi K^{*0}$ control sample and then applying the same scale factor in the fit to the signal bins. The difference with respect to the nominal result is taken as a systematic uncertainty.

Systematic uncertainties are assessed for the signal and background mass shapes. To evaluate the uncertainty related to the choice of the signal mass model, a fit is performed to the data using the correctly identified and misidentified mass spectra from MC simulation as templates for the S^C and S^M components of the PDF. To evaluate the uncertainty related to the choice of the background mass model, a fit is performed to the data using Bernstein polynomials. For the highest q^2 bin only, the exponential function corrected for the threshold effect at the low edge of the mass range was also used as an alternative function.

The efficiency is evaluated on a finite sample of simulated events. A set of efficiency functions reflecting the statistical variation of the simulated event sample is generated, and used to fit the data. The standard deviation of the results obtained on each angular parameter is included as a systematic uncertainty. As the angular parameters of the background function are fixed in the fit, the statistical uncertainty related to the background parameters is evaluated separately. This is done by varying the parameters using the covariance matrix of the fit to the sidebands to generate 100 different background functions. The uncertainty is obtained by taking the standard deviation of the results obtained by fitting the data with each of the alternative background functions.

The effect of data-to-simulation differences on the angular efficiency and resolution is evaluated by performing an additional fit with the F_L parameter fixed to the value obtained from the nominal fit plus the difference between the results of the fit to the $B^0 \rightarrow J/\psi K^{*0}$ control sample on the F_L parameter and its current world average [36]. The difference with respect to the nominal result is taken as a systematic uncertainty.

The bias introduced by the contamination from partially reconstructed background sources in the lower-mass sideband on the background angular shape is evaluated in the J/ψ control sample, excluding from the mass sidebands events with the B^0 candidate mass lower than 5.1 GeV to define the background angular shape. The difference between the new results and the nominal ones is considered as a systematic uncertainty. Finally, the uncertainty due to the possible contamination of events from the two resonant channels in the adjoining q^2 bins is evaluated by adding a component with a simulation-based shape to the PDF to describe the contribution of the resonant channels, and using the difference of the results as the systematic uncertainty.

The systematic uncertainties are determined for each q^2 bin, and in cases where the systematic uncertainty is obtained as the difference between a nominal and alternative configuration, the systematic uncertainty is symmetrized. Their impact on the various angular parameters is summarized in Table 1. The individual systematic uncertainties are considered independent and added in quadrature.

7 Results

The projections of the fitted PDF on the mass and angular distributions are shown in Fig. 2 for the $2 < q^2 < 4.3 \text{ GeV}^2$ and $4.3 < q^2 < 6 \text{ GeV}^2$ bins. The fit quality has been verified in all q^2 bins by assessing the agreement between the PDF projections and the data.

Table 2 and Fig. 3 provide the measured values of the CP -averaged observables in bins of q^2 along with the corresponding uncertainties. Figure 3 also contains four predictions based on the SM. The q^2 range of validity of the predictions varies depending on the methods used to estimate the form factors. The ABCDMN predictions are obtained using the theoretical framework discussed in Ref. [18], which combines local form factors derived from lattice quantum chromodynamics [41, 42] and light-cone sum rules [43], and non-local form factors from Ref. [44]. The EOS predictions are obtained using the procedure described in Ref. [45] and the EOS [46] software package, using the same sources as ABCDMN for the form-factor calculations, but with a different approach to combining them. Additional predictions are calculated using the `flavio` [47] software package, which combines local form factors from lattice quantum chromodynamics [41, 42] and light-cone sum rules [48], and quantum chromodynamics factorization for nonlocal form factors [49, 50]. The predictions calculated with the `HEPfit` [51] software package adopt a conservative estimation of non-local hadronic matrix

Table 1: The uncertainties considered in the analysis on the various angular observables. For each source of uncertainty, the range covers the absolute variation observed across the q^2 bins.

| Source | $F_L (\times 10^{-3})$ | $P_1 (\times 10^{-3})$ | $P_2 (\times 10^{-3})$ | $P_3 (\times 10^{-3})$ |
|-----------------------------|------------------------|------------------------|------------------------|------------------------|
| Efficiency modeling | 1–9 | 7–44 | 3–11 | 0–46 |
| Fit bias | 1–2 | 0–6 | 2–62 | 1–12 |
| Misidentification fraction | 0–2 | 1–4 | 1–3 | 0–14 |
| Signal mass resolution | 1–10 | 1–12 | 2–11 | 1–21 |
| Signal mass shape | 0–9 | 1–22 | 0–10 | 3–70 |
| Background mass shape | 0–5 | 1–16 | 1–13 | 0–8 |
| Efficiency (statistical) | 1–10 | 5–31 | 1–64 | 4–45 |
| Background (statistical) | 2–6 | 4–20 | 1–21 | 2–16 |
| Data/simulation differences | 8 | 0–23 | 0–16 | 0–13 |
| Partially reco background | 1 | 1 | 0 | 1 |
| Resonant background | 0–1 | 0–6 | 0–5 | 0–2 |

| Source | $P'_4 (\times 10^{-3})$ | $P'_5 (\times 10^{-3})$ | $P'_6 (\times 10^{-3})$ | $P'_8 (\times 10^{-3})$ |
|-----------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| Efficiency modeling | 3–87 | 2–13 | 5–16 | 6–28 |
| Fit bias | 9–54 | 0–8 | 0–3 | 0–24 |
| Misidentification fraction | 1–5 | 1–10 | 0–4 | 0–12 |
| Signal mass resolution | 4–23 | 0–12 | 0–5 | 0–16 |
| Signal mass shape | 2–16 | 1–15 | 0–7 | 0–91 |
| Background mass shape | 6–30 | 1–13 | 0–7 | 1–10 |
| Efficiency (statistical) | 5–47 | 4–22 | 4–13 | 10–59 |
| Background (statistical) | 6–37 | 4–24 | 3–9 | 5–23 |
| Data/simulation differences | 0–11 | 0–13 | 0–3 | 0–30 |
| Partially reco background | 25 | 0 | 0 | 2 |
| Resonant background | 0–30 | 0–11 | 0–5 | 0–12 |




Figure 2: Mass and angular distributions for (upper two rows) $2 < q^2 < 4.3 \text{ GeV}^2$ and (lower two rows) $4.3 < q^2 < 6 \text{ GeV}^2$. The projections of the total fitted distribution (blue solid line) and its different components are overlaid. The signal is shown by the red dotted line, and the background by the orange dashed line.

elements and their uncertainties, to account for a possible large impact from charm-loop penguin diagrams [17, 52].

Table 2: The measured CP -averaged angular observables, in the corresponding q^2 bins. The first uncertainty is statistical and the second is systematic.

| | $1.1 < q^2 < 2 \text{ GeV}^2$ | $2 < q^2 < 4.3 \text{ GeV}^2$ | $4.3 < q^2 < 6 \text{ GeV}^2$ |
|--------|-------------------------------------|--------------------------------------|--------------------------------------|
| F_L | $0.709^{+0.073}_{-0.054} \pm 0.021$ | $0.810^{+0.036}_{-0.030} \pm 0.016$ | $0.714^{+0.032}_{-0.030} \pm 0.012$ |
| P_1 | $0.09^{+0.23}_{-0.20} \pm 0.04$ | $-0.29^{+0.19}_{-0.21} \pm 0.05$ | $-0.30^{+0.15}_{-0.17} \pm 0.04$ |
| P_2 | $-0.37^{+0.17}_{-0.13} \pm 0.10$ | $-0.244^{+0.094}_{-0.077} \pm 0.039$ | $0.121^{+0.080}_{-0.076} \pm 0.030$ |
| P_3 | $-0.05^{+0.21}_{-0.22} \pm 0.04$ | $-0.19^{+0.20}_{-0.22} \pm 0.09$ | $-0.03 \pm 0.14 \pm 0.08$ |
| P'_4 | $-0.44^{+0.29}_{-0.32} \pm 0.11$ | $-0.43^{+0.16}_{-0.19} \pm 0.08$ | $-0.72^{+0.15}_{-0.16} \pm 0.07$ |
| P'_5 | $0.36^{+0.17}_{-0.13} \pm 0.03$ | $-0.14^{+0.10}_{-0.09} \pm 0.04$ | $-0.44 \pm 0.10 \pm 0.03$ |
| P'_6 | $0.000^{+0.094}_{-0.097} \pm 0.021$ | $0.108^{+0.075}_{-0.071} \pm 0.018$ | $0.129^{+0.074}_{-0.071} \pm 0.011$ |
| P'_8 | $0.16 \pm 0.37 \pm 0.11$ | $0.73^{+0.18}_{-0.19} \pm 0.06$ | $-0.01 \pm 0.22 \pm 0.04$ |
| | $6 < q^2 < 8.68 \text{ GeV}^2$ | $10.09 < q^2 < 12.86 \text{ GeV}^2$ | $14.18 < q^2 < 16 \text{ GeV}^2$ |
| F_L | $0.627 \pm 0.016 \pm 0.011$ | $0.474^{+0.011}_{-0.013} \pm 0.009$ | $0.394 \pm 0.012 \pm 0.009$ |
| P_1 | $-0.06 \pm 0.10 \pm 0.05$ | $-0.439^{+0.051}_{-0.047} \pm 0.030$ | $-0.465 \pm 0.037 \pm 0.025$ |
| P_2 | $0.188^{+0.039}_{-0.040} \pm 0.014$ | $0.386^{+0.021}_{-0.019} \pm 0.018$ | $0.440^{+0.008}_{-0.010} \pm 0.008$ |
| P_3 | $0.099^{+0.092}_{-0.090} \pm 0.014$ | $0.013^{+0.041}_{-0.043} \pm 0.007$ | $-0.034^{+0.037}_{-0.038} \pm 0.010$ |
| P'_4 | $-0.95 \pm 0.10 \pm 0.06$ | $-1.025^{+0.064}_{-0.066} \pm 0.059$ | $-1.159^{+0.042}_{-0.038} \pm 0.041$ |
| P'_5 | $-0.495 \pm 0.067 \pm 0.023$ | $-0.746^{+0.033}_{-0.032} \pm 0.014$ | $-0.688^{+0.038}_{-0.036} \pm 0.021$ |
| P'_6 | $0.010 \pm 0.052 \pm 0.016$ | $0.080^{+0.037}_{-0.041} \pm 0.011$ | $0.121^{+0.040}_{-0.039} \pm 0.011$ |
| P'_8 | $0.06 \pm 0.14 \pm 0.04$ | $0.09^{+0.09}_{-0.10} \pm 0.03$ | $0.011^{+0.089}_{-0.086} \pm 0.022$ |

The HEPfit predictions, given the large uncertainties, are compatible with the measurements. The EOS and ABCDMN predictions, while showing compatibility with the analysis results for most of the observables, present tensions with the measured values of P_2 and P'_5 , in the region of q^2 below the J/ψ resonance. The flavio predictions show similar tensions for the P'_5 parameter. The current measurements are compatible with those presented in previous analyses from the CMS Collaboration [13, 14] and other experiments [7, 11, 15, 16], as shown for P_2 and P'_5 in Fig. 4. The statistical uncertainty has significantly improved compared to previous CMS publications.

8 Summary

In summary, the study of the full angular distribution of the flavor-changing neutral-current $B^0 \rightarrow K^{*0} \mu^+ \mu^-$ decay has been performed using 140 fb^{-1} of proton-proton collision data recorded by the CMS detector at the LHC at $\sqrt{s} = 13 \text{ TeV}$. A complete set of observables has been measured via unbinned maximum likelihood fits to the B^0 candidate mass and angular variables, in bins of the squared invariant mass of the dimuon system ranging from 1.1 to 16 GeV^2 . The measurements are compared to a variety of predictions based on the standard model, with tension in a few of the angular observables seen for some of the predictions, as




Figure 3: Measurements of the angular observables versus q^2 . The inner vertical bars represent the statistical uncertainties, while the outer vertical bars give the total uncertainties. The horizontal bars show the bin widths. The vertical shaded regions correspond to the J/ψ and $\psi(2S)$ resonances. Predictions are shown, averaged in each bin, from Ref. [18] (labeled ABCDMN), and the EOS [46], flavio [47], and HEPfit [51] libraries.




Figure 4: Comparison of current measurements of P_2 and P_5' to a previous P_2 measurement by LHCb [16] and previous P_5' results from ATLAS [15], Belle [7], CMS [14], and LHCb [16]. The inner vertical bars represent the statistical uncertainties, while the outer vertical bars give the total uncertainties. The horizontal bars show the bin widths.

is also reported by other experiments. These results are among the most precise experimental measurements of the angular observables of this decay, and provide a valuable contribution to the understanding of the $b \rightarrow s\ell^+\ell^-$ processes.

Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centres and personnel of the Worldwide LHC Computing Grid and other centres for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: SC (Armenia), BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); ERC PRG, RVTT3 and MoER TK202 (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMBF, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LMTLT (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES and NSC (Poland); FCT (Portugal); MESTD (Serbia); MCIN/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI and NSTDA (Thailand); TUBITAK and TENMAK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (USA).

Individuals have received support from the Marie-Curie programme and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, 101115353, 101002207, and COST Action CA16108 (European Union); the Leventis

Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, project no. 22rl-037 (Armenia); the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the F.R.S.-FNRS and FWO (Belgium) under the "Excellence of Science – EOS" – be.h project n. 30820817; the Beijing Municipal Science & Technology Commission, No. Z191100007219010 and Fundamental Research Funds for the Central Universities (China); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Shota Rustaveli National Science Foundation, grant FR-22-985 (Georgia); the Deutsche Forschungsgemeinschaft (DFG), among others, under Germany's Excellence Strategy – EXC 2121 "Quantum Universe" – 390833306, and under project number 400140256 - GRK2497; the Hellenic Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Hungarian Academy of Sciences, the New National Excellence Program - ÚNKP, the NKFIH research grants K 131991, K 133046, K 138136, K 143460, K 143477, K 146913, K 146914, K 147048, 2020-2.2.1-ED-2021-00181, TKP2021-NKTA-64, and 2021-4.1.2-NEMZ.KI (Hungary); the Council of Science and Industrial Research, India; ICSC – National Research Centre for High Performance Computing, Big Data and Quantum Computing and FAIR – Future Artificial Intelligence Research, funded by the NextGenerationEU program (Italy); the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369 and 2021/43/B/ST2/01552 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MCIN/AEI/10.13039/501100011033, ERDF "a way of making Europe", and the Programa Estatal de Fomento de la Investigación Científica y Técnica de Excelencia María de Maeztu, grant MDM-2017-0765 and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, and the National Science, Research and Innovation Fund via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, grant B39G670016 (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (USA).

References

- [1] ATLAS Collaboration, "Study of the rare decays of B_s^0 and B^0 mesons into muon pairs using data collected during 2015 and 2016 with the ATLAS detector", *JHEP* **04** (2019) 098, doi:10.1007/JHEP04(2019)098, arXiv:1812.03017.
- [2] LHCb Collaboration, "Measurement of the $B_s^0 \rightarrow \mu^+ \mu^-$ decay properties and search for the $B^0 \rightarrow \mu^+ \mu^-$ and $B_s^0 \rightarrow \mu^+ \mu^- \gamma$ decays", *Phys. Rev. D* **105** (2022) 012010, doi:10.1103/PhysRevD.105.012010, arXiv:2108.09283.
- [3] LHCb Collaboration, "Analysis of neutral B-meson decays into two muons", *Phys. Rev. Lett.* **128** (2022) 041801, doi:10.1103/PhysRevLett.128.041801, arXiv:2108.09284.
- [4] CMS Collaboration, "Measurement of the $B_s^0 \rightarrow \mu^+ \mu^-$ decay properties and search for the $B^0 \rightarrow \mu^+ \mu^-$ decay in proton-proton collisions at $\sqrt{s} = 13$ TeV", *Phys. Lett. B* **842** (2023) 137955, doi:10.1016/j.physletb.2023.137955, arXiv:2212.10311.
- [5] BaBar Collaboration, "Angular distributions in the decays $B \rightarrow K^* \ell^+ \ell^-$ ", *Phys. Rev. D* **79** (2009) 031102, doi:10.1103/PhysRevD.79.031102, arXiv:0804.4412.

- [6] Belle Collaboration, “Measurement of the differential branching fraction and forward-backward asymmetry for $B \rightarrow K^{(*)}\ell^+\ell^-$ ”, *Phys. Rev. Lett.* **103** (2009) 171801, doi:10.1103/PhysRevLett.103.171801, arXiv:0904.0770.
- [7] Belle Collaboration, “Lepton-flavor-dependent angular analysis of $B \rightarrow K^*\ell^+\ell^-$ ”, *Phys. Rev. Lett.* **118** (2017) 111801, doi:10.1103/PhysRevLett.118.111801, arXiv:1612.05014.
- [8] CDF Collaboration, “Measurements of the angular distributions in the decays $B \rightarrow K^{(*)}\mu^+\mu^-$ at CDF”, *Phys. Rev. Lett.* **108** (2012) 081807, doi:10.1103/PhysRevLett.108.081807, arXiv:1108.0695.
- [9] LHCb Collaboration, “Differential branching fraction and angular analysis of the decay $B^0 \rightarrow K^{*0}\mu^+\mu^-$ ”, *JHEP* **08** (2013) 131, doi:10.1007/JHEP08(2013)131, arXiv:1304.6325.
- [10] LHCb Collaboration, “Measurement of form-factor-independent observables in the decay $B^0 \rightarrow K^{*0}\mu^+\mu^-$ ”, *Phys. Rev. Lett.* **111** (2013) 191801, doi:10.1103/PhysRevLett.111.191801, arXiv:1308.1707.
- [11] LHCb Collaboration, “Angular analysis of the $B^0 \rightarrow K^{*0}\mu^+\mu^-$ decay using 3 fb^{-1} of integrated luminosity”, *JHEP* **02** (2016) 104, doi:10.1007/JHEP02(2016)104, arXiv:1512.04442.
- [12] CMS Collaboration, “Angular analysis and branching fraction measurement of the decay $B^0 \rightarrow K^{*0}\mu^+\mu^-$ ”, *Phys. Lett. B* **727** (2013) 77, doi:10.1016/j.physletb.2013.10.017, arXiv:1308.3409.
- [13] CMS Collaboration, “Angular analysis of the decay $B^0 \rightarrow K^{*0}\mu^+\mu^-$ from pp collisions at $\sqrt{s} = 8\text{ TeV}$ ”, *Phys. Lett. B* **753** (2016) 424, doi:10.1016/j.physletb.2015.12.020, arXiv:1507.08126.
- [14] CMS Collaboration, “Measurement of angular parameters from the decay $B^0 \rightarrow K^{*0}\mu^+\mu^-$ in proton-proton collisions at $\sqrt{s} = 8\text{ TeV}$ ”, *Phys. Lett. B* **781** (2018) 517, doi:10.1016/j.physletb.2018.04.030, arXiv:1710.02846.
- [15] ATLAS Collaboration, “Angular analysis of $B_d^0 \rightarrow K^*\mu^+\mu^-$ decays in pp collisions at $\sqrt{s} = 8\text{ TeV}$ with the ATLAS detector”, *JHEP* **10** (2018) 047, doi:10.1007/JHEP10(2018)047, arXiv:1805.04000.
- [16] LHCb Collaboration, “Measurement of CP-averaged observables in the $B^0 \rightarrow K^{*0}\mu^+\mu^-$ decay”, *Phys. Rev. Lett.* **125** (2020) 011802, doi:10.1103/PhysRevLett.125.011802, arXiv:2003.04831.
- [17] M. Ciuchini et al., “Charming penguins and lepton universality violation in $b \rightarrow s\ell^+\ell^-$ decays”, *Eur. Phys. J. C* **83** (2023) 64, doi:10.1140/epjc/s10052-023-11191-w, arXiv:2110.10126.
- [18] M. Algueró et al., “To (b)e or not to (b)e: no electrons at LHCb”, *Eur. Phys. J. C* **83** (2023) 648, doi:10.1140/epjc/s10052-023-11824-0, arXiv:2304.07330.
- [19] M. Bordone, G. Isidori, S. Mächler, and A. Tinari, “Short- vs. long-distance physics in $B \rightarrow K^{(*)}\ell^+\ell^-$: a data-driven analysis”, *Eur. Phys. J. C* **84** (2024) 547, doi:10.1140/epjc/s10052-024-12869-5, arXiv:2401.18007.

-
- [20] HEPData record for this analysis, 2024. doi:10.17182/hepdata.154898.
- [21] W. Altmannshofer et al., “Symmetries and asymmetries of $B \rightarrow K^* \mu^+ \mu^-$ decays in the Standard Model and Beyond”, *JHEP* **01** (2009) 019, doi:10.1088/1126-6708/2009/01/019, arXiv:0811.1214.
- [22] S. Descotes-Genon, J. Matias, M. Ramon, and J. Virto, “Implications from clean observables for the binned analysis of $B \rightarrow K^* \mu^+ \mu^-$ at large recoil”, *JHEP* **01** (2013) 048, doi:10.1007/JHEP01(2013)048, arXiv:1207.2753.
- [23] S. Descotes-Genon, T. Hurth, J. Matias, and J. Virto, “Optimizing the basis of $B \rightarrow K^* \ell^+ \ell^-$ observables in the full kinematic range”, *JHEP* **05** (2013) 137, doi:10.1007/JHEP05(2013)137, arXiv:1303.5794.
- [24] CMS Collaboration, “Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **13** (2018) P06015, doi:10.1088/1748-0221/13/06/P06015, arXiv:1804.04528.
- [25] CMS Collaboration, “Description and performance of track and primary-vertex reconstruction with the CMS tracker”, *JINST* **9** (2014) P10009, doi:10.1088/1748-0221/9/10/P10009, arXiv:1405.6569.
- [26] CMS Tracker Group, “The CMS Phase-1 pixel detector upgrade”, *JINST* **16** (2021) P02027, doi:10.1088/1748-0221/16/02/P02027, arXiv:2012.14304.
- [27] CMS Collaboration, “Track impact parameter resolution for the full pseudo rapidity coverage in the 2017 dataset with the CMS Phase-1 Pixel detector”, CMS Detector Performance Summary CMS-DP-2020-049, 2020.
- [28] CMS Collaboration, “The CMS experiment at the CERN LHC”, *JINST* **3** (2008) S08004, doi:10.1088/1748-0221/3/08/S08004.
- [29] CMS Collaboration, “Performance of the CMS Level-1 trigger in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **15** (2020) P10017, doi:10.1088/1748-0221/15/10/P10017, arXiv:2006.10165.
- [30] CMS Collaboration, “The CMS trigger system”, *JINST* **12** (2017) P01020, doi:10.1088/1748-0221/12/01/P01020, arXiv:1609.02366.
- [31] T. Sjöstrand et al., “An introduction to PYTHIA 8.2”, *Comput. Phys. Commun.* **191** (2015) 159, doi:10.1016/j.cpc.2015.01.024, arXiv:1410.3012.
- [32] D. J. Lange, “The EvtGen particle decay simulation package”, *Nucl. Instrum. Meth. A* **462** (2001) 152, doi:10.1016/S0168-9002(01)00089-4.
- [33] E. Barberio, B. van Eijk, and Z. Waş, “PHOTOS — a universal Monte Carlo for QED radiative corrections in decays”, *Comput. Phys. Commun.* **66** (1991) 115, doi:10.1016/0010-4655(91)90012-A.
- [34] E. Barberio and Z. Waş, “PHOTOS—a universal Monte Carlo for QED radiative corrections: version 2.0”, *Comput. Phys. Commun.* **79** (1994) 291, doi:10.1016/0010-4655(94)90074-4.
- [35] GEANT4 Collaboration, “GEANT4—a simulation toolkit”, *Nucl. Instrum. Meth. A* **506** (2003) 250, doi:10.1016/S0168-9002(03)01368-8.

- [36] Particle Data Group, “Review of particle physics”, *Phys. Rev. D* **110** (2024) 030001, doi:10.1103/PhysRevD.110.030001.
- [37] M. Pivk and F. R. Le Diberder, “ \mathcal{P} lot: a statistical tool to unfold data distributions”, *Nucl. Instrum. Meth. A* **555** (2005) 356, doi:10.1016/j.nima.2005.08.106, arXiv:physics/0402083.
- [38] M. J. Oreglia, “A study of the reactions $\psi' \rightarrow \gamma\gamma\psi$ ”. PhD thesis, Stanford University, 1980. SLAC Report SLAC-R-236.
- [39] J. E. Gaiser, “Charmonium spectroscopy from radiative decays of the J/ψ and ψ' ”. PhD thesis, Stanford University, 1982. SLAC Report SLAC-R-255.
- [40] CMS Collaboration, “Precise determination of the mass of the Higgs boson and tests of compatibility of its couplings with the standard model predictions using proton collisions at 7 and 8 TeV”, *Eur. Phys. J. C* **75** (2015) 212, doi:10.1140/epjc/s10052-015-3351-7, arXiv:1412.8662.
- [41] R. R. Horgan, Z. Liu, S. Meinel, and M. Wingate, “Lattice QCD calculation of form factors describing the rare decays $B \rightarrow K^*\ell^+\ell^-$ and $B_s \rightarrow \phi\ell^+\ell^-$ ”, *Phys. Rev. D* **89** (2014) 094501, doi:10.1103/PhysRevD.89.094501, arXiv:1310.3722.
- [42] R. R. Horgan, Z. Liu, S. Meinel, and M. Wingate, “Rare B decays using lattice QCD form factors”, in *Proceedings of The 32nd International Symposium on Lattice Field Theory – PoS(LATTICE2014)*, p. 372. 2015. doi:10.22323/1.214.0372.
- [43] N. Gubernari, A. Kokulu, and D. van Dyk, “ $B \rightarrow P$ and $B \rightarrow V$ form factors from B-meson light-cone sum rules beyond leading twist”, *JHEP* **01** (2019) 150, doi:10.1007/JHEP01(2019)150, arXiv:1811.00983.
- [44] N. Gubernari, D. van Dyk, and J. Virto, “Non-local matrix elements in $B_{(s)} \rightarrow \{K^{(*)}, \phi\}\ell^+\ell^-$ ”, *JHEP* **02** (2021) 088, doi:10.1007/JHEP02(2021)088, arXiv:2011.09813.
- [45] N. Gubernari, M. Reboud, D. van Dyk, and J. Virto, “Improved theory predictions and global analysis of exclusive $b \rightarrow s\mu^+\mu^-$ processes”, *JHEP* **09** (2022) 133, doi:10.1007/JHEP09(2022)133, arXiv:2206.03797.
- [46] D. van Dyk et al., “eos/eos: EOS Version 1.0.10”, 2023. doi:10.5281/zenodo.8340859.
- [47] D. M. Straub, “flavio: a Python package for flavour and precision phenomenology in the Standard Model and beyond”, 10, 2018. arXiv:1810.08132.
- [48] A. Bharucha, D. M. Straub, and R. Zwicky, “ $B \rightarrow V\ell^+\ell^-$ in the Standard Model from light-cone sum rules”, *JHEP* **08** (2016) 098, doi:10.1007/JHEP08(2016)098, arXiv:1503.05534.
- [49] M. Beneke, T. Feldmann, and D. Seidel, “Exclusive radiative and electroweak $b \rightarrow d$ and $b \rightarrow s$ penguin decays at NLO”, *Eur. Phys. J. C* **41** (2005) 173, doi:10.1140/epjc/s2005-02181-5, arXiv:hep-ph/0412400.
- [50] M. Beneke, T. Feldmann, and D. Seidel, “Systematic approach to exclusive $B \rightarrow V\ell^+\ell^-$, $V\gamma$ decays”, *Nucl. Phys. B* **612** (2001) 25, doi:10.1016/S0550-3213(01)00366-2, arXiv:hep-ph/0106067.



















- [51] J. De Blas et al., “HEPfit: a code for the combination of indirect and direct constraints on high energy physics models”, *Eur. Phys. J. C* **80** (2020) 456, doi:10.1140/epjc/s10052-020-7904-z, arXiv:1910.14012.
- [52] M. Ciuchini et al., “ $B \rightarrow K^* \ell^+ \ell^-$ decays at large recoil in the Standard Model: a theoretical reappraisal”, *JHEP* **06** (2016) 116, doi:10.1007/JHEP06(2016)116, arXiv:1512.07157.

A The CMS Collaboration

Yerevan Physics Institute, Yerevan, Armenia

A. Hayrapetyan, A. Tumasyan¹ 













Institut für Hochenergiephysik, Vienna, Austria

W. Adam , J.W. Andrejkovic, L. Benato , T. Bergauer , S. Chatterjee , K. Damanakis , M. Dragicevic , P.S. Hussain , M. Jeitler² , N. Krammer , A. Li , D. Liko , I. Mikulec , J. Schieck² , R. Schöfbeck² , D. Schwarz , M. Sonawane , W. Waltenberger , C.-E. Wulz² 

















Universiteit Antwerpen, Antwerpen, Belgium

T. Janssen , T. Van Laer, P. Van Mechelen 













Vrije Universiteit Brussel, Brussel, Belgium

N. Breugelmans, J. D'Hondt , S. Dansana , A. De Moor , M. Delcourt , F. Heyen, S. Lowette , I. Makarenko , D. Müller , S. Tavernier , M. Tytgat³ , G.P. Van Onsem , S. Van Putte , D. Vannerom 














Université Libre de Bruxelles, Bruxelles, Belgium

B. Bilin , B. Clerbaux , A.K. Das, I. De Bruyn , G. De Lentdecker , H. Evard , L. Favart , P. Gianneios , J. Jaramillo , A. Khalilzadeh, F.A. Khan , K. Lee , A. Malara , S. Paredes , M.A. Shahzad, L. Thomas , M. Vanden Bemden , C. Vander Velde , P. Vanlaer 

Ghent University, Ghent, Belgium

M. De Coen , D. Dobur , G. Gokbulut , Y. Hong , J. Knolle , L. Lambrecht , D. Marckx , K. Mota Amarilo , K. Skovpen , N. Van Den Bossche , J. van der Linden , L. Wezenbeek 

















Université Catholique de Louvain, Louvain-la-Neuve, Belgium

A. Benecke , A. Bethani , G. Bruno , C. Caputo , J. De Favereau De Jeneret , C. Delaere , I.S. Donertas , A. Giammanco , A.O. Guzel , Sa. Jain , V. Lemaitre, J. Lidrych , P. Mastrapasqua , T.T. Tran 










Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil

G.A. Alves , E. Coelho , G. Correia Silva , C. Hensel , T. Menezes De Oliveira , C. Mora Herrera⁴ , P. Rebello Teles , M. Soeiro, A. Vilela Pereira⁴ 

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior , M. Barroso Ferreira Filho , H. Brandao Malbouisson , W. Carvalho , J. Chinellato⁵, E.M. Da Costa , G.G. Da Silveira⁶ , D. De Jesus Damiao , S. Fonseca De Souza , R. Gomes De Souza, T. Laux Kuhn⁶, M. Macedo , J. Martins , L. Mundim , H. Nogima , J.P. Pinheiro , A. Santoro , A. Sznajder , M. Thiel 

Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil




C.A. Bernardes⁶ , L. Calligaris , T.R. Fernandez Perez Tomei , E.M. Gregores , I. Maitto Silverio , P.G. Mercadante , S.F. Novaes , B. Orzari , Sandra S. Padula 

Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria


A. Aleksandrov , G. Antchev , R. Hadjiiska , P. Iaydjiev , M. Misheva , M. Shopova , G. Sultanov 












University of Sofia, Sofia, Bulgaria

A. Dimitrov , L. Litov , B. Pavlov , P. Petkov , A. Petrov , E. Shumka 


Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile
S. Keshri , D. Laroze , S. Thakur 


Beihang University, Beijing, China
T. Cheng , T. Javaid , L. Yuan 

Department of Physics, Tsinghua University, Beijing, China
Z. Hu , Z. Liang, J. Liu



Institute of High Energy Physics, Beijing, China
G.M. Chen⁷ , H.S. Chen⁷ , M. Chen⁷ , F. Iemmi , C.H. Jiang, A. Kapoor⁸ , H. Liao ,
Z.-A. Liu⁹ , R. Sharma¹⁰ , J.N. Song⁹, J. Tao , C. Wang⁷, J. Wang , Z. Wang⁷, H. Zhang ,
J. Zhao 

State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China
A. Agapitos , Y. Ban , A. Carvalho Antunes De Oliveira , S. Deng , B. Guo, C. Jiang ,
A. Levin , C. Li , Q. Li , Y. Mao, S. Qian, S.J. Qian , X. Qin, X. Sun , D. Wang , H. Yang,
L. Zhang , Y. Zhao, C. Zhou 

Guangdong Provincial Key Laboratory of Nuclear Science and Guangdong-Hong Kong Joint Laboratory of Quantum Matter, South China Normal University, Guangzhou, China
S. Yang 

Sun Yat-Sen University, Guangzhou, China
Z. You 

University of Science and Technology of China, Hefei, China
K. Jaffel , N. Lu 




Nanjing Normal University, Nanjing, China
G. Bauer¹¹, B. Li¹², K. Yi¹³ , J. Zhang 



Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China
Y. Li

Zhejiang University, Hangzhou, Zhejiang, China
Z. Lin , C. Lu , M. Xiao 

Universidad de Los Andes, Bogota, Colombia
C. Avila , D.A. Barbosa Trujillo, A. Cabrera , C. Florez , J. Fraga , J.A. Reyes Vega

Universidad de Antioquia, Medellin, Colombia
F. Ramirez , C. Rendón, M. Rodriguez , A.A. Ruales Barbosa , J.D. Ruiz Alvarez 

University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia
D. Giljanovic , N. Godinovic , D. Lelas , A. Sculac 




University of Split, Faculty of Science, Split, Croatia
M. Kovac , A. Petkovic, T. Sculac 

Institute Rudjer Boskovic, Zagreb, Croatia
P. Bargassa , V. Brigljevic , B.K. Chitroda , D. Ferencek , K. Jakovcic, A. Starodumov¹⁴ ,
T. Susa 

University of Cyprus, Nicosia, Cyprus

A. Attikis , K. Christoforou , A. Hadjiagapiou, C. Leonidou , J. Mousa , C. Nicolaou, L. Paizanos, F. Ptochos , P.A. Razis , H. Rykaczewski, H. Saka , A. Stepennov 


Charles University, Prague, Czech Republic

M. Finger , M. Finger Jr. , A. Kveton 


Escuela Politecnica Nacional, Quito, Ecuador

E. Ayala 

Universidad San Francisco de Quito, Quito, Ecuador

E. Carrera Jarrin 









Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt

H. Abdalla¹⁵ , Y. Assran^{16,17}, B. El-mahdy

Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt

M.A. Mahmoud , Y. Mohammed 















National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

K. Ehataht , M. Kadastik, T. Lange , S. Nandan , C. Nielsen , J. Pata , M. Raidal , L. Tani , C. Veelken 

Department of Physics, University of Helsinki, Helsinki, Finland

H. Kirschenmann , K. Osterberg , M. Voutilainen 



















Helsinki Institute of Physics, Helsinki, Finland

S. Bharthuar , N. Bin Norjoharuddeen , E. Brücken , F. Garcia , P. Inkaew , K.T.S. Kallonen , T. Lampén , K. Lassila-Perini , S. Lehti , T. Lindén , M. Myllymäki , M.m. Rantanen , H. Siikonen , J. Tuominiemi 





























Lappeenranta-Lahti University of Technology, Lappeenranta, Finland

P. Luukka , H. Petrow 


















IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France

M. Besancon , F. Couderc , M. Dejardin , D. Denegri, J.L. Faure, F. Ferri , S. Ganjour , P. Gras , G. Hamel de Monchenault , M. Kumar , V. Lohezic , J. Malcles , F. Orlandi , L. Portales , A. Rosowsky , M.Ö. Sahin , A. Savoy-Navarro¹⁸ , P. Simkina , M. Titov , M. Tornago 

Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

F. Beaudette , G. Boldrini , P. Busson , A. Cappati , C. Charlot , M. Chiusi , T.D. Cuisset , F. Damas , O. Davignon , A. De Wit , I.T. Ehle , B.A. Fontana Santos Alves , S. Ghosh , A. Gilbert , R. Granier de Cassagnac , A. Hakimi , B. Harikrishnan , L. Kalipoliti , G. Liu , M. Nguyen , C. Ochando , R. Salerno , J.B. Sauvan , Y. Sirois , L. Urda Gómez , E. Vernazza , A. Zabi , A. Zghiche 


















Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France

J.-L. Agram¹⁹ , J. Andrea , D. Apparú , D. Bloch , J.-M. Brom , E.C. Chabert , C. Collard , S. Falke , U. Goerlach , R. Haeberle , A.-C. Le Bihan , M. Meena , O. Poncet , G. Saha , M.A. Sessini , P. Van Hove , P. Vaucelle 

Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France

A. Di Florio 

Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France

D. Amram, S. Beauceron , B. Blancon , G. Boudoul , N. Chanon , D. Contardo , P. Depasse , C. Dozen²⁰ , H. El Mamouni, J. Fay , S. Gascon , M. Gouzevitch , C. Greenberg, G. Grenier , B. Ille , E. Jourd'huy, I.B. Laktineh, M. Lethuillier , L. Mirabito, S. Perries, A. Purohit , M. Vander Donckt , P. Verdier , J. Xiao 






























Georgian Technical University, Tbilisi, Georgia

I. Lomidze , T. Toriashvili²¹ , Z. Tsamalaidze¹⁴ 









RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany

V. Botta , S. Consuegra Rodríguez , L. Feld , K. Klein , M. Lipinski , D. Meuser , A. Pauls , D. Pérez Adán , N. Röwert , M. Teroerde 













































RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

S. Diekmann , A. Dodonova , N. Eich , D. Eliseev , F. Engelke , J. Erdmann , M. Erdmann , P. Fackeldey , B. Fischer , T. Hebbeker , K. Hoepfner , F. Ivone , A. Jung , M.y. Lee , F. Mausolf , M. Merschmeyer , A. Meyer , S. Mukherjee , D. Noll , F. Nowotny, A. Pozdnyakov , Y. Rath, W. Redjeb , F. Rehm, H. Reithler , V. Sarkisovi , A. Schmidt , C. Seth, A. Sharma , J.L. Spah , A. Stein , F. Torres Da Silva De Araujo²² , S. Wiedenbeck , S. Zaleski






































RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany

C. Dziwok , G. Flügge , T. Kress , A. Nowack , O. Pooth , A. Stahl , T. Ziemons , A. Zotz 



































Deutsches Elektronen-Synchrotron, Hamburg, Germany

H. Aarup Petersen , M. Aldaya Martin , J. Alimena , S. Amoroso, Y. An , J. Bach , S. Baxter , M. Bayatmakou , H. Becerril Gonzalez , O. Behnke , A. Belvedere , F. Blekman²³ , K. Borrás²⁴ , A. Campbell , A. Cardini , C. Cheng, F. Colombina , G. Eckerlin, D. Eckstein , L.I. Estevez Banos , E. Gallo²³ , A. Geiser , V. Guglielmi , M. Guthoff , A. Hinzmann , L. Jeppe , B. Kaech , M. Kasemann , C. Kleinwort , R. Kogler , M. Komm , D. Krücker , W. Lange, D. Leyva Pernia , K. Lipka²⁵ , W. Lohmann²⁶ , F. Lorkowski , R. Mankel , I.-A. Melzer-Pellmann , M. Mendizabal Morentin , A.B. Meyer , G. Milella , K. Moral Figueroa , A. Mussgiller , L.P. Nair , J. Niedziela , A. Nürnberg , Y. Otariid, J. Park , E. Ranken , A. Raspereza , D. Rastorguev , J. Rübenach, L. Rygaard, A. Saggio , M. Scham^{27,24} , S. Schnake²⁴ , P. Schütze , C. Schwanenberger²³ , D. Selivanova , K. Sharko , M. Shchedrolosiev , D. Stafford, F. Vazzoler , A. Ventura Barroso , R. Walsh , D. Wang , Q. Wang , K. Wichmann, L. Wiens²⁴ , C. Wissing , Y. Yang , A. Zimmermann Castro Santos 

University of Hamburg, Hamburg, Germany

A. Albrecht , S. Albrecht , M. Antonello , S. Bein , S. Bollweg, M. Bonanomi , P. Connor , K. El Morabit , Y. Fischer , E. Garutti , A. Grohsjean , J. Haller , D. Hundhausen, H.R. Jabusch , G. Kasieczka , P. Keicher, R. Klanner , W. Korcari , T. Kramer , C.c. Kuo, V. Kutzner , F. Labe , J. Lange , A. Lobanov , C. Matthies , L. Moureaux , M. Mrowietz, A. Nigamova , Y. Nissan, A. Paasch , K.J. Pena Rodriguez , T. Quadfasel , B. Raciti , M. Rieger , D. Savoie , J. Schindler , P. Schleper , M. Schröder , J. Schwandt , M. Sommerhalder , H. Stadie , G. Steinbrück , A. Tews, B. Wiederspan, M. Wolf 


Karlsruher Institut fuer Technologie, Karlsruhe, Germany

S. Brommer , E. Butz , T. Chwalek , A. Dierlamm , A. Droll, U. Elicabuk, N. Faltermann , M. Giffels , A. Gottmann , F. Hartmann²⁸ , R. Hofsaess , M. Horzela , U. Husemann , J. Kieseler , M. Klute , R. Koppenhöfer , O. Lavoryk, J.M. Lawhorn , M. Link, A. Lintuluoto , S. Maier , S. Mitra , M. Mormile , Th. Müller , M. Neukum, M. Oh , E. Pfeffer , M. Presilla , G. Quast , K. Rabbertz , B. Regnery , N. Shadskiy , I. Shvetsov , H.J. Simonis , L. Sowa, L. Stockmeier, K. Tauqeer, M. Toms , N. Trevisani , R.F. Von Cube , M. Wassmer , S. Wieland , F. Wittig, R. Wolf , X. Zuo 

Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

G. Anagnostou, G. Daskalakis , A. Kyriakis, A. Papadopoulos²⁸, A. Stakia 







National and Kapodistrian University of Athens, Athens, Greece

P. Kontaxakis , G. Melachroinos, Z. Painesis , I. Papavergou , I. Paraskevas , N. Saoulidou , K. Theofilatos , E. Tziaferi , K. Vellidis , I. Zisopoulos 






National Technical University of Athens, Athens, Greece

G. Bakas , T. Chatzistavrou, G. Karapostoli , K. Kousouris , I. Papakrivopoulos , E. Siamarkou, G. Tsipolitis , A. Zacharopoulou

University of Ioánnina, Ioánnina, Greece

I. Bestintzanos, I. Evangelou , C. Foudas, C. Kamtsikis, P. Katsoulis, P. Kokkas , P.G. Kosmoglou Kioseglou , N. Manthos , I. Papadopoulos , J. Strologas 

HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

C. Hajdu , D. Horvath^{29,30} , K. Márton, A.J. Rádl³¹ , F. Sikler , V. Veszpremi 

MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

M. Csanád , K. Farkas , A. Fehérkuti³² , M.M.A. Gadallah³³ , Á. Kadlecik , P. Major , G. Pásztor , G.I. Veres 




Faculty of Informatics, University of Debrecen, Debrecen, Hungary

L. Olah , G. Zilizi 













HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary

G. Bencze, S. Czellar, J. Molnar, Z. Szillasi

Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary

T. Csorgo³² , F. Nemes³² , T. Novak 

Panjab University, Chandigarh, India

S. Bansal , S.B. Beri, V. Bhatnagar , G. Chaudhary , S. Chauhan , N. Dhingra³⁴ , A. Kaur , A. Kaur , H. Kaur , M. Kaur , S. Kumar , T. Sheokand, J.B. Singh , A. Singla 

University of Delhi, Delhi, India






A. Ahmed , A. Bhardwaj , A. Chhetri , B.C. Choudhary , A. Kumar , A. Kumar , M. Naimuddin , K. Ranjan , M.K. Saini, S. Saumya 

Saha Institute of Nuclear Physics, HBNI, Kolkata, India

S. Baradia , S. Barman³⁵ , S. Bhattacharya , S. Das Gupta, S. Dutta , S. Dutta, S. Sarkar

Indian Institute of Technology Madras, Madras, India


M.M. Ameen , P.K. Behera , S.C. Behera , S. Chatterjee , G. Dash , P. Jana , P. Kalbhor , S. Kamble , J.R. Komaragiri³⁶ , D. Kumar³⁶ , T. Mishra , B. Parida³⁷ 

P.R. Pujahari , N.R. Saha , A. Sharma , A.K. Sikdar , R.K. Singh, P. Verma, S. Verma ,
A. Vijay











Tata Institute of Fundamental Research-A, Mumbai, India

S. Dugad, G.B. Mohanty , M. Shelake, P. Suryadevara

Tata Institute of Fundamental Research-B, Mumbai, India

A. Bala , S. Banerjee , R.M. Chatterjee, M. Guchait , Sh. Jain , A. Jaiswal, S. Kumar ,
G. Majumder , K. Mazumdar , S. Parolia , A. Thachayath 

National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India

S. Bahinipati³⁸ , C. Kar , D. Maity³⁹ , P. Mal , V.K. Muraleedharan Nair Bindhu³⁹ ,
K. Naskar³⁹ , A. Nayak³⁹ , S. Nayak, K. Pal, P. Sadangi, S.K. Swain , S. Varghese³⁹ ,
D. Vats³⁹ 







Indian Institute of Science Education and Research (IISER), Pune, India

S. Acharya⁴⁰ , A. Alpana , S. Dube , B. Gomber⁴⁰ , P. Hazarika , B. Kansal ,
A. Laha , B. Sahu⁴⁰ , S. Sharma , K.Y. Vaish 

Isfahan University of Technology, Isfahan, Iran

H. Bakhshiansohi⁴¹ , A. Jafari⁴² , M. Zeinali⁴³ 


























Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

S. Bashiri, S. Chenarani⁴⁴ , S.M. Etesami , Y. Hosseini , M. Khakzad , E. Khazaie ,
M. Mohammadi Najafabadi , S. Tizchang⁴⁵ 










University College Dublin, Dublin, Ireland

M. Felcini , M. Grunewald 







INFN Sezione di Bari^a, Università di Bari^b, Politecnico di Bari^c, Bari, Italy

M. Abbrescia^{a,b} , A. Colaleo^{a,b} , D. Creanza^{a,c} , B. D'Anzi^{a,b} , N. De Filippis^{a,c} ,
M. De Palma^{a,b} , W. Elmetenawee^{a,b,46} , N. Ferrara^{a,b} , L. Fiore^a , G. Iaselli^{a,c} ,
L. Longo^a , M. Louka^{a,b} , G. Maggi^{a,c} , M. Maggi^a , I. Margjeka^a , V. Mastrapasqua^{a,b} ,
S. My^{a,b} , S. Nuzzo^{a,b} , A. Pellecchia^{a,b} , A. Pompili^{a,b} , G. Pugliese^{a,c} ,
R. Radogna^{a,b} , D. Ramos^a , A. Ranieri^a , L. Silvestris^a , F.M. Simone^{a,c} ,
Ü. Sözbilir^a , A. Stamerra^{a,b} , D. Troiano^{a,b} , R. Venditti^{a,b} , P. Verwilligen^a ,
A. Zaza^{a,b}










INFN Sezione di Bologna^a, Università di Bologna^b, Bologna, Italy

G. Abbiendi^a , C. Battilana^{a,b} , D. Bonacorsi^{a,b} , P. Capiluppi^{a,b} , A. Castro^{+a,b} ,
F.R. Cavallo^a , M. Cuffiani^{a,b} , G.M. Dallavalle^a , T. Diotallevi^{a,b} , F. Fabbri^a ,
A. Fanfani^{a,b} , D. Fasanella^a , P. Giacomelli^a , L. Giommi^{a,b} , C. Grandi^a ,
L. Guiducci^{a,b} , S. Lo Meo^{a,47} , M. Lorusso^{a,b} , L. Lunerti^a , S. Marcellini^a ,
G. Masetti^a , F.L. Navarra^{a,b} , G. Paggi^{a,b} , A. Perrotta^a , F. Primavera^{a,b} ,
A.M. Rossi^{a,b} , S. Rossi Tisbeni^{a,b} , T. Rovelli^{a,b} , G.P. Siroli^{a,b}

INFN Sezione di Catania^a, Università di Catania^b, Catania, Italy

S. Costa^{a,b,48} , A. Di Mattia^a , A. Lapertosa^a , R. Potenza^{a,b} , A. Tricomi^{a,b,48} ,
C. Tuve^{a,b} 

INFN Sezione di Firenze^a, Università di Firenze^b, Firenze, Italy

P. Assiouras^a , G. Barbagli^a , G. Bardelli^{a,b} , B. Camaiani^{a,b} , A. Cassese^a ,
R. Ceccarelli^a , V. Ciulli^{a,b} , C. Civinini^a , R. D'Alessandro^{a,b} , E. Focardi^{a,b} ,
T. Kello^a , G. Latino^{a,b} , P. Lenzi^{a,b} , M. Lizzo^a , M. Meschini^a , S. Paoletti^a

A. Papanastassiou^{a,b}, G. Sguazzoni^a , L. Vilianni^a 












INFN Laboratori Nazionali di Frascati, Frascati, Italy

L. Benussi , S. Bianco , S. Meola⁴⁹ , D. Piccolo 



INFN Sezione di Genova^a, Università di Genova^b, Genova, Italy

M. Alves Gallo Pereira^a , F. Ferro^a , E. Robutti^a , S. Tosi^{a,b} 










INFN Sezione di Milano-Bicocca^a, Università di Milano-Bicocca^b, Milano, Italy

A. Benaglia^a , F. Brivio^a , F. Cetorelli^{a,b} , F. De Guio^{a,b} , M.E. Dinardo^{a,b} , P. Dini^a , S. Gennai^a , R. Gerosa^{a,b} , A. Ghezzi^{a,b} , P. Govoni^{a,b} , L. Guzzi^a , M.T. Lucchini^{a,b} , M. Malberti^a , S. Malvezzi^a , A. Massironi^a , D. Menasce^a , L. Moroni^a , M. Paganoni^{a,b} , S. Palluotto^{a,b} , D. Pedrini^a , A. Perego^{a,b} , B.S. Pinolini^a, G. Pizzati^{a,b}, S. Ragazzi^{a,b} , T. Tabarelli de Fatis^{a,b} 










INFN Sezione di Napoli^a, Università di Napoli 'Federico II'^b, Napoli, Italy; Università della Basilicata^c, Potenza, Italy; Scuola Superiore Meridionale (SSM)^d, Napoli, Italy

S. Buontempo^a , A. Cagnotta^{a,b} , F. Carnevali^{a,b}, N. Cavallo^{a,c} , F. Fabozzi^{a,c} , A.O.M. Iorio^{a,b} , L. Lista^{a,b,50} , P. Paolucci^{a,28} , B. Rossi^a 










INFN Sezione di Padova^a, Università di Padova^b, Padova, Italy; Università di Trento^c, Trento, Italy

R. Ardino^a , P. Azzi^a , N. Bacchetta^{a,51} , D. Bisello^{a,b} , P. Bortignon^a , G. Bortolato^{a,b}, A. Bragagnolo^{a,b} , A.C.M. Bulla^a , R. Carlin^{a,b} , T. Dorigo^{a,52} , F. Gasparini^{a,b} , U. Gasparini^{a,b} , S. Giorgetti^a, M. Gulmini^{a,53} , E. Lusiani^a , M. Margoni^{a,b} , G. Maron^{a,53} , A.T. Meneguzzo^{a,b} , M. Migliorini^{a,b} , J. Pazzini^{a,b} , P. Ronchese^{a,b} , R. Rossin^{a,b} , F. Simonetto^{a,b} , M. Tosi^{a,b} , A. Triossi^{a,b} , M. Zanetti^{a,b} , P. Zotto^{a,b} , A. Zucchetta^{a,b} , G. Zumerle^{a,b}










INFN Sezione di Pavia^a, Università di Pavia^b, Pavia, Italy

A. Braghieri^a , S. Calzaferri^a , D. Fiorina^a , P. Montagna^{a,b} , V. Re^a , C. Riccardi^{a,b} , P. Salvini^a , I. Vai^{a,b} , P. Vitulo^{a,b} 










INFN Sezione di Perugia^a, Università di Perugia^b, Perugia, Italy

S. Ajmal^{a,b} , M.E. Ascioti^{a,b}, G.M. Bilei^a , C. Carrivale^{a,b}, D. Ciangottini^{a,b} , L. Fanò^{a,b} , M. Magherini^{a,b} , V. Mariani^{a,b} , M. Menichelli^a , F. Moscatelli^{a,54} , A. Rossi^{a,b} , A. Santocchia^{a,b} , D. Spiga^a , T. Tedeschi^{a,b}









































INFN Sezione di Pisa^a, Università di Pisa^b, Scuola Normale Superiore di Pisa^c, Pisa, Italy; Università di Siena^d, Siena, Italy

C. Aimè^a , C.A. Alexe^{a,c} , P. Asenov^{a,b} , P. Azzurri^a , G. Bagliesi^a , R. Bhattacharya^a , L. Bianchini^{a,b} , T. Boccali^a , E. Bossini^a , D. Bruschini^{a,c} , R. Castaldi^a , M.A. Ciocci^{a,b} , M. Cipriani^{a,b} , V. D'Amante^{a,d} , R. Dell'Orso^a , S. Donato^a , A. Giassi^a , F. Ligabue^{a,c} , A.C. Marini^a , D. Matos Figueiredo^a , A. Messineo^{a,b} , S. Mishra^a , M. Musich^{a,b} , F. Palla^a , A. Rizzi^{a,b} , G. Rolandi^{a,c} , S. Roy Chowdhury^a , T. Sarkar^a , A. Scribano^a , P. Spagnolo^a , R. Tenchini^a , G. Tonelli^{a,b} , N. Turini^{a,d} , F. Vaselli^{a,c} , A. Venturi^a , P.G. Verdini^a








INFN Sezione di Roma^a, Sapienza Università di Roma^b, Roma, Italy

C. Baldenegro Barrera^{a,b} , P. Barria^a , C. Basile^{a,b} , F. Cavallari^a , L. Cunqueiro Mendez^{a,b} , D. Del Re^{a,b} , E. Di Marco^{a,b} , M. Diemoz^a , F. Errico^{a,b} , R. Gargiulo^{a,b}, E. Longo^{a,b} , L. Martikainen^{a,b} , J. Mijuskovic^{a,b} , G. Organtini^{a,b} , F. Pandolfi^a , R. Paramatti^{a,b} , C. Quaranta^{a,b} , S. Rahatlou^{a,b} , C. Rovelli^a , F. Santanastasio^{a,b} , L. Soffi^a , V. Vladimirov^{a,b}

INFN Sezione di Torino^a, Università di Torino^b, Torino, Italy; Università del Piemonte Orientale^c, Novara, Italy

N. Amapane^{a,b} , R. Arcidiacono^{a,c} , S. Argiro^{a,b} , M. Arneodo^{a,c} , N. Bartosik^a , R. Bellan^{a,b} , A. Bellora^{a,b} , C. Biino^a , C. Borca^{a,b} , N. Cartiglia^a , M. Costa^{a,b} , R. Covarelli^{a,b} , N. Demaria^a , L. Finco^a , M. Grippo^{a,b} , B. Kiani^{a,b} , F. Legger^a , F. Luongo^{a,b} , C. Mariotti^a , L. Markovic^{a,b} , S. Maselli^a , A. Mecca^{a,b} , L. Menzio^{a,b}, P. Meridiani^a , E. Migliore^{a,b} , M. Monteno^a , R. Mulargia^a , M.M. Obertino^{a,b} , G. Ortona^a , L. Pacher^{a,b} , N. Pastrone^a , M. Pelliccioni^a , M. Ruspa^{a,c} , F. Siviero^{a,b} , V. Sola^{a,b} , A. Solano^{a,b} , A. Staiano^a , C. Tarricone^{a,b} , D. Trocino^a , G. Umoret^{a,b} , R. White^{a,b} 


INFN Sezione di Trieste^a, Università di Trieste^b, Trieste, Italy

J. Babbar^{a,b} , S. Belforte^a , V. Candelise^{a,b} , M. Casarsa^a , F. Cossutti^a , K. De Leo^a , G. Della Ricca^{a,b} 

Kyungpook National University, Daegu, Korea

S. Dogra , J. Hong , B. Kim , J. Kim, D. Lee, H. Lee, S.W. Lee , C.S. Moon , Y.D. Oh , M.S. Ryu , S. Sekmen , B. Tae, Y.C. Yang 

Department of Mathematics and Physics - GWNu, Gangneung, Korea

M.S. Kim 

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

G. Bak , P. Gwak , H. Kim , D.H. Moon 

Hanyang University, Seoul, Korea

E. Asilar , J. Choi⁵⁵ , D. Kim , T.J. Kim , J.A. Merlin, Y. Ryou

Korea University, Seoul, Korea

S. Choi , S. Han, B. Hong , K. Lee, K.S. Lee , S. Lee , J. Yoo 













Kyung Hee University, Department of Physics, Seoul, Korea

J. Goh , S. Yang 








Sejong University, Seoul, Korea

H. S. Kim , Y. Kim, S. Lee



Seoul National University, Seoul, Korea

J. Almond, J.H. Bhyun, J. Choi , J. Choi, W. Jun , J. Kim , Y.W. Kim, S. Ko , H. Kwon , H. Lee , J. Lee , J. Lee , B.H. Oh , S.B. Oh , H. Seo , U.K. Yang, I. Yoon 

University of Seoul, Seoul, Korea

W. Jang , D.Y. Kang, Y. Kang , S. Kim , B. Ko, J.S.H. Lee , Y. Lee , I.C. Park , Y. Roh, I.J. Watson 

Yonsei University, Department of Physics, Seoul, Korea














































S. Ha , K. Hwang, H.D. Yoo 





















































Sungkyunkwan University, Suwon, Korea

M. Choi , M.R. Kim , H. Lee, Y. Lee , I. Yu 













College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait

T. Beyrouthy, Y. Gharbia











Kuwait University - College of Science - Department of Physics, Safat, KuwaitF. Alazemi **Riga Technical University, Riga, Latvia**K. Dreimanis , A. Gaile , C. Munoz Diaz, D. Osite , G. Pikurs, A. Potrebko , M. Seidel ,
D. Sidiropoulos Kontos**University of Latvia (LU), Riga, Latvia**N.R. Strautnieks **Vilnius University, Vilnius, Lithuania**M. Ambrozas , A. Juodagalvis , A. Rinkevicius , G. Tamulaitis **National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia**I. Yusuff⁵⁶ , Z. Zolkapli**Universidad de Sonora (UNISON), Hermosillo, Mexico**J.F. Benitez , A. Castaneda Hernandez , H.A. Encinas Acosta, L.G. Gallegos Maríñez,
M. León Coello , J.A. Murillo Quijada , A. Sehrawat , L. Valencia Palomo **Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico**G. Ayala , H. Castilla-Valdez , H. Crotte Ledesma, E. De La Cruz-Burelo , I. Heredia-
De La Cruz⁵⁷ , R. Lopez-Fernandez , J. Mejia Guisao , C.A. Mondragon Herrera,
A. Sánchez Hernández **Universidad Iberoamericana, Mexico City, Mexico**C. Oropeza Barrera , D.L. Ramirez Guadarrama, M. Ramírez García **Benemerita Universidad Autonoma de Puebla, Puebla, Mexico**I. Bautista , I. Pedraza , H.A. Salazar Ibarguen , C. Uribe Estrada **University of Montenegro, Podgorica, Montenegro**I. Bubanja , N. Raicevic **University of Canterbury, Christchurch, New Zealand**P.H. Butler **National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan**A. Ahmad , M.I. Asghar, A. Awais , M.I.M. Awan, H.R. Hoorani , W.A. Khan **AGH University of Krakow, Krakow, Poland**V. Avati, L. Grzanka , M. Malawski **National Centre for Nuclear Research, Swierk, Poland**H. Bialkowska , M. Bluj , M. Górski , M. Kazana , M. Szeleper , P. Zalewski **Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland**K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolikowski ,
A. Muhammad **Warsaw University of Technology, Warsaw, Poland**P. Fokow , K. Pozniak , W. Zabolotny **Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal**M. Araujo , D. Bastos , C. Beirão Da Cruz E Silva , A. Boletti , M. Bozzo ,
T. Camporesi , G. Da Molin , P. Faccioli , M. Gallinaro , J. Hollar , N. Leonardo ,
G.B. Marozzo, A. Petrilli , M. Pisano , J. Seixas , J. Varela , J.W. Wulff

Faculty of Physics, University of Belgrade, Belgrade, SerbiaP. Adzic , P. Milenovic **VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia**D. Devetak, M. Dordevic , J. Milosevic , L. Nadderd , V. Rekovic**Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain**J. Alcaraz Maestre , Cristina F. Bedoya , J.A. Brochero Cifuentes , Oliver M. Carretero , M. Cepeda , M. Cerrada , N. Colino , B. De La Cruz , A. Delgado Peris , A. Escalante Del Valle , D. Fernández Del Val , J.P. Fernández Ramos , J. Flix , M.C. Fouz , O. Gonzalez Lopez , S. Goy Lopez , J.M. Hernandez , M.I. Josa , J. Llorente Merino , C. Martin Perez , E. Martin Viscasillas , D. Moran , C. M. Morcillo Perez , Á. Navarro Tobar , C. Perez Dengra , A. Pérez-Calero Yzquierdo , J. Puerta Pelayo , I. Redondo , S. Sánchez Navas , J. Sastre , J. Vazquez Escobar **Universidad Autónoma de Madrid, Madrid, Spain**J.F. de Trocóniz **Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain**B. Alvarez Gonzalez , J. Cuevas , J. Fernandez Menendez , S. Folgueras , I. Gonzalez Caballero , P. Leguina , E. Palencia Cortezon , J. Prado Pico, C. Ramón Álvarez , V. Rodríguez Bouza , A. Soto Rodríguez , A. Trapote , C. Vico Villalba , P. Vischia **Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain**S. Bhowmik , S. Blanco Fernández , I.J. Cabrillo , A. Calderon , J. Duarte Campderros , M. Fernandez , G. Gomez , C. Lasiosa García , R. Lopez Ruiz , C. Martinez Rivero , P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas , E. Navarrete Ramos , J. Piedra Gomez , L. Scodellaro , I. Vila , J.M. Vizán Garcia **University of Colombo, Colombo, Sri Lanka**B. Kailasapathy⁵⁸ , D.D.C. Wickramarathna **University of Ruhuna, Department of Physics, Matara, Sri Lanka**W.G.D. Dharmaratna⁵⁹ , K. Liyanage , N. Perera **CERN, European Organization for Nuclear Research, Geneva, Switzerland**D. Abbaneo , C. Amendola , E. Auffray , G. Auzinger , J. Baechler, D. Barney , A. Bermúdez Martínez , M. Bianco , A.A. Bin Anuar , A. Bocci , L. Boronovi , C. Botta , E. Brondolin , C. Caillol , G. Cerminara , N. Chernyavskaya , D. d'Enterria , A. Dabrowski , A. David , A. De Roeck , M.M. Defranchis , M. Deile , M. Dobson , G. Franzoni , W. Funk , S. Giani, D. Gigi, K. Gill , F. Glege , J. Hegeman , J.K. Heikkilä , B. Huber, V. Innocente , T. James , P. Janot , O. Kaluzinska , O. Karacheban²⁶ , S. Laurila , P. Lecoq , E. Leutgeb , C. Lourenço , L. Malgeri , M. Mannelli , M. Matthewman, A. Mehta , F. Meijers , S. Mersi , E. Meschi , V. Milosevic , F. Monti , F. Moortgat , M. Mulders , I. Neutelings , S. Orfanelli, F. Pantaleo , G. Petrucciani , A. Pfeiffer , M. Pierini , H. Qu , D. Rabady , B. Ribeiro Lopes , F. Riti , M. Rovere , H. Sakulin , R. Salvatico , S. Sanchez Cruz , S. Scarfi , C. Schwick, M. Selvaggi , A. Sharma , K. Shchelina , P. Silva , P. Sphicas⁶⁰ , A.G. Stahl Leiton , A. Steen , S. Summers , D. Treille , P. Tropea , D. Walter , J. Wanczyk⁶¹ , J. Wang, K.A. Wozniak⁶² , S. Wuchterl , P. Zehetner , P. Zejdl , W.D. Zeuner



















PSI Center for Neutron and Muon Sciences, Villigen, Switzerland

T. Bevilacqua⁶³ , L. Caminada⁶³ , A. Ebrahimi , W. Erdmann , R. Horisberger ,
Q. Ingram , H.C. Kaestli , D. Kotlinski , C. Lange , M. Missiroli⁶³ , L. Noehte⁶³ ,
T. Rohe , A. Samalan




ETH Zurich - Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

T.K. Aarrestad , M. Backhaus , G. Bonomelli, A. Calandri , C. Cazzaniga , K. Datta ,
P. De Bryas Dexmiers D'archiac⁶¹ , A. De Cosa , G. Dissertori , M. Dittmar,
M. Donegà , F. Eble , M. Galli , K. Gedia , F. Glessgen , C. Grab , N. Härringer ,
T.G. Harte, D. Hits , W. Lustermann , A.-M. Lyon , R.A. Manzoni , M. Marchegiani ,
L. Marchese , A. Mascellani⁶¹ , F. Nessi-Tedaldi , F. Pauss , V. Perovic , S. Pigazzini ,
B. Ristic , R. Seidita , J. Steggemann⁶¹ , A. Tarabini , D. Valsecchi , R. Wallny 












Universität Zürich, Zurich, Switzerland

C. Amsler⁶⁴ , P. Bärttschi , M.F. Canelli , K. Cormier , M. Huwiler , W. Jin ,
A. Jofrehei , B. Kilminster , S. Leontsinis , S.P. Liehti , A. Macchiolo , P. Meiring ,
F. Meng , J. Motta , A. Reimers , P. Robmann, M. Senger , E. Shokr, F. Stäger ,
R. Tramontano 

National Central University, Chung-Li, Taiwan

C. Adloff⁶⁵ , D. Bhowmik, C.M. Kuo, W. Lin, P.K. Rout , P.C. Tiwari³⁶ 




















National Taiwan University (NTU), Taipei, Taiwan

L. Ceard, K.F. Chen , Z.g. Chen, A. De Iorio , W.-S. Hou , T.h. Hsu, Y.w. Kao,
S. Karmakar , G. Kole , Y.y. Li , R.-S. Lu , E. Paganis , X.f. Su , J. Thomas-Wilsker ,
L.s. Tsai, D. Tsionou, H.y. Wu, E. Yazgan 

High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

C. Asawatangtrakuldee , N. Srimanobhas , V. Wachirapusanand 

Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

D. Agyel , F. Boran , F. Dolek , I. Dumanoglu⁶⁶ , E. Eskut , Y. Guler⁶⁷ ,
E. Gurpinar Guler⁶⁷ , C. Isik , O. Kara, A. Kayis Topaksu , U. Kiminsu , Y. Komurcu ,
G. Onengut , K. Ozdemir⁶⁸ , A. Polatoz , B. Tali⁶⁹ , U.G. Tok , S. Turkcapar ,
E. Uslan , I.S. Zorbakir 

Middle East Technical University, Physics Department, Ankara, Turkey

G. Sokmen, M. Yalvac⁷⁰ 

Bogazici University, Istanbul, Turkey

B. Akgun , I.O. Atakisi , E. Gülmez , M. Kaya⁷¹ , O. Kaya⁷² , S. Tekten⁷³ 





Istanbul Technical University, Istanbul, Turkey

A. Cakir , K. Cankocak^{66,74} , G.G. Dincer⁶⁶ , S. Sen⁷⁵ 

Istanbul University, Istanbul, Turkey

O. Aydilek⁷⁶ , B. Hacisahinoglu , I. Hos⁷⁷ , B. Kaynak , S. Ozkorucuklu , O. Potok ,
H. Sert , C. Simsek , C. Zorbilmez 

Yildiz Technical University, Istanbul, Turkey














S. Cerci , B. Isildak⁷⁸ , D. Sunar Cerci , T. Yetkin 

Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine




















A. Boyaryntsev , B. Grynyov 

National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine
L. Levchuk 





































University of Bristol, Bristol, United Kingdom

D. Anthony , J.J. Brooke , A. Bundock , F. Bury , E. Clement , D. Cussans ,
H. Flacher , M. Glowacki, J. Goldstein , H.F. Heath , M.-L. Holmberg , L. Kreczko ,
S. Paramesvaran , L. Robertshaw, V.J. Smith , K. Walkingshaw Pass



Rutherford Appleton Laboratory, Didcot, United Kingdom

A.H. Ball, K.W. Bell , A. Belyaev⁷⁹ , C. Brew , R.M. Brown , D.J.A. Cockerill ,
C. Cooke , A. Elliot , K.V. Ellis, K. Harder , S. Harper , J. Linacre , K. Manolopoulos,
D.M. Newbold , E. Olaiya, D. Petyt , T. Reis , A.R. Sahasransu , G. Salvi , T. Schuh,
C.H. Shepherd-Themistocleous , I.R. Tomalin , K.C. Whalen , T. Williams 













Imperial College, London, United Kingdom

I. Andreou , R. Bainbridge , P. Bloch , C.E. Brown , O. Buchmuller, C.A. Carrillo Montoya , G.S. Chahal⁸⁰ , D. Colling , J.S. Dancu, I. Das , P. Dauncey , G. Davies ,
M. Della Negra , S. Fayer, G. Fedi , G. Hall , A. Howard, G. Iles , C.R. Knight ,
P. Krueper, J. Langford , K.H. Law , J. León Holgado , L. Lyons , A.-M. Magnan ,
B. Maier , S. Mallios, M. Mieskolainen , J. Nash⁸¹ , M. Pesaresi , P.B. Pradeep,
B.C. Radburn-Smith , A. Richards, A. Rose , K. Savva , C. Seez , R. Shukla ,
A. Tapper , K. Uchida , G.P. Uttley , T. Virdee²⁸ , M. Vojinovic , N. Wardle ,
D. Winterbottom 

Brunel University, Uxbridge, United Kingdom

J.E. Cole , A. Khan, P. Kyberd , I.D. Reid 

Baylor University, Waco, Texas, USA

S. Abdullin , A. Brinkerhoff , E. Collins , M.R. Darwish , J. Dittmann ,
K. Hatakeyama , J. Hiltbrand , B. McMaster , J. Samudio , S. Sawant ,
C. Sutantawibul , J. Wilson 



















Catholic University of America, Washington, DC, USA

R. Bartek , A. Dominguez , A.E. Simsek , S.S. Yu 


















The University of Alabama, Tuscaloosa, Alabama, USA

B. Bam , A. Buchot Perraguin , R. Chudasama , S.I. Cooper , C. Crovella ,
S.V. Gleyzer , E. Pearson, C.U. Perez , P. Rumerio⁸² , E. Usai , R. Yi 






Boston University, Boston, Massachusetts, USA

A. Akpinar , C. Cosby , G. De Castro, Z. Demiragli , C. Erice , C. Fangmeier ,
C. Fernandez Madrazo , E. Fontanesi , D. Gastler , F. Golf , S. Jeon , J. O'cain,
I. Reed , J. Rohlf , K. Salyer , D. Sperka , D. Spitzbart , I. Suarez , A. Tsatsos ,
A.G. Zecchinelli 

Brown University, Providence, Rhode Island, USA












G. Barone , G. Benelli , D. Cutts , L. Gouskos , M. Hadley , U. Heintz ,
J.M. Hogan⁸³ , T. Kwon , G. Landsberg , K.T. Lau , D. Li , J. Luo , S. Mondal ,
T. Russell, S. Sagir⁸⁴ , X. Shen, F. Simpson , M. Stamenkovic , N. Venkatasubramanian,
X. Yan 

University of California, Davis, Davis, California, USA




S. Abbott , B. Barton , C. Brainerd , R. Breedon , H. Cai 

M. Calderon De La Barca Sanchez , M. Chertok , M. Citron , J. Conway , P.T. Cox , R. Erbacher , F. Jensen , O. Kukral , G. Mocellin , M. Mulhearn , S. Ostrom , W. Wei , S. Yoo , F. Zhang 

















University of California, Los Angeles, California, USA

K. Adamidis, M. Bachtis , D. Campos, R. Cousins , A. Datta , G. Flores Avila , J. Hauser , M. Ignatenko , M.A. Iqbal , T. Lam , Y.f. Lo, E. Manca , A. Nunez Del Prado, D. Saltzberg , V. Valuev 

University of California, Riverside, Riverside, California, USA

R. Clare , J.W. Gary , G. Hanson 










University of California, San Diego, La Jolla, California, USA

A. Aportela, A. Arora , J.G. Branson , S. Cittolin , S. Cooperstein , D. Diaz , J. Duarte , L. Giannini , Y. Gu, J. Guiang , R. Kansal , V. Krutelyov , R. Lee , J. Letts , M. Masciovecchio , F. Mokhtar , S. Mukherjee , M. Pieri , M. Quinnan , B.V. Sathia Narayanan , V. Sharma , M. Tadel , E. Vourliotis , F. Würthwein , Y. Xiang , A. Yagil















University of California, Santa Barbara - Department of Physics, Santa Barbara, California, USA

A. Barzdukas , L. Brennan , C. Campagnari , K. Downham , C. Grieco , J. Incandela , J. Kim , A.J. Li , P. Masterson , H. Mei , J. Richman , S.N. Santpur , U. Sarica , R. Schmitz , F. Setti , J. Sheplock , D. Stuart , T.Á. Vámi , S. Wang , D. Zhang















California Institute of Technology, Pasadena, California, USA

S. Bhattacharya , A. Bornheim , O. Cerri, A. Latorre, J. Mao , H.B. Newman , G. Reales Gutiérrez, M. Spiropulu , J.R. Vlimant , C. Wang , S. Xie , R.Y. Zhu 
















Carnegie Mellon University, Pittsburgh, Pennsylvania, USA

J. Alison , S. An , P. Bryant , M. Cremonesi, V. Dutta , T. Ferguson , T.A. Gómez Espinosa , A. Harilal , A. Kallil Tharayil, C. Liu , T. Mudholkar , S. Murthy , P. Palit , K. Park, M. Paulini , A. Roberts , A. Sanchez , W. Terrill


















University of Colorado Boulder, Boulder, Colorado, USA










J.P. Cumalat , W.T. Ford , A. Hart , A. Hassani , G. Karathanasis , N. Manganelli , J. Parkes , C. Savard , N. Schonbeck , K. Stenson , K.A. Ulmer , S.R. Wagner , N. Zipper , D. Zuolo 

Cornell University, Ithaca, New York, USA










J. Alexander , S. Bright-Thonney , X. Chen , D.J. Cranshaw , J. Fan , X. Fan , S. Hogan , P. Kotamnives, J. Monroy , M. Oshiro , J.R. Patterson , M. Reid , A. Ryd , J. Thom , P. Wittich , R. Zou 

Fermi National Accelerator Laboratory, Batavia, Illinois, USA










M. Albrow , M. Alyari , O. Amram , G. Apollinari , A. Apresyan , L.A.T. Bauerdick , D. Berry , J. Berryhill , P.C. Bhat , K. Burkett , J.N. Butler , A. Canepa , G.B. Cerati , H.W.K. Cheung , F. Chlebana , G. Cummings , J. Dickinson , I. Dutta , V.D. Elvira , Y. Feng , J. Freeman , A. Gandrakota , Z. Gecse , L. Gray , D. Green, A. Grummer , S. Grünendahl , D. Guerrero , O. Gutsche , R.M. Harris , R. Heller , T.C. Herwig , J. Hirschauer , B. Jayatilaka , S. Jindariani , M. Johnson , U. Joshi , T. Klijnsma , B. Klima , K.H.M. Kwok , S. Lammel , C. Lee , D. Lincoln , R. Lipton , T. Liu , C. Madrid , K. Maeshima , C. Mantilla , D. Mason , P. McBride , P. Merkel , S. Mrenna , S. Nahn , J. Ngadiuba , D. Noonan , S. Norberg, V. Papadimitriou

N. Pastika , K. Pedro , C. Pena⁸⁵ , F. Ravera , A. Reinsvold Hall⁸⁶ , L. Ristori , M. Safdari , E. Sexton-Kennedy , N. Smith , A. Soha , L. Spiegel , S. Stoynev , J. Strait , L. Taylor , S. Tkaczyk , N.V. Tran , L. Uplegger , E.W. Vaandering , I. Zoi






University of Florida, Gainesville, Florida, USA

C. Aruta , P. Avery , D. Bourilkov , P. Chang , V. Cherepanov , R.D. Field, C. Huh , E. Koenig , M. Kolosova , J. Konigsberg , A. Korytov , K. Matchev , N. Menendez , G. Mitselmakher , K. Mohrman , A. Muthirakalayil Madhu , N. Rawal , S. Rosenzweig , Y. Takahashi , J. Wang










Florida State University, Tallahassee, Florida, USA

T. Adams , A. Al Kadhim , A. Askew , S. Bower , R. Hashmi , R.S. Kim , S. Kim , T. Kolberg , G. Martinez, H. Prosper , P.R. Prova, M. Wulansatiti , R. Yohay , J. Zhang










Florida Institute of Technology, Melbourne, Florida, USA

B. Alsufyani, M.M. Baarmand , S. Butalla , S. Das , T. Elkafrawy⁸⁷ , M. Hohmann , E. Yanes










University of Illinois Chicago, Chicago, Illinois, USA

M.R. Adams , A. Baty , C. Bennett, R. Cavanaugh , R. Escobar Franco , O. Evdokimov , C.E. Gerber , M. Hawksworth, A. Hingrajiya, D.J. Hofman , J.h. Lee , D. S. Lemos , A.H. Merrit , C. Mills , S. Nanda , G. Oh , B. Ozek , D. Pilipovic , R. Pradhan , E. Prifti, T. Roy , S. Rudrabhatla , N. Singh, M.B. Tonjes , N. Varelas , M.A. Wadud , Z. Ye , J. Yoo










The University of Iowa, Iowa City, Iowa, USA

M. Alhousseini , D. Blend, K. Dilsiz⁸⁸ , L. Emediato , G. Karaman , O.K. Köseyan , J.-P. Merlo, A. Mestvirishvili⁸⁹ , O. Neogi, H. Ogul⁹⁰ , Y. Onel , A. Penzo , C. Snyder, E. Tiras⁹¹








Johns Hopkins University, Baltimore, Maryland, USA

B. Blumenfeld , L. Corcodilos , J. Davis , A.V. Gritsan , L. Kang , S. Kyriacou , P. Maksimovic , M. Roguljic , J. Roskes , S. Sekhar , M. Swartz










The University of Kansas, Lawrence, Kansas, USA

A. Abreu , L.F. Alcerro Alcerro , J. Anguiano , S. Arteaga Escatel , P. Baringer , A. Bean , Z. Flowers , D. Grove , J. King , G. Krintiras , M. Lazarovits , C. Le Mahieu , J. Marquez , M. Murray , M. Nickel , M. Pitt , S. Popescu⁹² , C. Rogan , C. Royon , S. Sanders , C. Smith , G. Wilson










Kansas State University, Manhattan, Kansas, USA










B. Allmond , R. Gujju Gurunadha , A. Ivanov , K. Kaadze , Y. Maravin , J. Natoli , D. Roy , G. Sorrentino 

University of Maryland, College Park, Maryland, USA

A. Baden , A. Belloni , J. Bistany-riebman, Y.M. Chen , S.C. Eno , N.J. Hadley , S. Jabeen , R.G. Kellogg , T. Koeth , B. Kronheim, Y. Lai , S. Lascio , A.C. Mignerey , S. Nabili , C. Palmer , C. Papageorgakis , M.M. Paranjpe, E. Popova⁹³ , A. Shevelev , L. Wang

Massachusetts Institute of Technology, Cambridge, Massachusetts, USA













J. Bendavid , I.A. Cali , P.c. Chou , M. D'Alfonso , J. Eysermans , C. Freer , G. Gomez-Ceballos , M. Goncharov, G. Grosso, P. Harris, D. Hoang, D. Kovalskyi , J. Krupa , L. Lavezzo , Y.-J. Lee , K. Long , C. Mcginn, A. Novak , M.I. Park

C. Paus , C. Reissel , C. Roland , G. Roland , S. Rothman , G.S.F. Stephans ,
Z. Wang , B. Wyslouch , T. J. Yang 











University of Minnesota, Minneapolis, Minnesota, USA

B. Crossman , B.M. Joshi , C. Kapsiak , M. Krohn , D. Mahon , J. Mans ,
B. Marzocchi , M. Revering , R. Rusack , R. Saradhy , N. Strobbe 



















University of Nebraska-Lincoln, Lincoln, Nebraska, USA

K. Bloom , D.R. Claes , G. Haza , J. Hossain , C. Joo , I. Kravchenko , J.E. Siado ,
W. Tabb , A. Vagnerini , A. Wightman , F. Yan , D. Yu 

State University of New York at Buffalo, Buffalo, New York, USA

H. Bandyopadhyay , L. Hay , H.w. Hsia, I. Iashvili , A. Kalogeropoulos ,
A. Kharchilava , M. Morris , D. Nguyen , S. Rappoccio , H. Rejeb Sfar, A. Williams ,
P. Young 



























Northeastern University, Boston, Massachusetts, USA

G. Alverson , E. Barberis , J. Bonilla , B. Bylsma, M. Campana , J. Dervan, Y. Haddad ,
Y. Han , I. Israr , A. Krishna , J. Li , M. Lu , G. Madigan , R. Mccarthy ,
D.M. Morse , V. Nguyen , T. Orimoto , A. Parker , L. Skinnari , D. Wood 









Northwestern University, Evanston, Illinois, USA

J. Bueghly, S. Dittmer , K.A. Hahn , Y. Liu , M. McGinnis , Y. Miao , D.G. Monk ,
M.H. Schmitt , A. Taliercio , M. Velasco

















University of Notre Dame, Notre Dame, Indiana, USA

G. Agarwal , R. Band , R. Bucci, S. Castells , A. Das , R. Goldouzian , M. Hildreth ,
K.W. Ho , K. Hurtado Anampa , T. Ivanov , C. Jessop , K. Lannon , J. Lawrence ,
N. Loukas , L. Lutton , J. Mariano, N. Marinelli, I. Mcalister, T. McCauley , C. Mcgrady ,
C. Moore , Y. Musienko¹⁴ , H. Nelson , M. Osherson , A. Piccinelli , R. Ruchti ,
A. Townsend , Y. Wan, M. Wayne , H. Yockey, M. Zarucki , L. Zygalá 

The Ohio State University, Columbus, Ohio, USA

A. Basnet , M. Carrigan , L.S. Durkin , C. Hill , M. Joyce , M. Nunez Ornelas , K. Wei,
D.A. Wenzl, B.L. Winer , B. R. Yates 




















Princeton University, Princeton, New Jersey, USA

H. Bouchamaoui , K. Coldham, P. Das , G. Dezoort , P. Elmer , A. Frankenthal ,
B. Greenberg , N. Haubrich , K. Kennedy, G. Kopp , S. Kwan , D. Lange ,
A. Loeliger , D. Marlow , I. Ojalvo , J. Olsen , D. Stickland , C. Tully , L.H. Vage




University of Puerto Rico, Mayaguez, Puerto Rico, USA

S. Malik , R. Sharma

Purdue University, West Lafayette, Indiana, USA

A.S. Bakshi , S. Chandra , R. Chawla , A. Gu , L. Gutay, M. Jones , A.W. Jung ,
A.M. Koshy, M. Liu , G. Negro , N. Neumeister , G. Paspalaki , S. Piperov ,
V. Scheurer, J.F. Schulte , M. Stojanovic , J. Thieman , A. K. Viridi , F. Wang ,
A. Wildridge , W. Xie , Y. Yao 

Purdue University Northwest, Hammond, Indiana, USA

J. Dolen , N. Parashar , A. Pathak 

Rice University, Houston, Texas, USA





















D. Acosta , T. Carnahan , K.M. Ecklund , P.J. Fernández Manteca , S. Freed, P. Gardner,

F.J.M. Geurts , I. Krommydas , W. Li , J. Lin , O. Miguel Colin , B.P. Padley , R. Redjimi, J. Rotter , E. Yigitbasi , Y. Zhang 











University of Rochester, Rochester, New York, USA

A. Bodek , P. de Barbaro , R. Demina , J.L. Dulemba , A. Garcia-Bellido , O. Hindrichs , A. Khukhunaishvili , N. Parmar, P. Parygin⁹³ , R. Taus 
















Rutgers, The State University of New Jersey, Piscataway, New Jersey, USA

B. Chiarito, J.P. Chou , S.V. Clark , D. Gadkari , Y. Gershtein , E. Halkiadakis , M. Heindl , C. Houghton , D. Jaroslowski , S. Konstantinou , I. Laflotte , A. Lath , R. Montalvo, K. Nash, J. Reichert , H. Routray , P. Saha , S. Salur , S. Schnetzer, S. Somalwar , R. Stone , S.A. Thayil , S. Thomas, J. Vora , H. Wang 

University of Tennessee, Knoxville, Tennessee, USA

D. Ally , A.G. Delannoy , S. Fiorendi , S. Higginbotham , T. Holmes , A.R. Kanuganti , N. Karunarathna , L. Lee , E. Nibigira , S. Spanier 





Texas A&M University, College Station, Texas, USA

D. Aebi , M. Ahmad , T. Akhter , K. Androsov⁶¹ , O. Bouhali⁹⁴ , R. Eusebi , J. Gilmore , T. Huang , T. Kamon⁹⁵ , H. Kim , S. Luo , R. Mueller , D. Overton , D. Rathjens , A. Safonov 

Texas Tech University, Lubbock, Texas, USA

N. Akchurin , J. Damgov , N. Gogate , V. Hegde , A. Hussain , Y. Kazhykarim, K. Lamichhane , S.W. Lee , A. Mankel , T. Peltola , I. Volobouev 

Vanderbilt University, Nashville, Tennessee, USA

E. Appelt , Y. Chen , S. Greene, A. Gurrola , W. Johns , R. Kunnawalkam Elayavalli , A. Melo , F. Romeo , P. Sheldon , S. Tuo , J. Velkovska , J. Viinikainen 























University of Virginia, Charlottesville, Virginia, USA

B. Cardwell , H. Chung, B. Cox , J. Hakala , R. Hirosky , A. Ledovskoy , C. Neu 



















































Wayne State University, Detroit, Michigan, USA




























S. Bhattacharya , P.E. Karchin 

University of Wisconsin - Madison, Madison, Wisconsin, USA

A. Aravind, S. Banerjee , K. Black , T. Bose , E. Chavez , S. Dasu , P. Everaerts , C. Galloni, H. He , M. Herndon , A. Herve , C.K. Koraka , A. Lanaro, R. Loveless , J. Madhusudanan Sreekala , A. Mallampalli , A. Mohammadi , S. Mondal, G. Parida , L. Pétré , D. Pinna, A. Savin, V. Shang , V. Sharma , W.H. Smith , D. Teague, H.F. Tsoi , W. Vetens , A. Warden 

Authors affiliated with an institute or an international laboratory covered by a cooperation agreement with CERN

S. Afanasiev , V. Alexakhin , D. Budkouski , I. Golutvin[†] , I. Gorbunov , V. Karjavine , V. Korenkov , A. Lanev , A. Malakhov , V. Matveev⁹⁶ , V. Palichik , V. Perelygin , M. Savina , V. Shalaev , S. Shmatov , S. Shulha , V. Smirnov , O. Teryaev , N. Voytishin , B.S. Yuldashev⁹⁷, A. Zarubin , I. Zhizhin , G. Gavrillov , V. Golovtsov , Y. Ivanov , V. Kim⁹⁶ , P. Levchenko⁹⁸ , V. Murzin , V. Oreshkin , D. Sosnov , V. Sulimov , L. Uvarov , A. Vorobyev[†], Yu. Andreev , A. Dermenev , S. Gninenko , N. Golubev , A. Karneyev , D. Kirpichnikov , M. Kirsanov , N. Krasnikov , I. Tlisova , A. Toropin , T. Aushev , K. Ivanov , V. Gavrillov , N. Lychkovskaya , A. Nikitenko^{99,100} , V. Popov , A. Zhokin , R. Chistov⁹⁶ , M. Danilov⁹⁶ 

S. Polikarpov⁹⁶ , V. Andreev , M. Azarkin , M. Kirakosyan, A. Terkulov , E. Boos , V. Bunichev , M. Dubinin⁸⁵ , L. Dudko , A. Ershov , A. Gribushin , V. Klyukhin , O. Kodolova¹⁰⁰ , S. Obraztsov , S. Petrushanko , V. Savrin , A. Snigirev , V. Blinov⁹⁶, T. Dimova⁹⁶ , A. Kozyrev⁹⁶ , O. Radchenko⁹⁶ , Y. Skovpen⁹⁶ , V. Kachanov , D. Konstantinov , S. Slabospitskii , A. Uzunian , A. Babaev , V. Borshch , D. Druzhkin¹⁰¹ 

†: Deceased

¹Also at Yerevan State University, Yerevan, Armenia

²Also at TU Wien, Vienna, Austria

³Also at Ghent University, Ghent, Belgium

⁴Also at Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

⁵Also at Universidade Estadual de Campinas, Campinas, Brazil

⁶Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil

⁷Also at University of Chinese Academy of Sciences, Beijing, China

⁸Also at China Center of Advanced Science and Technology, Beijing, China

⁹Also at University of Chinese Academy of Sciences, Beijing, China

¹⁰Also at China Spallation Neutron Source, Guangdong, China

¹¹Now at Henan Normal University, Xinxiang, China

¹²Also at University of Shanghai for Science and Technology, Shanghai, China

¹³Now at The University of Iowa, Iowa City, Iowa, USA

¹⁴Also at an institute or an international laboratory covered by a cooperation agreement with CERN

¹⁵Also at Cairo University, Cairo, Egypt

¹⁶Also at Suez University, Suez, Egypt

¹⁷Now at British University in Egypt, Cairo, Egypt

¹⁸Also at Purdue University, West Lafayette, Indiana, USA

¹⁹Also at Université de Haute Alsace, Mulhouse, France

²⁰Also at Istinye University, Istanbul, Turkey

²¹Also at Tbilisi State University, Tbilisi, Georgia

²²Also at The University of the State of Amazonas, Manaus, Brazil

²³Also at University of Hamburg, Hamburg, Germany

²⁴Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

²⁵Also at Bergische University Wuppertal (BUW), Wuppertal, Germany

²⁶Also at Brandenburg University of Technology, Cottbus, Germany

²⁷Also at Forschungszentrum Jülich, Juelich, Germany

²⁸Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland

²⁹Also at HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary

³⁰Now at Universitatea Babeş-Bolyai - Facultatea de Fizica, Cluj-Napoca, Romania

³¹Also at MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

³²Also at HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

³³Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt

³⁴Also at Punjab Agricultural University, Ludhiana, India

³⁵Also at University of Visva-Bharati, Santiniketan, India

³⁶Also at Indian Institute of Science (IISc), Bangalore, India

³⁷Also at Amity University Uttar Pradesh, Noida, India

³⁸Also at IIT Bhubaneswar, Bhubaneswar, India

³⁹Also at Institute of Physics, Bhubaneswar, India

⁴⁰Also at University of Hyderabad, Hyderabad, India

⁴¹Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany

- ⁴²Also at Isfahan University of Technology, Isfahan, Iran
- ⁴³Also at Sharif University of Technology, Tehran, Iran
- ⁴⁴Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran
- ⁴⁵Also at Department of Physics, Faculty of Science, Arak University, ARAK, Iran
- ⁴⁶Also at Helwan University, Cairo, Egypt
- ⁴⁷Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy
- ⁴⁸Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy
- ⁴⁹Also at Università degli Studi Guglielmo Marconi, Roma, Italy
- ⁵⁰Also at Scuola Superiore Meridionale, Università di Napoli 'Federico II', Napoli, Italy
- ⁵¹Also at Fermi National Accelerator Laboratory, Batavia, Illinois, USA
- ⁵²Also at Lulea University of Technology, Lulea, Sweden
- ⁵³Also at Laboratori Nazionali di Legnaro dell'INFN, Legnaro, Italy
- ⁵⁴Also at Consiglio Nazionale delle Ricerche - Istituto Officina dei Materiali, Perugia, Italy
- ⁵⁵Also at Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France
- ⁵⁶Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia
- ⁵⁷Also at Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico
- ⁵⁸Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka
- ⁵⁹Also at Saegis Campus, Nugegoda, Sri Lanka
- ⁶⁰Also at National and Kapodistrian University of Athens, Athens, Greece
- ⁶¹Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland
- ⁶²Also at University of Vienna, Vienna, Austria
- ⁶³Also at Universität Zürich, Zurich, Switzerland
- ⁶⁴Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria
- ⁶⁵Also at Laboratoire d'Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, France
- ⁶⁶Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey
- ⁶⁷Also at Konya Technical University, Konya, Turkey
- ⁶⁸Also at Izmir Bakircay University, Izmir, Turkey
- ⁶⁹Also at Adiyaman University, Adiyaman, Turkey
- ⁷⁰Also at Bozok Universitetesi Rektörlüğü, Yozgat, Turkey
- ⁷¹Also at Marmara University, Istanbul, Turkey
- ⁷²Also at Milli Savunma University, Istanbul, Turkey
- ⁷³Also at Kafkas University, Kars, Turkey
- ⁷⁴Now at Istanbul Okan University, Istanbul, Turkey
- ⁷⁵Also at Hacettepe University, Ankara, Turkey
- ⁷⁶Also at Erzincan Binali Yildirim University, Erzincan, Turkey
- ⁷⁷Also at Istanbul University - Cerrahpasa, Faculty of Engineering, Istanbul, Turkey
- ⁷⁸Also at Yildiz Technical University, Istanbul, Turkey
- ⁷⁹Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom
- ⁸⁰Also at IPPP Durham University, Durham, United Kingdom
- ⁸¹Also at Monash University, Faculty of Science, Clayton, Australia
- ⁸²Also at Università di Torino, Torino, Italy
- ⁸³Also at Bethel University, St. Paul, Minnesota, USA
- ⁸⁴Also at Karamanoğlu Mehmetbey University, Karaman, Turkey

⁸⁵Also at California Institute of Technology, Pasadena, California, USA

⁸⁶Also at United States Naval Academy, Annapolis, Maryland, USA

⁸⁷Also at Ain Shams University, Cairo, Egypt

⁸⁸Also at Bingol University, Bingol, Turkey

⁸⁹Also at Georgian Technical University, Tbilisi, Georgia

⁹⁰Also at Sinop University, Sinop, Turkey

⁹¹Also at Erciyes University, Kayseri, Turkey

⁹²Also at Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania

⁹³Now at another institute or international laboratory covered by a cooperation agreement with CERN

⁹⁴Also at Texas A&M University at Qatar, Doha, Qatar

⁹⁵Also at Kyungpook National University, Daegu, Korea

⁹⁶Also at another institute or international laboratory covered by a cooperation agreement with CERN

⁹⁷Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan

⁹⁸Also at Northeastern University, Boston, Massachusetts, USA

⁹⁹Also at Imperial College, London, United Kingdom

¹⁰⁰Now at Yerevan Physics Institute, Yerevan, Armenia

¹⁰¹Also at Universiteit Antwerpen, Antwerpen, Belgium