

Vector boson production in association with jets at ATLAS and CMS (including Heavy Flavour jets)*

GIOVANNI PADOVANO ON BEHALF OF THE ATLAS AND CMS
COLLABORATIONS

Dipartimento di Fisica, Sapienza Università di Roma and INFN Roma
Piazzale Aldo Moro 5, 00185 Rome, Italy

Received December 8, 2024

Vector boson production in association with jets is among the Standard Model processes leaving the clearest signatures in high energy physics experiments. The latest results on W/Z +jets measurements from the ATLAS and CMS experiments at the LHC are presented. Particular attention is given to how these measurements can be used to extract relevant electroweak and QCD parameters, and also be re-interpreted to set limits on exotic models.

1. Introduction

Vector boson production is one of the most fundamental processes in particle physics and plays a crucial role in the Standard Model. These processes constitute relevant backgrounds in many particle physics analyses, covering both searches for exotic phenomena, Higgs boson physics measurements and Standard Model measurements, motivating the need of a high-precision estimate of their cross-sections. They also enable to determine QCD parameters and to probe the quark structure of the proton. Finally, the high-precision study of these processes enables to test the accuracy of several Monte-Carlo event generators, including matrix-element, parton shower and parton distribution functions (PDFs) setups. These proceedings present the latest W/Z +jets production cross-sections measurements, performed by the ATLAS and CMS experiments [1, 2] at the LHC [3]. The measurements are performed using the respective experimental pp collisions datasets of 140 fb^{-1} (ATLAS) and 138 fb^{-1} (CMS), recorded with a centre-of-mass energy of $\sqrt{s} = 13 \text{ TeV}$.

* Presented at “Diffraction and Low- x 2024”, Trabia (Palermo, Italy), September 8-14, 2024.




Fig. 1. Measurement of the $W + c$ integral fiducial cross-section by the CMS experiment. The measured value is compared with MADGRAPH5_aMC@NLO prediction using different PDFs sets [4].

2. Measurement of $W + c$ production cross-sections

The associated production of a W boson and a single charm quark ($W+c$) in proton-proton collisions at the LHC can be effectively used as a probe to study the s -quark content of the proton. This sensitivity originates from the dominance of the $sg \rightarrow W + c$ leading diagram over the Cabibbo-suppressed $dg \rightarrow W + c$ process.

2.1. CMS experiment measurements

The CMS Collaboration has measured the integral and differential cross-sections of the W boson production in association with a c -quark [4]. Events have been selected requiring one electron or one muon, originating from the W boson decay and a c -jet. Charm-jets have been tagged via the presence of a muon or a secondary vertex inside the jet.

The combined measured fiducial cross-section is:

$$\sigma(W + c) = 148.7 \pm 0.4(\text{stat.}) \pm 5.6(\text{syst.}) \text{ pb.} \quad (1)$$

as also shown in Fig. 1. Together with the integral fiducial cross-section, a measurement of the cross-sections ratio

$$R_c^\pm = \frac{\sigma(pp \rightarrow W^+ + c)}{\sigma(pp \rightarrow W^- + c)} \quad (2)$$

has been provided, resulting in

$$R_c^\pm = 0.950 \pm 0.005(\text{stat.}) \pm 0.010(\text{syst.}) \quad (3)$$

and compared with different predictions obtained assuming several models for the $s\bar{s}$ quark content of the proton.




Fig. 2. Observables measured by the ATLAS experiment in final states involving W boson in association with a c -hadron: (a) fiducial cross-section for the $W^- + D^+$ final state (b) fiducial cross-section ratio R_c^\pm [5].

The integral cross-section measurement has been complemented with differential cross-section measurements, as functions of the transverse momentum and the pseudo-rapidity of the lepton originating from the W boson decay. The measurements have also been repeated applying a parton-level unfolding procedure, consisting in an additional correction to account for the c -quark fragmentation and hadronisation process.

2.2. ATLAS experiment measurements

A similar study of the $W + c$ process has been published by the ATLAS Collaboration [5]. In this case, the charm-quark has been tagged via the direct detection of a charmed hadron, reconstructed through a secondary vertex fit. The W boson, instead, has similarly been reconstructed via the presence of an electron or a muon, together with missing transverse momentum in the final state. Four reconstructed charmed mesons are considered, specifically: $D^+ \rightarrow K^- \pi^+ \pi^+$, $D^{*+} \rightarrow D^0 \pi^+ \rightarrow (K^- \pi^+) \pi^+$ and their charge conjugate decays.

Fig. 2(a) shows the value of the measured fiducial cross-section for final states involving $W^- + D^+$, which resulted:

$$\sigma_{\text{fid}}(W^- + D^+) = 50.2 \pm 0.2(\text{stat.})^{+2.4}_{-2.3}(\text{syst.}) \text{ pb.} \quad (4)$$

Similar values have been obtained for the cross-sections of the other decay modes: $\sigma_{\text{fid}}(W^+ + D^-)$, $\sigma_{\text{fid}}(W^- + D^{*+})$, $\sigma_{\text{fid}}(W^+ + D^{*-})$. Additionally, the ratio of charm to anti-charm production cross-sections R_c^\pm has been studied to probe the $s\bar{s}$ quark-content asymmetry, as shown in Fig. 2(b).




Fig. 3. Measured fiducial cross-sections for (a) $Z + \geq 1b$ -jets, (b) $Z + \geq 2b$ -jets, (c) $Z + \geq 1c$ -jets, compared with several Monte-Carlo predictions with the relative flavour schemes [6].

3. Measurements of $Z + b, c$ production cross-sections

The production of a Z boson associated with heavy flavour quarks - namely b -quarks and c -quarks - in pp collisions is among the processes most commonly involved in the background events of many Higgs boson measurements or searches for new physics. The measurement of the production cross-section for $Z + b$ -jets and $Z + c$ -jets constitutes therefore an important benchmark to test the accuracy of the Monte Carlo generators predictions to be used in such analyses, and also provides tests of perturbative quantum chromodynamics (pQCD) and of the proton internal structure.

3.1. ATLAS experiment measurements

The ATLAS Collaboration has measured integral and differential production cross-sections of events containing a Z boson decaying into electrons or muons and produced in association with b - or c -jets [6].

Events have been categorised according to the associated jets multiplicity as $Z + \geq 1b$ -jets, $Z + \geq 2b$ -jets, $Z + \geq 1c$ -jets, and separate total and

differential cross-sections have been evaluated corresponding to these categories, as shown in Fig. 3. The $Z + \geq 1b$ -jets and $Z + \geq 2b$ -jets resulted optimally described by MADGRAPH5_aMC@NLO+PYTHIA8 with the choice of the 5 flavour scheme (FS). Instead, the $Z + \geq 1c$ -jets cross-section resulted optimally described by MADGRAPH5_aMC@NLO+PYTHIA8 with the 4 FS.

Some of the presented differential cross-section measurements have been optimised in bins of variables sensible to the possible presence of a valence-like charm quark in the proton (*intrinsic charm*). Data have been compared to Monte Carlo predictions obtained with PDFs choices testing different intrinsic charm hypotheses, without observing a significant evidence for its presence.

3.2. CMS experiment measurements

The integral and differential cross-sections of the $Z + b$ -jets production have been measured also by the CMS experiment and reported in Ref. [7]. Similarly to the ATLAS Collaboration measurement, events are categorised according to the multiplicity of b -jets associated with the Z boson productions: $Z + \geq 1 b$ -jets, $Z + \geq 2 b$ -jets and considering decays of the Z into electrons and muons.

The measured fiducial integral cross-sections are:

$$\begin{aligned}\sigma(Z \geq 1b\text{-jets}) &= 6.52 \pm 0.04(\text{stat.}) \pm 0.40(\text{syst.}) \pm 0.14(\text{theo.}) \text{ pb} \\ \sigma(Z \geq 2b\text{-jets}) &= 0.65 \pm 0.03(\text{stat.}) \pm 0.07(\text{syst.}) \pm 0.02(\text{theo.}) \text{ pb}.\end{aligned}\quad (5)$$

Normalised cross-sections are measured in bins of several kinematic variables and are used to test the accuracy of the corresponding Monte Carlo predictions. Two exemplary variables are reported in Fig. 4 where the cross-sections are differentiated in bins of vector boson transverse momentum p_{T}^Z and angular separation between the vector boson and the leading b -jet $\Delta R^{(Z,b\text{-jet})}$. The differential spectrum in p_{T}^Z is best described by the MADGRAPH5_aMC@LO, which is however yielding slight mismodelling at high values of $\Delta R^{(Z,b\text{-jet})}$, thus showing the complexity of the modelling of angular variables.

4. Other vector bosons in association with jets measurements

4.1. Measurement of $p_{\text{T}}^{\text{miss}} + \text{jets}$ production cross-sections with the ATLAS experiment

The ATLAS Collaboration has also published a measurement of inclusive and differential cross-sections for the production of events with missing




Fig. 4. Normalised differential cross-sections measurements for the $Z + \geq 1$ b -jets production in bins of (a) vector boson transverse momentum (b) angular separation between the vector boson and the leading b -jet [7].

transverse momentum in association with jets [8]. The measurement includes the determination of di-jet distributions in a region in which vector boson fusion processes are enhanced. Additionally, measurements of the $Z \rightarrow \nu\nu$ process are provided.

Ratios between the measured $p_T^{\text{miss}} + \text{jets}$ cross-section distributions are derived, to take advantage of the cancellations in modelling effects and some major systematic uncertainties, and are used to constraint new physics phenomenological models involving Dark Matter particles.

One common benchmark for Dark Matter involves extending the Standard Model with an additional $U(1)$ symmetry group, associated with a Z' vector boson, and involving Dark Matter Dirac fermions χ [9]. Fig. 5(a) shows a recasted contour in the $m_{Z'} - m_\chi$ plane, yielding an exclusion competitive with the previous dedicated detector-level ATLAS search [10]. Additionally, the 2HDM+ a model [11] has been studied involving an additional Higgs doublet, along with a pseudoscalar a which couples to Dark Matter. Fig. 5(b) shows the exclusion obtained for this model, in the plane $(m_a, \tan\beta)$, where m_a is the mass of the pseudoscalar and $\tan\beta$ is a parameter of the model.




Fig. 5. Exclusion limits at 95%, obtained recasting the $p_T^{\text{miss}} + \text{jets}$ cross-sections measurements for: (a) a simplified Dark Matter model involving a Dark Matter Dirac fermion χ (b) the 2HDM+a model [8].

4.2. Measurement of the $W \rightarrow cq/W \rightarrow q\bar{q}'$ branching fraction ratio with the CMS experiment

Vector boson production can also be exploited to perform flavour physics measurements. The CMS Collaboration has recently published a measurement of the W boson hadronic decay branching fraction ratio $R_c^W = \mathcal{B}(W \rightarrow cq)/\mathcal{B}(W \rightarrow q\bar{q}')$ [12]. From this measurement, the sum of squared elements in the second row of the Cabibbo-Kobayashi-Maskawa matrix, 0.970 ± 0.041 , and the element $|V_{cs}| = 0.959 \pm 0.021$ have been derived.

5. Conclusions

These proceedings summarised the most recent measurements on vector boson production in association with jets, obtained by the ATLAS and CMS Collaborations at the Large Hadron Collider. Several analysis strategies have been described, evidencing how these measurements can be used to extract electroweak and QCD parameters, and can also be exploited to draw exclusion limits on exotic new physics models.

REFERENCES

- [1] ATLAS Collaboration, *The ATLAS Experiment at the CERN Large Hadron Collider*, JINST 3 (2008) S08003, DOI 10.1088/1748-0221/3/08/S08003
- [2] CMS Collaboration, *The CMS experiment at the CERN LHC*, JINST 3 (2008) S08004, DOI 10.1088/1748-0221/3/08/S08004

- [3] L. Evans and P. Bryant (editors), *LHC Machine*, JINST 3 (2008) S08001, DOI 10.1088/1748-0221/3/08/S08001
- [4] CMS Collaboration, *Measurement of the production cross-section for a W boson in association with a charm quark in proton-proton collisions at $\sqrt{s} = 13$ TeV*, Eur. Phys. J. C 84 (2024) 27, doi:10.1140/epjc/s10052-023-12258-4
- [5] ATLAS Collaboration, *Measurement of the production of a W boson in association with a charmed hadron in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, Phys. Rev. D 108 (2023) 032012, doi:10.1103/PhysRevD.108.032012
- [6] ATLAS Collaboration, *Measurements of the production cross-section for a Z boson in association with b - or c -jets in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, Eur. Phys. J. C 84 (2024) 984, doi:10.1140/epjc/s10052-024-13159-w
- [7] CMS Collaboration, *Measurement of the production cross-section for $Z+b$ jets in proton-proton collisions at $\sqrt{s} = 13$ TeV*, Phys. Rev. D 105 (2022) 092014, doi:10.1103/PhysRevD.105.092014
- [8] ATLAS Collaboration, *Differential cross-sections for events with missing transverse momentum and jets measured with the ATLAS detector in 13 TeV proton-proton collisions*, JHEP 08 (2024) 223, doi:10.1007/JHEP08(2024)223
- [9] J. Abdallah, et al., *Simplified Models for Dark Matter Searches at the LHC*, Phys. Dark Univ. 9-10 (2015) 8-23, doi:10.1016/j.dark.2015.08.001
- [10] ATLAS Collaboration, *Search for new phenomena in events with an energetic jet and missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, Phys. Rev. D 103 (2021) 112006, doi:10.1103/PhysRevD.103.112006
- [11] M. Bauer et al., *Simplified dark matter models with two Higgs doublets: I. Pseudoscalar mediators*, JHEP 05 (2017) 138, doi:10.1007/JHEP05(2017)138
- [12] CMS Collaboration, *Measurement of the $W \rightarrow cq/W \rightarrow q\bar{q}'$ decay branching fraction ratio in proton-proton collisions at $\sqrt{s} = 13$ TeV*, CMS-PAS-SMP-24-009, <https://cds.cern.ch/record/2904877>