

CERN-EP-2025-011
05 February 2025
arXiv:2502.01111

Branching fraction measurement of the decay $B^+ \rightarrow \psi(2S)\phi(1020)K^+$

LHCb collaboration[†]

Abstract

The branching fraction of the decay $B^+ \rightarrow \psi(2S)\phi(1020)K^+$, relative to the topologically similar decay $B^+ \rightarrow J/\psi\phi(1020)K^+$, is measured using proton-proton collision data collected by the LHCb experiment at center-of-mass energies of 7, 8, and 13 TeV, corresponding to an integrated luminosity of 9 fb^{-1} . The ratio is found to be $0.061 \pm 0.004 \pm 0.009$, where the first uncertainty is statistical and the second systematic. Using the world-average branching fraction for $B^+ \rightarrow J/\psi\phi(1020)K^+$, the branching fraction for the decay $B^+ \rightarrow \psi(2S)\phi(1020)K^+$ is found to be $(3.0 \pm 0.2 \pm 0.5 \pm 0.2) \times 10^{-6}$, where the first uncertainty is statistical, the second systematic, and the third is due to the branching fraction of the normalization channel.

Submitted to Phys. Rev. D

© 2025 CERN for the benefit of the LHCb collaboration. CC BY 4.0 licence.

[†]Authors are listed at the end of this paper.

1 Introduction

The $B^+ \rightarrow \psi(2S)\phi K^+$ decay proceeds through the same quark transition $b \rightarrow c\bar{c}s$ as the topologically similar $B^+ \rightarrow J/\psi\phi K^+$ decay.¹ In the latter, states compatible with exotic mesons have been observed, such as the $\chi_{c1}(4140)$ state [1–4] and the $T_{c\bar{c}s1}(4000)$ state [5].² The $B^+ \rightarrow \psi(2S)\phi K^+$ phase space is about 80 times smaller than that of the $B^+ \rightarrow J/\psi\phi K^+$ decay, while the branching fraction $\mathcal{B}(B^+ \rightarrow \psi(2S)\phi K^+)$ is about 15 times smaller than that of the $B^+ \rightarrow J/\psi\phi K^+$ decay [7]. The average amplitude of the former process is therefore greater than that of the latter process. Thus, the $B^+ \rightarrow \psi(2S)\phi K^+$ decay provides the potential to make observations of high-mass resonant contributions that are complementary to those observed in the decay $B^+ \rightarrow J/\psi\phi K^+$, such as the $\chi_c(4700)$ state [8, 9].

This paper presents a measurement of the branching fraction of the decay $B^+ \rightarrow \psi(2S)\phi K^+$ relative to the normalization decay $B^+ \rightarrow J/\psi\phi K^+$. The intermediate resonances are reconstructed as $\psi(2S) \rightarrow \mu^+\mu^-$, $J/\psi \rightarrow \mu^+\mu^-$, and $\phi \rightarrow K^+K^-$. This analysis uses the full Run 1 (2011–2012) and Run 2 (2015–2016, 2017, and 2018) LHCb data samples of proton-proton (pp) collisions at center-of-mass energies of 7, 8, and 13 TeV, corresponding to an integrated luminosity of about 9 fb^{-1} . The branching fraction of $B^+ \rightarrow \psi(2S)\phi K^+$, calculated using the $\mathcal{B}(B^+ \rightarrow J/\psi\phi K^+)$ value provided by the PDG [6], is also reported. This LHCb result is compared with a measurement of the branching fraction of $B^+ \rightarrow \psi(2S)\phi K^+$ reported by the CMS collaboration [7].

2 Detector and simulation

The LHCb detector [10, 11] is a single-arm forward spectrometer covering the pseudorapidity range $2 < \eta < 5$, designed for the study of particles containing b or c quarks. The detector includes a high-precision tracking system consisting of a silicon-strip vertex detector surrounding the pp interaction region [12], a large-area silicon-strip detector located upstream of a dipole magnet with a bending power of about 4 T m , and three stations of silicon-strip detectors and straw drift tubes [13, 14] placed downstream of the magnet. The tracking system provides a measurement of the momentum, p , of charged particles with a relative uncertainty that varies from 0.5% at low momentum to 1.0% at $200 \text{ GeV}/c$. The minimum distance of a track to a primary vertex (PV), the impact parameter (IP), is measured with a resolution of $(15 + 29/p_T) \mu\text{m}$, where p_T is the component of the momentum transverse to the beam, in GeV/c . Different types of charged hadrons are distinguished using information from two ring-imaging Cherenkov detectors [15]. Photons, electrons, and hadrons are identified by a calorimeter system consisting of scintillating-pad and preshower detectors, an electromagnetic and a hadronic calorimeter. Muons are identified by a system composed of alternating layers of iron and multi-wire proportional chambers [16]. The online event selection is performed by a trigger [17], which consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a software stage, which applies a full event reconstruction.

¹Throughout this paper inclusion of the charge-conjugated decays is implied and the $\phi(1020)$ is referred to as ϕ .

²In this paper the new PDG naming conventions [6] for exotic hadrons are used. The states formerly known as the $X(4140)$, $Z_{cs}^+(4000)$ and $X(4700)$ are now referred to as the $\chi_{c1}(4140)$, $T_{c\bar{c}s1}(4000)$ and $\chi_c(4700)$, respectively.

Simulation is required to model the effects of the detector acceptance and the selection requirements. In the simulation, pp collisions are generated using PYTHIA [18] with a specific LHCb configuration [19]. Decays of unstable particles are described by EVTGEN [20], in which final-state radiation is generated using PHOTOS [21]. The interaction of the generated particles with the detector, and its response, are implemented using the GEANT4 toolkit [22] as described in Ref. [23].

3 Branching fraction measurement

The ratio of branching fractions \mathcal{R}_{BF} is measured as

$$\mathcal{R}_{\text{BF}} \equiv \frac{\mathcal{B}(B^+ \rightarrow \psi(2S)\phi K^+)}{\mathcal{B}(B^+ \rightarrow J/\psi\phi K^+)} = \frac{N_{\text{Signal}} F_{\text{Signal}} \epsilon_{\text{Norm}}}{N_{\text{Norm}} F_{\text{Norm}} \epsilon_{\text{Signal}}} \frac{\mathcal{B}(J/\psi \rightarrow \mu^+\mu^-)}{\mathcal{B}(\psi(2S) \rightarrow \mu^+\mu^-)}, \quad (1)$$

where $\mathcal{B}(J/\psi \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033)\%$ and $\mathcal{B}(B^+ \rightarrow J/\psi\phi K^+) = (5.0 \pm 0.4) \times 10^{-5}$ are the world-average branching fractions [6]. Assuming lepton flavor universality of the electroweak interaction, the branching fraction of the $\psi(2S) \rightarrow \mu^+\mu^-$ decay can be replaced by that of the $\psi(2S) \rightarrow e^+e^-$ decay, equal to $(0.793 \pm 0.017)\%$ [6], which has an uncertainty approximately three times smaller. The number of B^+ decays for the signal (normalization) mode with a charmonium decay combined with three kaons, including sidebands for the ϕ candidates in the K^+K^- mass spectra, is denoted by $N_{\text{Signal(Norm)}}$. This is measured by fitting the B^+ meson mass distribution. The fraction of candidates in the K^+K^- invariant mass spectrum attributed to the ϕ decays is denoted by $F_{\text{Signal(Norm)}}$, and is obtained by fitting the K^+K^- mass distribution. Finally, $\epsilon_{\text{Signal(Norm)}}$ is the efficiency of the signal (normalization) channel, and is obtained from simulation.

3.1 Selection

Candidate $B^+ \rightarrow \psi(2S)\phi K^+$ signal and $B^+ \rightarrow J/\psi\phi K^+$ normalization decays are selected using a hardware trigger, a two-stage software trigger, and offline event reconstruction criteria followed by two Boosted Decision Tree (BDT) [24, 25] classifiers that suppress background significantly. Selection criteria are identical for both channels, with the exception of the dimuon mass windows for the ψ , where ψ denotes either the $\psi(2S)$ or the J/ψ meson.

The hardware trigger selects events with at least one high- p_{T} muon candidate later associated with the ψ in the B^+ decay or selects events independently of any track used in the decay. The first stage of the software trigger selects a pair of oppositely charged muons with high invariant mass or a muon with high p_{T} and high χ_{IP}^2 for each track (or composite particle). The variable χ_{IP}^2 is defined as the difference in the vertex-fit χ^2 of the primary-vertex fit reconstructed with and without the track under consideration. The second stage of the software trigger selects pairs of muons with invariant mass within $120 \text{ MeV}/c^2$ of the ψ mass that form vertices which are significantly detached from the closest primary vertex.

Offline, B^+ meson candidates are formed from combinations of ψ candidates with three kaons where one of the K^+K^- pairs has an invariant mass between $1000 \text{ MeV}/c^2$ and $1040 \text{ MeV}/c^2$. Kaon candidates are required to have p_{T} larger than $100 \text{ MeV}/c$, $\chi_{\text{IP}}^2 > 1$, and to have particle-identification signatures compatible with the kaon hypothesis. The

ϕ candidates are formed from pairs of oppositely charged kaons that originate from a common vertex and have p_T greater than 200 MeV/ c . The B^+ candidates are required to have good vertices, to have invariant masses within 60 MeV/ c^2 of the nominal B^+ mass, and to have momentum vectors aligned with the B^+ line of flight. A vertex fit is performed constraining the ψ mass to its nominal value as given by the PDG [6] and requiring the B^+ momentum vector to point back to its primary vertex. Candidates satisfying these criteria are retained for further consideration.

Background is suppressed using two BDT classifiers [24, 25] implemented in the TMVA toolkit [26]. The first classifier is trained using fully simulated $B^+ \rightarrow \psi(2S)\phi K^+$ samples as a proxy for the signal and events from the far-upper sideband ($5.8 < M(\psi(2S)\phi K^+) < 6.0$ GeV/ c^2) as a proxy for the background. The variables with highest discrimination power are used as input to the classifier. These include: the p_T and χ_{IP}^2 of the kaons, muons, ψ , ϕ , and the B^+ candidates, the cosine of the angle between the B^+ momentum and the vector connecting the PV and the B^+ decay vertex, the sum of the χ_{IP}^2 of all final-state particles with respect to the PV, the vertex χ^2 of the B^+ candidate, and the B^+ decay time and its significance. The chosen BDT threshold corresponds to a signal efficiency close to 90% and a background rejection of about 99%.

After all previous steps, more than one $B^+ \rightarrow \psi(2S)\phi K^+$ candidate in a single event can satisfy the selection criteria. In the majority of cases (around 30% and 3% of events in the signal and normalization channels, respectively), this occurs when the masses of the two K^+K^- combinations in a single $B^+ \rightarrow \psi K^+K^-K^+$ candidate fall within the nominal ϕ window. Combinatorial background may also produce additional candidates in a single event, though on a smaller scale. To retain one B^+ candidate per event, a second BDT is trained. The input variables include properties of the B^+ meson, such as the χ_{IP}^2 , the decay-time significance, the decay-vertex quality, the cosine of the angle between its momentum vector and the trajectory. Additionally, the input variables include the significance of the maximum distance between the trajectories of final-state particles. The candidate with the highest BDT output in each event is selected and its mass is used in the B^+ mass fit described below. The correlation between the mass of the B^+ candidate and the BDT output is found to be negligible. When the masses of both K^+K^- pairs in a $B^+ \rightarrow \psi K^+K^-K^+$ candidate fall within the nominal ϕ window, both K^+K^- combinations are used in the fit to determine F_{Signal} (F_{Norm}).

3.2 Mass fits

Unbinned maximum-likelihood mass fits are performed separately for the $B^+ \rightarrow \psi(2S)\phi K^+$ and $B^+ \rightarrow J/\psi\phi K^+$ samples, and for each data-taking period as the detector operating-conditions varied for each interval. For both the signal and normalization channels, the B^+ peak is modeled by a double-sided Crystal Ball function (DSCB) [27], while the background is described by an exponential function. The DSCB tail parameters are obtained from simulation and are fixed in the fits to the data. The yields of signal and background, the signal peak position M_{B^+} and width σ_{B^+} , and the parameters describing the exponential backgrounds, are floated in the fits. Mass distributions of $B^+ \rightarrow \psi(2S)\phi K^+$ and $B^+ \rightarrow J/\psi\phi K^+$ candidates are shown in Figs. 1 and 2, respectively, along with the results of the fits. The fit parameters of each data taking periods are reported in Table 1. The yields of the signal and normalization models determined from the fits over the combined samples are 289 ± 19 and $35\,315 \pm 237$ events, respectively, which




Figure 1: Invariant-mass distributions of $B^+ \rightarrow \psi(2S)\phi K^+$ candidates with fit projections for (top left) Run 1, (top right) 2015–2016, (bottom left) 2017, and (bottom right) 2018 samples.

are compatible with the sum of the yields obtained independently, providing additional confidence in the consistency of these values.

Although the samples are selected with K^+K^- invariant masses in the 1000–1040 MeV/c^2 mass window, significant non- ϕ contributions are present. The fraction of ϕ candidates is obtained from a binned χ^2 fit to the extended K^+K^- mass distribution of 1000–1060 MeV/c^2 . The selection criteria are the same as for the B^+ samples except for this relaxed K^+K^- mass range. In these fits, background is statistically subtracted by means of the *sPlot* technique [28], using the results of the fit to the B^+ mass distributions. The mass distribution of the ϕ component is described by a P-wave Breit–Wigner function, convolved with a Gaussian function with resolution fixed to 1 MeV/c^2 (taken from simulation), while that of the non- ϕ component is described by an Argus function [29]. As the yields of signal in the disjoint data samples are small while the event-selection efficiency does not vary much in the K^+K^- mass window, these fits are performed over the integrated Run 1 + Run 2 data samples. Therefore, only one $F_{\text{Signal(Norm)}}$ is calculated for each of the signal and normalization channels and used for all disjoint data samples. The K^+K^- mass fit results are shown in Fig. 3, in which the different line shapes of the non- ϕ component are mainly due to the different phase space of the signal and normalization modes. The fractions of ϕ contribution within the nominal K^+K^- mass window are determined to be $F_{\text{Signal}} = 0.73 \pm 0.10$ and $F_{\text{Norm}} = 0.91 \pm 0.01$, where the uncertainties are statistical only.




Figure 2: Invariant-mass distribution of $B^+ \rightarrow J/\psi\phi K^+$ candidates with fit projections for (top left) Run 1, (top right) 2015–2016, (bottom left) 2017, and (bottom right) 2018 samples.

Table 1: Results of the fit to the invariant mass of signal and normalization decays.

Period	Parameter	$B^+ \rightarrow \psi(2S)\phi K^+$	$B^+ \rightarrow J/\psi\phi K^+$
Run 1	M_{B^+} [MeV/ c^2]	5279.45 ± 0.40	5279.30 ± 0.07
	σ_{B^+} [MeV/ c^2]	2.04 ± 0.41	4.38 ± 0.07
	$N_{\text{Signal(Norm)}}$	43 ± 7	6835 ± 97
2015–2016	M_{B^+} [MeV/ c^2]	5279.79 ± 0.22	5279.28 ± 0.06
	σ_{B^+} [MeV/ c^2]	1.60 ± 0.23	4.46 ± 0.07
	$N_{\text{Signal(Norm)}}$	95 ± 11	9772 ± 129
2017	M_{B^+} [MeV/ c^2]	5279.40 ± 0.22	5279.36 ± 0.06
	σ_{B^+} [MeV/ c^2]	1.33 ± 0.23	4.26 ± 0.07
	$N_{\text{Signal(Norm)}}$	64 ± 9	8515 ± 118
2018	M_{B^+} [MeV/ c^2]	5279.65 ± 0.19	5279.27 ± 0.06
	σ_{B^+} [MeV/ c^2]	1.48 ± 0.16	4.29 ± 0.06
	$N_{\text{Signal(Norm)}}$	90 ± 11	10196 ± 128




Figure 3: Distributions of the background-subtracted K^+K^- mass spectra from the full (Run 1 + Run 2) data sample for (left) $B^+ \rightarrow \psi(2S)\phi K^+$ and (right) $B^+ \rightarrow J/\psi\phi K^+$ decays.

3.3 Efficiencies

Particle-identification efficiencies are obtained from calibration data samples [30]. The remaining efficiencies, for both the signal and normalization decay modes, are obtained from simulated samples and calculated separately for each data-taking period. The normalization simulation sample, generated as pure phase space, is weighted to account for the structures in the $\psi\phi$ and ψK^+ masses observed in data. This correction is not applied to the signal sample, as no significant intermediate structure is seen given its smaller size. A systematic uncertainty is estimated to address possible residual effects. In the normalization sample, per-candidate weights are calculated by training a gradient-boosted decision-tree reweighter, GBReweighter, implemented in `hep_ml` [31]. The data distributions used here have the background subtracted statistically by means of the *sPlot* technique [28], using the results of the fit to the $B^+ \rightarrow J/\psi\phi K^+$ invariant-mass distribution.

Two factors distort the distribution of events in phase space (the ψ, ϕ, K^+ Dalitz plot), especially at the edges. First, the resolution of the reconstructed B^+ invariant mass cannot be neglected. To mitigate this problem, the masses of the B^+ and ψ candidates are constrained to their nominal values. Second, the natural line width of the ϕ meson cannot be neglected. Therefore, the simulated samples are reweighted in the $(s_{\psi K}, \cos(\theta))$ space, where s_{XY} denotes the invariant-mass squared of the XY system and θ is the angle between the ψ and ϕ , in the ϕK^+ rest frame, calculated as

$$\cos(\theta) = \frac{(m_{B^+}^2 - s_{\phi K} - m_{\psi}^2)(s_{\phi K} + m_{\phi}^2 - m_K^2) + 2s_{\phi K}(m_{\psi}^2 + m_{\phi}^2 - s_{\psi\phi})}{\lambda^{\frac{1}{2}}(m_{B^+}^2, s_{\phi K}, m_{\psi}^2) \lambda^{\frac{1}{2}}(s_{\phi K}, m_{\phi}^2, m_K^2)}, \quad (2)$$

where m_{B^+} , m_{ψ} and m_{ϕ} are the reconstructed mass of the B^+ , ψ and ϕ mesons, respectively, m_K is the known K^+ mass [6], and $\lambda^{\frac{1}{2}}(x, y, z)$ is defined as

$$\lambda^{\frac{1}{2}}(x, y, z) \equiv \sqrt{(x - y - z)(x - y + z) - 4yz}. \quad (3)$$

As this parametrization is not directly correlated with the ϕ mass, and $-1 < \cos(\theta) < 1$ by definition, the weighting at the limits of phase space is better behaved than if the usual Dalitz plot variables are used.

Table 2: Ratios of efficiencies for the signal and normalization channels for each data-taking period, as described in the text, where the uncertainties are statistical only.

Period	$\epsilon_{\text{Signal}}/\epsilon_{\text{Norm}}$
Run 1	0.823 ± 0.004
2015–2016	0.801 ± 0.007
2017	0.790 ± 0.004
2018	0.834 ± 0.003

Ratios of efficiencies for the signal and normalization channels are derived from weighted, simulated data for each running period and are reported in Table 2. Although the hardware and first-stage software triggers are more efficient for signal channel decays, the second-stage software trigger and offline selection criteria are more efficient for the normalization channel decays. This results from the much smaller phase space for the kaons in the signal channel and the use of the track χ_{IP}^2 to exclude candidates with kaon momentum and B^+ flight directions consistent with each other.

3.4 Systematic uncertainties

Six sources of systematic uncertainty are discussed. Others were considered, but judged to be negligible compared to the statistical and overall systematic uncertainties reported.

The model used to describe the signal and background shapes may affect the results. Where the baseline fit uses a DSCB function to model the signal, a Johnson S_U function [32] is used as an alternative model to fit the data. Similarly, the baseline exponential background shape is replaced with a second order Chebychev polynomial. Using the full dataset, results obtained with Johnson S_U show a difference of $\sim 2\%$ in both signal and normalization samples. Added in quadrature, the total systematic uncertainty associated with the signal and background shapes is $\sim 3\%$.

The fractions of ϕ contributions in the nominal $m(K^+K^-)$ signal windows, F_{Signal} and F_{Norm} , are also subject to fluctuation. Their statistical uncertainties are propagated to the branching fraction ratio measurements. These lead to a systematic effect of around 14%, almost entirely associated with the signal channel. This is the dominant systematic uncertainty.

The Breit–Wigner function used to model the ϕ mass distribution is not corrected for the $B^+ \rightarrow \psi K^- K^+ K^+$ four-body phase-space shape. Fitting the reweighted simulation samples with the same procedure yields a 3% deviation in the ratio of the fraction of ϕ events between signal and normalization samples.

The limited sizes of simulation samples lead to statistical uncertainties in the efficiencies. These are propagated to the final results as systematic uncertainties, explicitly accounting for the weighting procedure discussed below. The relative uncertainty from this source varies between 3% and 4%.

The weighting procedure accounting for resonant structure in phase space requires a set of four parameters as input to train the gradient-boosted decision-tree reweighter [31]. On trialing a large number of alternative parameter sets, the distribution of results shows a negligible spread. The distribution of results shows a negligible spread. Therefore, no

Table 3: Systematic uncertainties on the ratio of branching fractions of the decays $B^+ \rightarrow \psi(2S)\phi K^+$ and $B^+ \rightarrow J/\psi\phi K^+$, by origin and data-taking period.

Period	Mass model	ϕ fraction	Phase-space correction	Simulation statistics	Signal efficiency correction	External inputs [6]	Total
Run 1	3%	14%	3%	4%	2%	2%	15%
2015–2016	3%	14%	3%	3%	1%	2%	15%
2017	3%	14%	3%	3%	3%	2%	15%
2018	3%	14%	3%	3%	4%	2%	15%
Combined	3%	14%	3%	1%	1%	2%	15%

systematic uncertainty is associated with weighting the normalization channel. As the weighting procedure is not applied to the signal model in the baseline fit, the associated systematic uncertainty is estimated by taking the difference between the efficiency with and without this weighting. Uncertainties of 2%, 1%, 3% and 4% are found for the Run 1, 2015–2016, 2017 and 2018 samples, respectively.

The branching fractions $\mathcal{B}(J/\psi \rightarrow \mu^+\mu^-)$ and $\mathcal{B}(\psi(2S) \rightarrow \mu^+\mu^-)$ are required to calculate the relative branching fraction $\mathcal{B}(B^+ \rightarrow \psi(2S)\phi K^+)/\mathcal{B}(B^+ \rightarrow J/\psi\phi K^+)$, and the branching fraction of the $B^+ \rightarrow J/\psi\phi K^+$ is additionally required for the $\mathcal{B}(B^+ \rightarrow \psi(2S)\phi K^+)$ calculation. These values are taken from the PDG [6] and their uncertainties are propagated to the final results. The contribution to the systematic uncertainty of the branching fraction ratio is $\sim 2\%$ and the corresponding contribution to the $B^+ \rightarrow \psi(2S)\phi K^+$ branching fraction is $\sim 8\%$.

Table 3 summarizes the systematic uncertainties. With the exception of the uncertainties due to simulation sample sizes and signal weighting, systematic uncertainties are assumed to be fully correlated throughout all data-taking periods.

4 Results and summary

Using the yields, the ϕ fractions and efficiency ratios reported in Tables 1 and 2, respectively, the ratio of branching fractions between the $B^+ \rightarrow \psi(2S)\phi K^+$ and $B^+ \rightarrow J/\psi\phi K^+$ decays is calculated. The results are reported in Table 4 and illustrated in Fig. 4. The derived $B^+ \rightarrow \psi(2S)\phi K^+$ branching fraction results are illustrated in Fig. 5. Results for each data-taking period are combined using the Best Linear Unbiased Estimator method BLUE [33], which accounts for correlations among systematic effects. The reported results are consistent throughout the data-taking periods.

The combined value obtained for the ratio of branching fractions is

$$\mathcal{R}_{\text{BF}} \equiv \frac{\mathcal{B}(B^+ \rightarrow \psi(2S)\phi K^+)}{\mathcal{B}(B^+ \rightarrow J/\psi\phi K^+)} = 0.061 \pm 0.004 \pm 0.009,$$

where the first uncertainty is statistical and the second is systematic. The corresponding value for the $B^+ \rightarrow \psi(2S)\phi K^+$ branching fraction is

$$\mathcal{B}(B^+ \rightarrow \psi(2S)\phi K^+) = (3.0 \pm 0.2 \pm 0.5 \pm 0.2) \times 10^{-6},$$

Table 4: Ratio of branching fractions \mathcal{R}_{BF} for each data-taking period and for the combined sample, where the first uncertainty is statistical and the second is systematic.

Period	\mathcal{R}_{BF}
Run 1	$0.046 \pm 0.008 \pm 0.007$
2015–2016	$0.072 \pm 0.009 \pm 0.011$
2017	$0.056 \pm 0.008 \pm 0.008$
2018	$0.064 \pm 0.008 \pm 0.010$
Combined	$0.061 \pm 0.004 \pm 0.009$




Figure 4: Ratio \mathcal{R}_{BF} of branching fractions for the decays $B^+ \rightarrow \psi(2S)\phi K^+$ and $B^+ \rightarrow J/\psi\phi K^+$ for each data-taking period and the confidence level region at one standard deviation for the combined result in green. The inner error bars (in blue) and the outer error bars (in red) delimit the statistical and total (statistical plus systematic, added in quadrature) uncertainties, respectively.

where the third uncertainty is associated with the uncertainty of the branching fraction of the normalization channel. The result is compatible with the CMS measurement $(4.0 \pm 0.4 \pm 0.6 \pm 0.2) \times 10^{-6}$ [7].

The phase space in $B^+ \rightarrow \psi(2S)\phi K^+$ decays is about 80 times smaller than that in $B^+ \rightarrow J/\psi\phi K^+$ decays. Considering Fermi's golden rule that the transition rate is proportional to the phase space times the magnitude of the amplitude squared, one may conclude that the average amplitude squared for the $B^+ \rightarrow \psi(2S)\phi K^+$ decay is approximately 5 times larger than that for the $B^+ \rightarrow J/\psi\phi K^+$ decay. This suggests the presence of resonant amplitudes in the $B^+ \rightarrow \psi(2S)\phi K^+$ decay that enhance its average decay rate. States such as the $\chi_c(4700)$ [8,9] have the right masses and quark content to play this role, so an amplitude analysis made possible with an increased data sample size represents a promising avenue for further investigation.




Figure 5: Branching fraction of the decay $B^+ \rightarrow \psi(2S)\phi K^+$ for each data-taking period and the confidence level region at one standard deviation for the combined result in green. The inner error bars (in blue) and the outer error bars (in red) delimit the statistical and total (statistical plus systematic, added in quadrature) uncertainties, respectively. The CMS result [7] is included for comparison.

Acknowledgements

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: ARC (Australia); CAPES, CNPq, FAPERJ and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); NWO (Netherlands); MNiSW and NCN (Poland); MCID/IFA (Romania); MICIU and AEI (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); DOE NP and NSF (USA). We acknowledge the computing resources that are provided by ARDC (Australia), CBPF (Brazil), CERN, IHEP and LZU (China), IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (Netherlands), Polish WLCG (Poland), IFIN-HH (Romania), PIC (Spain), CSCS (Switzerland), and GridPP (United Kingdom). We are indebted to the communities behind the multiple open-source software packages on which we depend. Individual groups or members have received support from Key Research Program of Frontier Sciences of CAS, CAS PIFI, CAS CCEPP, Fundamental Research Funds for the Central Universities, and Sci. & Tech. Program of Guangzhou (China); Minciencias (Colombia); EPLANET, Marie Skłodowska-Curie Actions, ERC and NextGenerationEU (European Union); A*MIDEX, ANR, IPhU and Labex P2IO, and Région Auvergne-Rhône-Alpes (France); Alexander-von-Humboldt Foundation (Germany); ICSC (Italy); Severo Ochoa and María de Maeztu Units of Excellence, GVA, XuntaGal, GENCAT, InTalent-Inditex and Prog. Atracción Talento CM (Spain); SRC (Sweden); the Leverhulme Trust, the Royal Society and UKRI (United Kingdom).

References

- [1] CDF collaboration, T. Aaltonen *et al.*, *Evidence for a narrow near-threshold structure in the $J/\psi\phi$ mass spectrum in $B^+ \rightarrow J/\psi\phi K^+$ decays*, Phys. Rev. Lett. **102** (2009) 242002, arXiv:0903.2229.
- [2] CMS collaboration, S. Chatrchyan *et al.*, *Observation of a peaking structure in the $J/\psi\phi$ mass spectrum from $B^\pm \rightarrow J/\psi\phi K^\pm$ decays*, Phys. Lett. **B734** (2014) 261, arXiv:1309.6920.
- [3] D0 collaboration, V. M. Abazov *et al.*, *Search for the $X(4140)$ state in $B^+ \rightarrow J/\psi\phi K^+$ decays with the D0 Detector*, Phys. Rev. **D89** (2014) 012004, arXiv:1309.6580.
- [4] LHCb collaboration, R. Aaij *et al.*, *Amplitude analysis of $B^+ \rightarrow J/\psi\phi K^+$ decays*, Phys. Rev. **D95** (2017) 012002, arXiv:1606.07898.
- [5] LHCb collaboration, R. Aaij *et al.*, *Observation of new resonances decaying to $J/\psi K^+$ and $J/\psi\phi$* , Phys. Rev. Lett. **127** (2021) 082001, arXiv:2103.01803.
- [6] Particle Data Group, S. Navas *et al.*, *Review of particle physics*, Phys. Rev. **D110** (2024) 030001.
- [7] CMS collaboration, V. Khachatryan *et al.*, *Observation of the decay $B^+ \rightarrow \psi(2S)\phi(1020)K^+$ in pp collisions at $\sqrt{s} = 8$ TeV*, Phys. Lett. **B764** (2017) 66, arXiv:1607.02638.
- [8] J. M. Dias, X. Liu, and M. Nielsen, *Prediction for the decay width of a charged state near the $D_s\bar{D}^*/D_s^*\bar{D}$ threshold*, Phys. Rev. **D88** (2013) 096014, arXiv:1307.7100.
- [9] M. B. Voloshin, *Strange hadrocharmonium*, Phys. Lett. **B798** (2019) 135022, arXiv:1901.01936.
- [10] LHCb collaboration, A. A. Alves Jr. *et al.*, *The LHCb detector at the LHC*, JINST **3** (2008) S08005.
- [11] LHCb collaboration, R. Aaij *et al.*, *LHCb detector performance*, Int. J. Mod. Phys. **A30** (2015) 1530022, arXiv:1412.6352.
- [12] R. Aaij *et al.*, *Performance of the LHCb Vertex Locator*, JINST **9** (2014) P09007, arXiv:1405.7808.
- [13] R. Arink *et al.*, *Performance of the LHCb Outer Tracker*, JINST **9** (2014) P01002, arXiv:1311.3893.
- [14] P. d'Argent *et al.*, *Improved performance of the LHCb Outer Tracker in LHC Run 2*, JINST **12** (2017) P11016, arXiv:1708.00819.
- [15] M. Adinolfi *et al.*, *Performance of the LHCb RICH detector at the LHC*, Eur. Phys. J. **C73** (2013) 2431, arXiv:1211.6759.
- [16] A. A. Alves Jr. *et al.*, *Performance of the LHCb muon system*, JINST **8** (2013) P02022, arXiv:1211.1346.

- [17] R. Aaij *et al.*, *The LHCb trigger and its performance in 2011*, JINST **8** (2013) P04022, [arXiv:1211.3055](#).
- [18] T. Sjöstrand, S. Mrenna, and P. Skands, *A brief introduction to PYTHIA 8.1*, Comput. Phys. Commun. **178** (2008) 852, [arXiv:0710.3820](#); T. Sjöstrand, S. Mrenna, and P. Skands, *PYTHIA 6.4 physics and manual*, JHEP **05** (2006) 026, [arXiv:hep-ph/0603175](#).
- [19] I. Belyaev *et al.*, *Handling of the generation of primary events in Gauss, the LHCb simulation framework*, J. Phys. Conf. Ser. **331** (2011) 032047.
- [20] D. J. Lange, *The EvtGen particle decay simulation package*, Nucl. Instrum. Meth. **A462** (2001) 152.
- [21] P. Golonka and Z. Was, *PHOTOS Monte Carlo: A precision tool for QED corrections in Z and W decays*, Eur. Phys. J. **C45** (2006) 97, [arXiv:hep-ph/0506026](#).
- [22] Geant4 collaboration, J. Allison *et al.*, *Geant4 developments and applications*, IEEE Trans. Nucl. Sci. **53** (2006) 270; Geant4 collaboration, S. Agostinelli *et al.*, *Geant4: A simulation toolkit*, Nucl. Instrum. Meth. **A506** (2003) 250.
- [23] M. Clemencic *et al.*, *The LHCb simulation application, Gauss: Design, evolution and experience*, J. Phys. Conf. Ser. **331** (2011) 032023.
- [24] L. Breiman, J. H. Friedman, R. A. Olshen, and C. J. Stone, *Classification and regression trees*, Wadsworth international group, Belmont, California, USA, 1984.
- [25] Y. Freund and R. E. Schapire, *A decision-theoretic generalization of on-line learning and an application to boosting*, J. Comput. Syst. Sci. **55** (1997) 119.
- [26] H. Voss, A. Hoecker, J. Stelzer, and F. Tegenfeldt, *TMVA - Toolkit for Multivariate Data Analysis with ROOT*, PoS **ACAT** (2007) 040; A. Hoecker *et al.*, *TMVA 4 — Toolkit for Multivariate Data Analysis with ROOT. Users Guide.*, [arXiv:physics/0703039](#).
- [27] T. Skwarnicki, *A study of the radiative cascade transitions between the Upsilon-prime and Upsilon resonances*, PhD thesis, Institute of Nuclear Physics, Krakow, 1986, DESY-F31-86-02.
- [28] M. Pivk and F. R. Le Diberder, *sPlot: A statistical tool to unfold data distributions*, Nucl. Instrum. Meth. **A555** (2005) 356, [arXiv:physics/0402083](#).
- [29] ARGUS collaboration, H. Albrecht *et al.*, *Search for hadronic $b \rightarrow u$ decays*, Phys. Lett. **B241** (1990) 278.
- [30] R. Aaij *et al.*, *Selection and processing of calibration samples to measure the particle identification performance of the LHCb experiment in Run 2*, EPJ Tech. Instrum. **6** (2019) 1, [arXiv:1803.00824](#).
- [31] A. Rogozhnikov, *Reweighting with boosted decision trees*, Journal of Physics: Conference Series **762** (2016) 012036.

- [32] N. L. Johnson, *Systems of frequency curves generated by methods of translation*, *Biometrika* **36** (1949) 149.
- [33] R. Nisius, *BLUE: combining correlated estimates of physics observables within ROOT using the Best Linear Unbiased Estimate method*, *SoftwareX* **11** (2020) 100468, [arXiv:2001.10310](https://arxiv.org/abs/2001.10310).

LHCb collaboration

R. Aaij³⁸, A.S.W. Abdelmotteleb⁵⁷, C. Abellan Beteta⁵¹, F. Abudinén⁵⁷, T. Ackernley⁶¹, A. A. Adefisoye⁶⁹, B. Adeva⁴⁷, M. Adinolfi⁵⁵, P. Adlarson⁸², C. Agapopoulou¹⁴, C.A. Aidala⁸³, Z. Ajaltouni¹¹, S. Akar⁶⁶, K. Akiba³⁸, P. Albicocco²⁸, J. Albrecht^{19,f}, F. Alessio⁴⁹, M. Alexander⁶⁰, Z. Aliouche⁶³, P. Alvarez Cartelle⁵⁶, R. Amalric¹⁶, S. Amato³, J.L. Amey⁵⁵, Y. Amhis¹⁴, L. An⁶, L. Anderlini²⁷, M. Andersson⁵¹, A. Andreianov⁴⁴, P. Andreola⁵¹, M. Andreotti²⁶, D. Andreou⁶⁹, A. Anelli^{31,o}, D. Ao⁷, F. Archilli^{37,u}, M. Argenton²⁶, S. Arguedas Cuendis^{9,49}, A. Artamonov⁴⁴, M. Artuso⁶⁹, E. Aslanides¹³, R. Ataíde Da Silva⁵⁰, M. Atzeni⁶⁵, B. Audurier¹², D. Bacher⁶⁴, I. Bachiller Perea¹⁰, S. Bachmann²², M. Bachmayer⁵⁰, J.J. Back⁵⁷, P. Baladron Rodriguez⁴⁷, V. Balagura¹⁵, A. Balboni²⁶, W. Baldini²⁶, L. Balzani¹⁹, H. Bao⁷, J. Baptista de Souza Leite⁶¹, C. Barbero Pretel^{47,12}, M. Barbetti²⁷, I. R. Barbosa⁷⁰, R.J. Barlow⁶³, M. Barnyakov²⁵, S. Barsuk¹⁴, W. Barter⁵⁹, M. Bartolini⁵⁶, J. Bartz⁶⁹, J.M. Basels¹⁷, S. Bashir⁴⁰, G. Bassi^{35,r}, B. Batsukh⁵, P. B. Battista¹⁴, A. Bay⁵⁰, A. Beck⁵⁷, M. Becker¹⁹, F. Bedeschi³⁵, I.B. Bediaga², N. A. Behling¹⁹, S. Belin⁴⁷, K. Belous⁴⁴, I. Belov²⁹, I. Belyaev³⁶, G. Benane¹³, G. Bencivenni²⁸, E. Ben-Haim¹⁶, A. Berezhnoy⁴⁴, R. Bernet⁵¹, S. Bernet Andres⁴⁶, A. Bertolin³³, C. Betancourt⁵¹, F. Betti⁵⁹, J. Bex⁵⁶, I.a. Bezshyiko⁵¹, J. Bhom⁴¹, M.S. Bieker¹⁹, N.V. Biesuz²⁶, P. Billoir¹⁶, A. Biolchini³⁸, M. Birch⁶², F.C.R. Bishop¹⁰, A. Bitadze⁶³, A. Bizzeti¹⁶, T. Blake⁵⁷, F. Blanc⁵⁰, J.E. Blank¹⁹, S. Blusk⁶⁹, V. Bocharnikov⁴⁴, J.A. Boelhaeve¹⁹, O. Boente Garcia¹⁵, T. Boettcher⁶⁶, A. Bohare⁵⁹, A. Boldyrev⁴⁴, C.S. Bolognani⁷⁹, R. Bolzonella^{26,l}, R. B. Bonacci¹, N. Bondar⁴⁴, A. Bordeliu⁴⁹, F. Borgato^{33,p}, S. Borghi⁶³, M. Borsato^{31,o}, J.T. Borsuk⁴¹, S.A. Bouchiba⁵⁰, M. Bovill⁶⁴, T.J.V. Bowcock⁶¹, A. Boyer⁴⁹, C. Bozzi²⁶, A. Brea Rodriguez⁵⁰, N. Breer¹⁹, J. Brodzicka⁴¹, A. Brossa Gonzalo^{47,†}, J. Brown⁶¹, D. Brundu³², E. Buchanan⁵⁹, A. Buonaura⁵¹, L. Buonincontri^{33,p}, A.T. Burke⁶³, C. Burr⁴⁹, J.S. Butter⁵⁶, J. Buytaert⁴⁹, W. Byczynski⁴⁹, S. Cadeddu³², H. Cai⁷⁴, A. Caillet¹⁶, R. Calabrese^{26,l}, S. Calderon Ramirez⁹, L. Calefice⁴⁵, S. Cali²⁸, M. Calvi^{31,o}, M. Calvo Gomez⁴⁶, P. Camargo Magalhaes^{2,y}, J. I. Cambon Bouzas⁴⁷, P. Campana²⁸, D.H. Campora Perez⁷⁹, A.F. Campoverde Quezada⁷, S. Capelli³¹, L. Capriotti²⁶, R. Caravaca-Mora⁹, A. Carbone^{25,j}, L. Carcedo Salgado⁴⁷, R. Cardinale^{29,m}, A. Cardini³², P. Carniti^{31,o}, L. Carus²², A. Casais Vidal⁶⁵, R. Caspary²², G. Casse⁶¹, M. Cattaneo⁴⁹, G. Cavallero^{26,49}, V. Cavallini^{26,l}, S. Celani²², D. Cervenkov⁶⁴, S. Cesare^{30,n}, A.J. Chadwick⁶¹, I. Chahrour⁸³, M. Charles¹⁶, Ph. Charpentier⁴⁹, E. Chatzianagnostou³⁸, M. Chefdeville¹⁰, C. Chen¹³, S. Chen⁵, Z. Chen⁷, A. Chernov⁴¹, S. Chernyshenko⁵³, X. Chiotopoulos⁷⁹, V. Chobanova⁸¹, S. Cholak⁵⁰, M. Chrzascz⁴¹, A. Chubykin⁴⁴, V. Chulikov²⁸, P. Ciambriano²⁸, X. Cid Vidal⁴⁷, G. Ciezarek⁴⁹, P. Cifra⁴⁹, P.E.L. Clarke⁵⁹, M. Clemencic⁴⁹, H.V. Cliff⁵⁶, J. Closier⁴⁹, C. Cocha Toapaxi²², V. Coco⁴⁹, J. Cogan¹³, E. Cogneras¹¹, L. Cojocariu⁴³, S. Collaviti⁵⁰, P. Collins⁴⁹, T. Colombo⁴⁹, M. Colonna¹⁹, A. Comerma-Montells⁴⁵, L. Congedo²⁴, A. Contu³², N. Cooke⁶⁰, I. Corredoira⁴⁷, A. Correia¹⁶, G. Corti⁴⁹, J. Cottee Meldrum⁵⁵, B. Couturier⁴⁹, D.C. Craik⁵¹, M. Cruz Torres^{2,g}, E. Curras Rivera⁵⁰, R. Currie⁵⁹, C.L. Da Silva⁶⁸, S. Dadabaev⁴⁴, L. Dai⁷¹, X. Dai⁶, E. Dall’Occo⁴⁹, J. Dalseno⁴⁷, C. D’Ambrosio⁴⁹, J. Daniel¹¹, A. Danilina⁴⁴, P. d’Argent²⁴, G. Darze³, A. Davidson⁵⁷, J.E. Davies⁶³, A. Davis⁶³, O. De Aguiar Francisco⁶³, C. De Angelis^{32,k}, F. De Benedetti⁴⁹, J. de Boer³⁸, K. De Bruyn⁷⁸, S. De Capua⁶³, M. De Cian²²,

U. De Freitas Carneiro Da Graca^{2,a} , E. De Lucia²⁸ , J.M. De Miranda² , L. De Paula³ ,
M. De Serio^{24,h} , P. De Simone²⁸ , F. De Vellis¹⁹ , J.A. de Vries⁷⁹ , F. Debernardis²⁴ ,
D. Decamp¹⁰ , V. Dedu¹³ , S. Dekkers¹ , L. Del Buono¹⁶ , B. Delaney⁶⁵ ,
H.-P. Dembinski¹⁹ , J. Deng⁸ , V. Denysenko⁵¹ , O. Deschamps¹¹ , F. Dettori^{32,k} ,
B. Dey⁷⁷ , P. Di Nezza²⁸ , I. Diachkov⁴⁴ , S. Didenko⁴⁴ , S. Ding⁶⁹ , L. Dittmann²² ,
V. Dobishuk⁵³ , A. D. Docheva⁶⁰ , C. Dong^{4,b} , A.M. Donohoe²³ , F. Dordei³² ,
A.C. dos Reis² , A. D. Dowling⁶⁹ , W. Duan⁷² , P. Duda⁸⁰ , M.W. Dudek⁴¹ ,
L. Dufour⁴⁹ , V. Duk³⁴ , P. Durante⁴⁹ , M. M. Duras⁸⁰ , J.M. Durham⁶⁸ , O. D.
Durmus⁷⁷ , A. Dziurda⁴¹ , A. Dzyuba⁴⁴ , S. Easo⁵⁸ , E. Eckstein¹⁸ , U. Egede¹ ,
A. Egorychev⁴⁴ , V. Egorychev⁴⁴ , S. Eisenhardt⁵⁹ , E. Ejopu⁶³ , L. Eklund⁸² ,
M. Elashri⁶⁶ , J. Ellbracht¹⁹ , S. Ely⁶² , A. Ene⁴³ , J. Eschle⁶⁹ , S. Esen²² ,
T. Evans⁶³ , F. Fabiano^{32,k} , L.N. Falcao² , Y. Fan⁷ , B. Fang⁷ , L. Fantini^{34,q,49} ,
M. Faria⁵⁰ , K. Farmer⁵⁹ , D. Fazzini^{31,o} , L. Felkowski⁸⁰ , M. Feng^{5,7} , M. Feo¹⁹ ,
A. Fernandez Casani⁴⁸ , M. Fernandez Gomez⁴⁷ , A.D. Fernez⁶⁷ , F. Ferrari^{25,j} ,
F. Ferreira Rodrigues³ , M. Ferrillo⁵¹ , M. Ferro-Luzzi⁴⁹ , S. Filippov⁴⁴ , R.A. Fini²⁴ ,
M. Fiorini^{26,l} , M. Firlej⁴⁰ , K.L. Fischer⁶⁴ , D.S. Fitzgerald⁸³ , C. Fitzpatrick⁶³ ,
T. Fiutowski⁴⁰ , F. Fleuret¹⁵ , M. Fontana²⁵ , L. F. Foreman⁶³ , R. Forty⁴⁹ ,
D. Foulds-Holt⁵⁶ , V. Franco Lima³ , M. Franco Sevilla⁶⁷ , M. Frank⁴⁹ ,
E. Franzoso^{26,l} , G. Frau⁶³ , C. Frei⁴⁹ , D.A. Friday⁶³ , J. Fu⁷ , Q. Führung^{19,f,56} ,
Y. Fujii¹ , T. Fulghesu¹⁶ , E. Gabriel³⁸ , G. Galati²⁴ , M.D. Galati³⁸ ,
A. Gallas Torreira⁴⁷ , D. Galli^{25,j} , S. Gambetta⁵⁹ , M. Gandelman³ , P. Gandini³⁰ , B.
Ganie⁶³ , H. Gao⁷ , R. Gao⁶⁴ , T.Q. Gao⁵⁶ , Y. Gao⁸ , Y. Gao⁶ , Y. Gao⁸ ,
L.M. Garcia Martin⁵⁰ , P. Garcia Moreno⁴⁵ , J. García Pardiñas⁴⁹ , P. Gardner⁶⁷ , K. G.
Garg⁸ , L. Garrido⁴⁵ , C. Gaspar⁴⁹ , R.E. Geertsema³⁸ , L.L. Gerken¹⁹ ,
E. Gersabeck⁶³ , M. Gersabeck²⁰ , T. Gershon⁵⁷ , S. Ghizzo^{29,m} ,
Z. Ghorbanimoghaddam⁵⁵ , L. Giambastiani^{33,p} , F. I. Giasemis^{16,e} , V. Gibson⁵⁶ ,
H.K. Giemza⁴² , A.L. Gilman⁶⁴ , M. Giovannetti²⁸ , A. Gioventù⁴⁵ , L. Girardey^{63,58} ,
P. Gironella Gironell⁴⁵ , C. Giugliano^{26,l} , M.A. Giza⁴¹ , E.L. Gkoukousis⁶² ,
F.C. Glaser^{14,22} , V.V. Gligorov^{16,49} , C. Göbel⁷⁰ , E. Golobardes⁴⁶ , D. Golubkov⁴⁴ ,
A. Golutvin^{62,49,44} , S. Gomez Fernandez⁴⁵ , W. Gomulka⁴⁰ , F. Goncalves Abrantes⁶⁴ ,
M. Goncerz⁴¹ , G. Gong^{4,b} , J. A. Gooding¹⁹ , I.V. Gorelov⁴⁴ , C. Gotti³¹ ,
J.P. Grabowski¹⁸ , L.A. Granado Cardoso⁴⁹ , E. Graugés⁴⁵ , E. Graverini^{50,s} ,
L. Grazette⁵⁷ , G. Graziani , A. T. Grecu⁴³ , L.M. Greeven³⁸ , N.A. Grieser⁶⁶ ,
L. Grillo⁶⁰ , S. Gromov⁴⁴ , C. Gu¹⁵ , M. Guarise²⁶ , L. Guerry¹¹ , M. Guittiere¹⁴ ,
V. Guliaeva⁴⁴ , P. A. Günther²² , A.-K. Guseinov⁵⁰ , E. Gushchin⁴⁴ , Y. Guz^{6,49,44} ,
T. Gys⁴⁹ , K. Habermann¹⁸ , T. Hadavizadeh¹ , C. Hadjivasiliou⁶⁷ , G. Haefeli⁵⁰ ,
C. Haen⁴⁹ , M. Hajheidari⁴⁹ , G. Hallett⁵⁷ , M.M. Halvorsen⁴⁹ , P.M. Hamilton⁶⁷ ,
J. Hammerich⁶¹ , Q. Han⁸ , X. Han^{22,49} , S. Hansmann-Menzemer²² , L. Hao⁷ ,
N. Harnew⁶⁴ , T. H. Harris¹ , M. Hartmann¹⁴ , S. Hashmi⁴⁰ , J. He^{7,c} ,
F. Hemmer⁴⁹ , C. Henderson⁶⁶ , R.D.L. Henderson^{1,57} , A.M. Hennequin⁴⁹ ,
K. Hennessy⁶¹ , L. Henry⁵⁰ , J. Herd⁶² , P. Herrero Gascon²² , J. Heuel¹⁷ ,
A. Hicheur³ , G. Hijano Mendizabal⁵¹ , J. Horswill⁶³ , R. Hou⁸ , Y. Hou¹¹ ,
N. Howarth⁶¹ , J. Hu⁷² , W. Hu⁶ , X. Hu^{4,b} , W. Huang⁷ , W. Hulsbergen³⁸ ,
R.J. Hunter⁵⁷ , M. Hushchyn⁴⁴ , D. Hutchcroft⁶¹ , M. Idzik⁴⁰ , D. Ilin⁴⁴ , P. Ilten⁶⁶ ,
A. Inglessi⁴⁴ , A. Iniuukhin⁴⁴ , A. Ishteev⁴⁴ , K. Ivshin⁴⁴ , R. Jacobsson⁴⁹ , H. Jage¹⁷ ,
S.J. Jaimes Elles^{75,49,48} , S. Jakobsen⁴⁹ , E. Jans³⁸ , B.K. Jashal⁴⁸ , A. Jawahery^{67,49} ,
V. Jevtic^{19,f} , E. Jiang⁶⁷ , X. Jiang^{5,7} , Y. Jiang⁷ , Y. J. Jiang⁶ , M. John⁶⁴ , A.
John Rubesh Rajan²³ , D. Johnson⁵⁴ , C.R. Jones⁵⁶ , T.P. Jones⁵⁷ , S. Joshi⁴² ,
B. Jost⁴⁹ , J. Juan Castilla⁵⁶ , N. Jurik⁴⁹ , I. Juszczak⁴¹ , D. Kaminaris⁵⁰ ,
S. Kandybei⁵² , M. Kane⁵⁹ , Y. Kang^{4,b} , C. Kar¹¹ , M. Karacson⁴⁹ ,

D. Karpenkov⁴⁴ , A. Kauniskangas⁵⁰ , J.W. Kautz⁶⁶ , M.K. Kazanecki⁴¹ , F. Keizer⁴⁹ ,
M. Kenzie⁵⁶ , T. Ketel³⁸ , B. Khanji⁶⁹ , A. Kharisova⁴⁴ , S. Kholodenko^{35,49} ,
G. Khreich¹⁴ , T. Kirn¹⁷ , V.S. Kirsebom^{31,o} , O. Kitouni⁶⁵ , S. Klaver³⁹ ,
N. Kleijne^{35,r} , K. Klimaszewski⁴² , M.R. Kmiec⁴² , S. Koliiev⁵³ , L. Kolk¹⁹ ,
A. Konoplyannikov⁴⁴ , P. Kopciwicz^{40,49} , P. Koppenburg³⁸ , M. Korolev⁴⁴ ,
I. Kostiuk³⁸ , O. Kot⁵³ , S. Kotriakhova , A. Kozachuk⁴⁴ , P. Kravchenko⁴⁴ ,
L. Kravchuk⁴⁴ , M. Kreps⁵⁷ , P. Krokovny⁴⁴ , W. Krupa⁶⁹ , W. Krzemien⁴² ,
O. Kshyvanskyi⁵³ , S. Kubis⁸⁰ , M. Kucharczyk⁴¹ , V. Kudryavtsev⁴⁴ , E. Kulikova⁴⁴ ,
A. Kupsc⁸² , B. K. Kutsenko¹³ , D. Lacarrere⁴⁹ , P. Laguarda Gonzalez⁴⁵ , A. Lai³² ,
A. Lampis³² , D. Lancierini⁵⁶ , C. Landesa Gomez⁴⁷ , J.J. Lane¹ , R. Lane⁵⁵ ,
G. Lanfranchi²⁸ , C. Langenbruch²² , J. Langer¹⁹ , O. Lantwin⁴⁴ , T. Latham⁵⁷ ,
F. Lazzari^{35,s} , C. Lazzeroni⁵⁴ , R. Le Gac¹³ , H. Lee⁶¹ , R. Lefèvre¹¹ , A. Leflat⁴⁴ ,
S. Legotin⁴⁴ , M. Lehurax⁵⁷ , E. Lemos Cid⁴⁹ , O. Leroy¹³ , T. Lesiak⁴¹ , E.
D. Lesser⁴⁹ , B. Leverington²² , A. Li^{4,b} , C. Li¹³ , H. Li⁷² , K. Li⁸ , L. Li⁶³ ,
M. Li⁸ , P. Li⁷ , P.-R. Li⁷³ , Q. Li^{5,7} , S. Li⁸ , T. Li^{5,d} , T. Li⁷² , Y. Li⁸ ,
Y. Li⁵ , Z. Lian^{4,b} , X. Liang⁶⁹ , S. Libralon⁴⁸ , C. Lin⁷ , T. Lin⁵⁸ , R. Lindner⁴⁹ ,
H. Linton⁶² , V. Lisovskyi⁵⁰ , R. Litvinov^{32,49} , F. L. Liu¹ , G. Liu⁷² , K. Liu⁷³ ,
S. Liu^{5,7} , W. Liu⁸ , Y. Liu⁵⁹ , Y. Liu⁷³ , Y. L. Liu⁶² , A. Lobo Salvia⁴⁵ ,
A. Loi³² , T. Long⁵⁶ , J.H. Lopes³ , A. Lopez Huertas⁴⁵ , S. López Soliño⁴⁷ ,
Q. Lu¹⁵ , C. Lucarelli²⁷ , D. Lucchesi^{33,p} , M. Lucio Martinez⁷⁹ , V. Lukashenko^{38,53} ,
Y. Luo⁶ , A. Lupato^{33,i} , E. Luppi^{26,l} , K. Lynch²³ , X.-R. Lyu⁷ , G. M. Ma^{4,b} ,
S. Maccolini¹⁹ , F. Machefert¹⁴ , F. Maciuc⁴³ , B. Mack⁶⁹ , I. Mackay⁶⁴ , L. M.
Mackey⁶⁹ , L.R. Madhan Mohan⁵⁶ , M. J. Madurai⁵⁴ , A. Maevskiy⁴⁴ ,
D. Magdalinski³⁸ , D. Maisuzenko⁴⁴ , M.W. Majewski⁴⁰ , J.J. Malczewski⁴¹ , S. Malde⁶⁴ ,
L. Malentacca⁴⁹ , A. Malinin⁴⁴ , T. Maltsev⁴⁴ , G. Manca^{32,k} , G. Mancinelli¹³ ,
C. Mancuso^{30,14,n} , R. Manera Escalero⁴⁵ , F. M. Manganella³⁷ , D. Manuzzi²⁵ ,
D. Marangotto^{30,n} , J.F. Marchand¹⁰ , R. Marchevski⁵⁰ , U. Marconi²⁵ , E. Mariani¹⁶ ,
S. Mariani⁴⁹ , C. Marin Benito^{45,49} , J. Marks²² , A.M. Marshall⁵⁵ , L. Martel⁶⁴ ,
G. Martelli^{34,q} , G. Martellotti³⁶ , L. Martinazzoli⁴⁹ , M. Martinelli^{31,o} , D.
Martinez Gomez⁷⁸ , D. Martinez Santos⁸¹ , F. Martinez Vidal⁴⁸ , A.
Martorell i Granollers⁴⁶ , A. Massafferri² , R. Matev⁴⁹ , A. Mathad⁴⁹ , V. Matiuinin⁴⁴ ,
C. Matteuzzi⁶⁹ , K.R. Mattioli¹⁵ , A. Mauri⁶² , E. Maurice¹⁵ , J. Mauricio⁴⁵ ,
P. Mayencourt⁵⁰ , J. Mazorra de Cos⁴⁸ , M. Mazurek⁴² , M. McCann⁶² ,
L. McConnell²³ , T.H. McGrath⁶³ , N.T. McHugh⁶⁰ , A. McNab⁶³ , R. McNulty²³ ,
B. Meadows⁶⁶ , G. Meier¹⁹ , D. Melnychuk⁴² , F. M. Meng^{4,b} , M. Merk^{38,79} ,
A. Merli⁵⁰ , L. Meyer Garcia⁶⁷ , D. Miao^{5,7} , H. Miao⁷ , M. Mikhasenko⁷⁶ ,
D.A. Milanés⁷⁵ , A. Minotti^{31,o} , E. Minucci²⁸ , T. Miralles¹¹ , B. Mitreska¹⁹ ,
D.S. Mitzel¹⁹ , A. Modak⁵⁸ , R.A. Mohammed⁶⁴ , R.D. Moise¹⁷ , S. Mokhenko⁴⁴ , E.
F. Molina Cardenas⁸³ , T. Mombächer⁴⁹ , M. Monk^{57,1} , S. Monteil¹¹ ,
A. Morcillo Gomez⁴⁷ , G. Morello²⁸ , M.J. Morello^{35,r} , M.P. Morgenthaler²² ,
J. Moron⁴⁰ , W. Morren³⁸ , A.B. Morris⁴⁹ , A.G. Morris¹³ , R. Mountain⁶⁹ ,
H. Mu^{4,b} , Z. M. Mu⁶ , E. Muhammad⁵⁷ , F. Muheim⁵⁹ , M. Mulder⁷⁸ , K. Müller⁵¹ ,
F. Muñoz-Rojas⁹ , R. Murta⁶² , P. Naik⁶¹ , T. Nakada⁵⁰ , R. Nandakumar⁵⁸ ,
T. Nanut⁴⁹ , I. Nasteva³ , M. Needham⁵⁹ , N. Neri^{30,n} , S. Neubert¹⁸ , N. Neufeld⁴⁹ ,
P. Neustroev⁴⁴ , J. Nicolini^{19,14} , D. Nicotra⁷⁹ , E.M. Niel⁴⁹ , N. Nikitin⁴⁴ , Q. Niu⁷³,
P. Nogaroli³ , P. Nogga¹⁸ , C. Normand⁵⁵ , J. Novoa Fernandez⁴⁷ , G. Nowak⁶⁶ ,
C. Nunez⁸³ , H. N. Nur⁶⁰ , A. Oblakowska-Mucha⁴⁰ , V. Obraztsov⁴⁴ , T. Oeser¹⁷ ,
S. Okamura^{26,l} , A. Okhotnikov⁴⁴ , O. Okhrimenko⁵³ , R. Oldeman^{32,k} , F. Oliva⁵⁹ ,
M. Olocco¹⁹ , C.J.G. Onderwater⁷⁹ , R.H. O'Neil⁵⁹ , D. Osthues¹⁹ ,
J.M. Otalora Goicochea³ , P. Owen⁵¹ , A. Oyanguren⁴⁸ , O. Ozcelik⁵⁹ , F. Paciolla^{35,v}

A. Padee⁴², K.O. Padeken¹⁸, B. Pagare⁵⁷, P.R. Pais²², T. Pajero⁴⁹, A. Palano²⁴,
 M. Palutan²⁸, X. Pan^{4,b}, G. Panshin⁴⁴, L. Paolucci⁵⁷, A. Papanestis^{58,49},
 M. Pappagallo^{24,h}, L.L. Pappalardo^{26,l}, C. Pappenheimer⁶⁶, C. Parkes⁶³, D.
 Parmar⁷⁶, B. Passalacqua^{26,l}, G. Passaleva²⁷, D. Passaro^{35,r}, A. Pastore²⁴,
 M. Patel⁶², J. Patoc⁶⁴, C. Patrignani^{25,j}, A. Paul⁶⁹, C.J. Pawley⁷⁹,
 A. Pellegrino³⁸, J. Peng^{5,7}, M. Pepe Altarelli²⁸, S. Perazzini²⁵, D. Pereima⁴⁴, H.
 Pereira Da Costa⁶⁸, A. Pereiro Castro⁴⁷, P. Perret¹¹, A. Perrevoort⁷⁸,
 A. Perro^{49,13}, M.J. Peters⁶⁶, K. Petridis⁵⁵, A. Petrolini^{29,m}, J. P. Pfaller⁶⁶,
 H. Pham⁶⁹, L. Pica^{35,r}, M. Piccini³⁴, L. Piccolo³², B. Pietrzyk¹⁰, G. Pietrzyk¹⁴,
 D. Pinci³⁶, F. Pisani⁴⁹, M. Pizzichemi^{31,o,49}, V. Placinta⁴³, M. Plo Casasus⁴⁷,
 T. Poeschl⁴⁹, F. Polci^{16,49}, M. Poli Lener²⁸, A. Poluektov¹³, N. Polukhina⁴⁴,
 I. Polyakov⁴⁴, E. Polycarpo³, S. Ponce⁴⁹, D. Popov⁷, S. Poslavskii⁴⁴,
 K. Prasanth⁵⁹, C. Prouve⁸¹, D. Provenzano^{32,k}, V. Pugatch⁵³, G. Punzi^{35,s}, S.
 Qasim⁵¹, Q. Q. Qian⁶, W. Qian⁷, N. Qin^{4,b}, S. Qu^{4,b}, R. Quagliani⁴⁹,
 R.I. Rabadan Trejo⁵⁷, J.H. Rademacker⁵⁵, M. Rama³⁵, M. Ramírez García⁸³,
 V. Ramos De Oliveira⁷⁰, M. Ramos Pernas⁵⁷, M.S. Rangel³, F. Ratnikov⁴⁴,
 G. Raven³⁹, M. Rebollo De Miguel⁴⁸, F. Redi^{30,i}, J. Reich⁵⁵, F. Reiss⁶³, Z. Ren⁷,
 P.K. Resmi⁶⁴, R. Ribatti⁵⁰, G. Ricart^{15,12}, D. Ricciardi^{35,r}, S. Ricciardi⁵⁸,
 K. Richardson⁶⁵, M. Richardson-Slipper⁵⁹, K. Rinnert⁶¹, P. Robbe^{14,49},
 G. Robertson⁶⁰, E. Rodrigues⁶¹, A. Rodriguez Alvarez⁴⁵, E. Rodriguez Fernandez⁴⁷,
 J.A. Rodriguez Lopez⁷⁵, E. Rodriguez Rodriguez⁴⁷, J. Roensch¹⁹, A. Rogachev⁴⁴,
 A. Rogovskiy⁵⁸, D.L. Rolf⁴⁹, P. Roloff⁴⁹, V. Romanovskiy⁶⁶, A. Romero Vidal⁴⁷,
 G. Romolini²⁶, F. Ronchetti⁵⁰, T. Rong⁶, M. Rotondo²⁸, S. R. Roy²²,
 M.S. Rudolph⁶⁹, M. Ruiz Diaz²², R.A. Ruiz Fernandez⁴⁷, J. Ruiz Vidal^{82,z},
 A. Ryzhikov⁴⁴, J. Ryzka⁴⁰, J. J. Saavedra-Arias⁹, J.J. Saborido Silva⁴⁷, R. Sadek¹⁵,
 N. Sagidova⁴⁴, D. Sahoo⁷⁷, N. Sahoo⁵⁴, B. Saïtta^{32,k}, M. Salomoni^{31,49,o},
 I. Sanderswood⁴⁸, R. Santacesaria³⁶, C. Santamarina Rios⁴⁷, M. Santimaria^{28,49},
 L. Santoro², E. Santovetti³⁷, A. Saputi^{26,49}, D. Saranin⁴⁴, A. Sarnatskiy⁷⁸,
 G. Sarpis⁵⁹, M. Sarpis⁶³, C. Satriano^{36,t}, A. Satta³⁷, M. Saur⁶, D. Savrina⁴⁴,
 H. Sazak¹⁷, F. Sborzacchi^{49,28}, L.G. Scantlebury Smead⁶⁴, A. Scarabotto¹⁹,
 S. Schael¹⁷, S. Scherl⁶¹, M. Schiller⁶⁰, H. Schindler⁴⁹, M. Schmelling²¹,
 B. Schmidt⁴⁹, S. Schmitt¹⁷, H. Schmitz¹⁸, O. Schneider⁵⁰, A. Schopper⁴⁹,
 N. Schulte¹⁹, S. Schulte⁵⁰, M.H. Schune¹⁴, R. Schwemmer⁴⁹, G. Schwering¹⁷,
 B. Sciascia²⁸, A. Sciuccati⁴⁹, I. Segal⁷⁶, S. Sellam⁴⁷, A. Semennikov⁴⁴,
 T. Senger⁵¹, M. Senghi Soares³⁹, A. Sergi^{29,m}, N. Serra⁵¹, L. Sestini³³,
 A. Seuthe¹⁹, Y. Shang⁶, D.M. Shangase⁸³, M. Shapkin⁴⁴, R. S. Sharma⁶⁹,
 I. Shchemerov⁴⁴, L. Shchutska⁵⁰, T. Shears⁶¹, L. Shekhtman⁴⁴, Z. Shen⁶,
 S. Sheng^{5,7}, V. Shevchenko⁴⁴, B. Shi⁷, Q. Shi⁷, Y. Shimizu¹⁴, E. Shmanin²⁵,
 R. Shorkin⁴⁴, J.D. Shupperd⁶⁹, R. Silva Coutinho⁶⁹, G. Simi^{33,p}, S. Simone^{24,h},
 N. Skidmore⁵⁷, T. Skwarnicki⁶⁹, M.W. Slater⁵⁴, J.C. Smallwood⁶⁴, E. Smith⁶⁵,
 K. Smith⁶⁸, M. Smith⁶², A. Snoch³⁸, L. Soares Lavra⁵⁹, M.D. Sokoloff⁶⁶,
 F.J.P. Soler⁶⁰, A. Solomin^{44,55}, A. Solovev⁴⁴, I. Solovyev⁴⁴, N. S. Sommerfeld¹⁸,
 R. Song¹, Y. Song⁵⁰, Y. Song^{4,b}, Y. S. Song⁶, F.L. Souza De Almeida⁶⁹,
 B. Souza De Paula³, E. Spadaro Norella^{29,m}, E. Spedicato²⁵, J.G. Speer¹⁹,
 E. Spiridenkov⁴⁴, P. Spradlin⁶⁰, V. Srisakaran⁴⁹, F. Stagni⁴⁹, M. Stahl⁴⁹, S. Stahl⁴⁹,
 S. Stanislaus⁶⁴, E.N. Stein⁴⁹, O. Steinkamp⁵¹, O. Stenyakin⁴⁴, H. Stevens¹⁹,
 D. Strelalina⁴⁴, Y. Su⁷, F. Suljik⁶⁴, J. Sun³², L. Sun⁷⁴, D. Sundfeld²,
 W. Sutcliffe⁵¹, P.N. Swallow⁵⁴, K. Swientek⁴⁰, F. Swystun⁵⁶, A. Szabelski⁴²,
 T. Szumlak⁴⁰, Y. Tan^{4,b}, Y. Tang⁷⁴, M.D. Tat⁶⁴, A. Terentev⁴⁴,
 F. Terzuoli^{35,v,49}, F. Teubert⁴⁹, E. Thomas⁴⁹, D.J.D. Thompson⁵⁴, H. Tilquin⁶²,

V. Tisserand¹¹ , S. T’Jampens¹⁰ , M. Tobin^{5,49} , L. Tomassetti^{26,l} , G. Tonani^{30,n,49} , X. Tong⁶ , D. Torres Machado² , L. Toscano¹⁹ , D.Y. Tou^{4,b} , C. Tripl⁴⁶ , G. Tuci²² , N. Tuning³⁸ , L.H. Uecker²² , A. Ukleja⁴⁰ , D.J. Unverzagt²² , B. Urbach⁵⁹ , E. Ursov⁴⁴ , A. Usachov³⁹ , A. Ustyuzhanin⁴⁴ , U. Uwer²² , V. Vagnoni²⁵ , V. Valcarce Cadenas⁴⁷ , G. Valenti²⁵ , N. Valls Canudas⁴⁹ , H. Van Hecke⁶⁸ , E. van Herwijnen⁶² , C.B. Van Hulse^{47,x} , R. Van Laak⁵⁰ , M. van Veghel³⁸ , G. Vasquez⁵¹ , R. Vazquez Gomez⁴⁵ , P. Vazquez Regueiro⁴⁷ , C. Vázquez Sierra⁴⁷ , S. Vecchi²⁶ , J.J. Velthuis⁵⁵ , M. Veltri^{27,w} , A. Venkateswaran⁵⁰ , M. Verdoggia³² , M. Vesterinen⁵⁷ , D. Vico Benet⁶⁴ , P. Vidrier Villalba⁴⁵ , M. Vieites Diaz⁴⁹ , X. Vilasis-Cardona⁴⁶ , E. Vilella Figueras⁶¹ , A. Villa²⁵ , P. Vincent¹⁶ , F.C. Volle⁵⁴ , D. vom Bruch¹³ , N. Voropaev⁴⁴ , K. Vos⁷⁹ , C. Vrahas⁵⁹ , J. Wagner¹⁹ , J. Walsh³⁵ , E.J. Walton^{1,57} , G. Wan⁶ , C. Wang²² , G. Wang⁸ , H. Wang⁷³ , J. Wang⁶ , J. Wang⁵ , J. Wang^{4,b} , J. Wang⁷⁴ , M. Wang³⁰ , N. W. Wang⁷ , R. Wang⁵⁵ , X. Wang⁸ , X. Wang⁷² , X. W. Wang⁶² , Y. Wang⁶ , Y. W. Wang⁷³ , Z. Wang¹⁴ , Z. Wang^{4,b} , Z. Wang³⁰ , J.A. Ward^{57,1} , M. Waterlaet⁴⁹ , N.K. Watson⁵⁴ , D. Websdale⁶² , Y. Wei⁶ , J. Wendel⁸¹ , B.D.C. Westhenry⁵⁵ , C. White⁵⁶ , M. Whitehead⁶⁰ , E. Whiter⁵⁴ , A.R. Wiederhold⁶³ , D. Wiedner¹⁹ , G. Wilkinson⁶⁴ , M.K. Wilkinson⁶⁶ , M. Williams⁶⁵ , M. J. Williams⁴⁹ , M.R.J. Williams⁵⁹ , R. Williams⁵⁶ , Z. Williams⁵⁵ , F.F. Wilson⁵⁸ , M. Winn¹² , W. Wislicki⁴² , M. Witek⁴¹ , L. Witola²² , G. Wormser¹⁴ , S.A. Wotton⁵⁶ , H. Wu⁶⁹ , J. Wu⁸ , X. Wu⁷⁴ , Y. Wu⁶ , Z. Wu⁷ , K. Wyllie⁴⁹ , S. Xian⁷² , Z. Xiang⁵ , Y. Xie⁸ , A. Xu³⁵ , J. Xu⁷ , L. Xu^{4,b} , L. Xu^{4,b} , M. Xu⁵⁷ , Z. Xu⁴⁹ , Z. Xu⁷ , Z. Xu⁵ , K. Yang⁶² , S. Yang⁷ , X. Yang⁶ , Y. Yang^{29,m} , Z. Yang⁶ , V. Yeroshenko¹⁴ , H. Yeung⁶³ , H. Yin⁸ , X. Yin⁷ , C. Y. Yu⁶ , J. Yu⁷¹ , X. Yuan⁵ , Y. Yuan^{5,7} , E. Zaffaroni⁵⁰ , M. Zavertyaev²¹ , M. Zdybal⁴¹ , F. Zenesini^{25,j} , C. Zeng^{5,7} , M. Zeng^{4,b} , C. Zhang⁶ , D. Zhang⁸ , J. Zhang⁷ , L. Zhang^{4,b} , S. Zhang⁷¹ , S. Zhang⁶⁴ , Y. Zhang⁶ , Y. Z. Zhang^{4,b} , Z. Zhang^{4,b} , Y. Zhao²² , A. Zharkova⁴⁴ , A. Zhelezov²² , S. Z. Zheng⁶ , X. Z. Zheng^{4,b} , Y. Zheng⁷ , T. Zhou⁶ , X. Zhou⁸ , Y. Zhou⁷ , V. Zhovkovska⁵⁷ , L. Z. Zhu⁷ , X. Zhu^{4,b} , X. Zhu⁸ , V. Zhukov¹⁷ , J. Zhuo⁴⁸ , Q. Zou^{5,7} , D. Zuliani^{33,p} , G. Zunica⁵⁰ .

¹School of Physics and Astronomy, Monash University, Melbourne, Australia

²Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

³Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

⁴Department of Engineering Physics, Tsinghua University, Beijing, China

⁵Institute Of High Energy Physics (IHEP), Beijing, China

⁶School of Physics State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

⁷University of Chinese Academy of Sciences, Beijing, China

⁸Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China

⁹Consejo Nacional de Rectores (CONARE), San Jose, Costa Rica

¹⁰Université Savoie Mont Blanc, CNRS, IN2P3-LAPP, Annecy, France

¹¹Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France

¹²Université Paris-Saclay, Centre d’Etudes de Saclay (CEA), IRFU, Saclay, France, Gif-Sur-Yvette, France

¹³Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France

¹⁴Université Paris-Saclay, CNRS/IN2P3, IJCLab, Orsay, France

¹⁵Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

¹⁶LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France

¹⁷I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany

¹⁸Universität Bonn - Helmholtz-Institut für Strahlen und Kernphysik, Bonn, Germany

¹⁹Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany

- ²⁰ *Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, Freiburg, Germany*
- ²¹ *Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany*
- ²² *Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany*
- ²³ *School of Physics, University College Dublin, Dublin, Ireland*
- ²⁴ *INFN Sezione di Bari, Bari, Italy*
- ²⁵ *INFN Sezione di Bologna, Bologna, Italy*
- ²⁶ *INFN Sezione di Ferrara, Ferrara, Italy*
- ²⁷ *INFN Sezione di Firenze, Firenze, Italy*
- ²⁸ *INFN Laboratori Nazionali di Frascati, Frascati, Italy*
- ²⁹ *INFN Sezione di Genova, Genova, Italy*
- ³⁰ *INFN Sezione di Milano, Milano, Italy*
- ³¹ *INFN Sezione di Milano-Bicocca, Milano, Italy*
- ³² *INFN Sezione di Cagliari, Monserrato, Italy*
- ³³ *INFN Sezione di Padova, Padova, Italy*
- ³⁴ *INFN Sezione di Perugia, Perugia, Italy*
- ³⁵ *INFN Sezione di Pisa, Pisa, Italy*
- ³⁶ *INFN Sezione di Roma La Sapienza, Roma, Italy*
- ³⁷ *INFN Sezione di Roma Tor Vergata, Roma, Italy*
- ³⁸ *Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands*
- ³⁹ *Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, Netherlands*
- ⁴⁰ *AGH - University of Krakow, Faculty of Physics and Applied Computer Science, Kraków, Poland*
- ⁴¹ *Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*
- ⁴² *National Center for Nuclear Research (NCBJ), Warsaw, Poland*
- ⁴³ *Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*
- ⁴⁴ *Authors affiliated with an institute formerly covered by a cooperation agreement with CERN.*
- ⁴⁵ *ICCUB, Universitat de Barcelona, Barcelona, Spain*
- ⁴⁶ *La Salle, Universitat Ramon Llull, Barcelona, Spain*
- ⁴⁷ *Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain*
- ⁴⁸ *Instituto de Física Corpuscular, Centro Mixto Universidad de Valencia - CSIC, Valencia, Spain*
- ⁴⁹ *European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- ⁵⁰ *Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- ⁵¹ *Physik-Institut, Universität Zürich, Zürich, Switzerland*
- ⁵² *NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- ⁵³ *Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- ⁵⁴ *School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom*
- ⁵⁵ *H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- ⁵⁶ *Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- ⁵⁷ *Department of Physics, University of Warwick, Coventry, United Kingdom*
- ⁵⁸ *STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- ⁵⁹ *School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- ⁶⁰ *School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- ⁶¹ *Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- ⁶² *Imperial College London, London, United Kingdom*
- ⁶³ *Department of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- ⁶⁴ *Department of Physics, University of Oxford, Oxford, United Kingdom*
- ⁶⁵ *Massachusetts Institute of Technology, Cambridge, MA, United States*
- ⁶⁶ *University of Cincinnati, Cincinnati, OH, United States*
- ⁶⁷ *University of Maryland, College Park, MD, United States*
- ⁶⁸ *Los Alamos National Laboratory (LANL), Los Alamos, NM, United States*
- ⁶⁹ *Syracuse University, Syracuse, NY, United States*
- ⁷⁰ *Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to ³*
- ⁷¹ *School of Physics and Electronics, Hunan University, Changsha City, China, associated to ⁸*
- ⁷² *Guangdong Provincial Key Laboratory of Nuclear Science, Guangdong-Hong Kong Joint Laboratory of Quantum Matter, Institute of Quantum Matter, South China Normal University, Guangzhou, China,*

associated to ⁴

⁷³ Lanzhou University, Lanzhou, China, associated to ⁵

⁷⁴ School of Physics and Technology, Wuhan University, Wuhan, China, associated to ⁴

⁷⁵ Departamento de Física , Universidad Nacional de Colombia, Bogota, Colombia, associated to ¹⁶

⁷⁶ Ruhr Universitaet Bochum, Fakultaet f. Physik und Astronomie, Bochum, Germany, associated to ¹⁹

⁷⁷ Eotvos Lorand University, Budapest, Hungary, associated to ⁴⁹

⁷⁸ Van Swinderen Institute, University of Groningen, Groningen, Netherlands, associated to ³⁸

⁷⁹ Universiteit Maastricht, Maastricht, Netherlands, associated to ³⁸

⁸⁰ Tadeusz Kosciuszko Cracow University of Technology, Cracow, Poland, associated to ⁴¹

⁸¹ Universidade da Coruña, A Coruña, Spain, associated to ⁴⁶

⁸² Department of Physics and Astronomy, Uppsala University, Uppsala, Sweden, associated to ⁶⁰

⁸³ University of Michigan, Ann Arbor, MI, United States, associated to ⁶⁹

^a Centro Federal de Educação Tecnológica Celso Suckow da Fonseca, Rio De Janeiro, Brazil

^b Center for High Energy Physics, Tsinghua University, Beijing, China

^c Hangzhou Institute for Advanced Study, UCAS, Hangzhou, China

^d School of Physics and Electronics, Henan University , Kaifeng, China

^e LIP6, Sorbonne Université, Paris, France

^f Lamarr Institute for Machine Learning and Artificial Intelligence, Dortmund, Germany

^g Universidad Nacional Autónoma de Honduras, Tegucigalpa, Honduras

^h Università di Bari, Bari, Italy

ⁱ Università di Bergamo, Bergamo, Italy

^j Università di Bologna, Bologna, Italy

^k Università di Cagliari, Cagliari, Italy

^l Università di Ferrara, Ferrara, Italy

^m Università di Genova, Genova, Italy

ⁿ Università degli Studi di Milano, Milano, Italy

^o Università degli Studi di Milano-Bicocca, Milano, Italy

^p Università di Padova, Padova, Italy

^q Università di Perugia, Perugia, Italy

^r Scuola Normale Superiore, Pisa, Italy

^s Università di Pisa, Pisa, Italy

^t Università della Basilicata, Potenza, Italy

^u Università di Roma Tor Vergata, Roma, Italy

^v Università di Siena, Siena, Italy

^w Università di Urbino, Urbino, Italy

^x Universidad de Alcalá, Alcalá de Henares , Spain

^y Facultad de Ciencias Físicas, Madrid, Spain

^z Department of Physics/Division of Particle Physics, Lund, Sweden

[†] Deceased