


HH

ISSN 0029-3865



CBPF - CENTRO BRASILEIRO DE PESQUISAS FÍSICAS
Rio de Janeiro


Notas de Física

CBPF-NF-057/96

September 1996

A Deformed Bose-Einstein Gas Near $q = 1$

A. B. Pinheiro and I. Roditi



sw9648



CNPq - Conselho Nacional de Desenvolvimento Científico e Tecnológico

A Deformed Bose-Einstein Gas Near $q = 1$

A. B. Pinheiro^a and I. Roditi^b

Centro Brasileiro de Pesquisas Físicas - CBPF
Rua Dr. Xavier Sigaud, 150
22290-180 - Rio de Janeiro, RJ - Brazil

ABSTRACT

We show that the Bose-Einstein condensation occurs, on the region near $q = 1$, for a modified ideal deformed gas.

Key-words: q-Deformation; Bose-Einstein condensation.

e-mail address:

^(a)baptista@CBPF.SU1.CAT.CBPF.BR

^(b)roditi@CBPF.SU1.CAT.CBPF.BR

Recently, a great deal of interest has been devoted to the study of quantum groups [1-4] and deformed quantum q -gases. Chaichian *et al* [5] investigated the deformed Bose-Einstein gas near $q = 1$ and found that in this limit some divergencies appear in the series expansion of the thermodynamic functions. This fact did not allow the study of the Bose-Einstein condensation (BEC). Later on, R-Monteiro *et al* [6, 7] showed that, for highly deformed q -gases, BEC is present and the specific heat has a λ -point transition behavior.

We propose here a new type of deformed system in which the parameter q is a function of the momentum, p . This procedure is implemented by means of a system where each oscillator is attached to a different value of q , and for particular choices it is possible to regulate the thermodynamic functions. When the Bose-Einstein gas is deformed this way, we show that it is possible to approach the region near $q = 1$ and BEC occurs on this system.

The Hamiltonian for an ideal q -gas is taken as [5, 6]

$$H = \sum_i \hbar\omega_i a_i^\dagger a_i = \sum_i \hbar\omega_i [N_i], \quad (1)$$

where a_i, a_i^\dagger, N_i are annihilation, creation and occupation operators, respectively, of particles in the state i . These operators have the following commutation relations:

$$a_i a_j^\dagger - ((q_i - 1)\delta_{ij} + 1)a_j^\dagger a_i = \delta_{ij} q^{-N_i}, \quad [N_i, a_j] = -\delta_{ij} a_j, \quad [N_i, a_j^\dagger] = \delta_{ij} a_j^\dagger. \quad (2)$$

We also have

$$[N_i] = (q_i^{N_i} - q_i^{-N_i}) / (q_i - q_i^{-1}). \quad (3)$$

As in the non-deformed case, the total energy can be written as a sum of single-particle energies :

$$E = \sum_i E_i(n_i) = \sum_i \hbar\omega_i [n_i]. \quad (4)$$

For real $q_i = \exp(t_i)$ we have for the state i

$$E_i(n) = \hbar\omega_i \frac{\sinh(nt_i)}{\sinh(t_i)}. \quad (5)$$

The grand canonical partition function of the system is

$$Z(z, V, T) = \text{Tr}\{\exp[-\beta(H - \mu N)]\} = \sum_{N=0}^{\infty} z^N Z_N(V, T), \quad (6)$$

where $z = \exp(\beta\mu)$ is the fugacity, μ is the chemical potential, N is the total number operator and Z_N is the canonical partition function :

$$Z_N = \sum_{n_i} \exp(-\beta\hbar \sum_i \omega_i [n_i]) \quad (7)$$

Further calculations provide that

$$Z = \prod_i \sum_{n=0}^{\infty} z^n \exp(-\beta \hbar \omega_i [n]) \quad (8)$$

So, the grand canonical potential is given by

$$\Omega = -\frac{1}{\beta} \ln Z = -\frac{1}{\beta} \sum_i \ln \left(\sum_{n=0}^{\infty} z^n \exp(-\beta \hbar \omega_i [n]) \right) \quad (9)$$

Making the assumption that all t_i are small enough, and then expanding equation (5) to order t_i^2 , we have :

$$E_i = \hbar \omega_i \left(n + \frac{1}{6} n(n+1)(n-1) t_i^2 + \dots \right). \quad (10)$$

The substitution of this in (8) gives

$$z^n Z_N(V, T) \simeq \sum_{n=0}^{\infty} e^{\beta(\mu - \hbar \omega) n} \left(1 - \frac{1}{6} n(n+1)(n-1) \beta \hbar \omega t_i^2 \right), \quad (11)$$

and makes it possible to perform the sum over n :

$$z^n Z_N(V, T) \simeq Z_0 (1 - x_i z^2 e^{-2x_i} Z_0^3 t_i^2), \quad (12)$$

where $x_i = \beta \hbar \omega_i$ and $Z_0 = 1/(1 - ze^{-x_i})$. Finally, the grand canonical potential is, to second order in t_i ,

$$\beta \Omega \simeq - \sum_i \ln Z_0 + z^2 \sum_i t_i^2 x_i e^{-2x_i} Z_0^3 \quad (13)$$

The next step is to enclose the system in a 3-dimensional volume V and replace the sum over levels by an integral over \vec{p} -space:

$$\sum_i \rightarrow \frac{4\pi V}{h^3} \int_0^{\infty} p^2 dp, \quad (14)$$

and make t a function of p by choosing

$$t(p, \epsilon) = \sqrt{\epsilon \frac{p^\eta}{(\Lambda)^\eta} e^{-\beta p^2/2m}}, \quad (15)$$

where $\Lambda = (2m/\beta)^{1/2}$, η is an integer, $\epsilon \ll 1$ and $\hbar \omega = p^2/2m$ defines the dispersion law for non-relativistic particles.

Remembering that $p = -\Omega/V$, we have

$$\beta p = \lambda^{-3} (g_{5/2}(z) - \epsilon G_{5/2}(\eta, z)), \quad (16)$$

where $\lambda = h/\sqrt{2\pi mkT}$, and the g_α and G functions are integrals of the form [8]

$$I(r, \alpha, \nu) = \int_0^\infty dx z^r \frac{x^{\alpha-1}}{(e^x - z)^\nu} = \frac{\Gamma(\alpha)}{(\nu-1)!} \sum_{k=0}^\infty \frac{(k+\nu-1)!}{k!} \frac{z^k}{(k+\nu)^\alpha} \quad (17)$$

For g_α , we have

$$g_\alpha = \frac{1}{\Gamma(\alpha)} I(1, \alpha, 1). \quad (18)$$

These functions have the following property:

$$z \frac{dg_\alpha(z)}{dz} = g_{\alpha-1}(z). \quad (19)$$

The G functions do not have the above property, as $\alpha = (\eta + 5)/2$ always. However, we can, for the sake of simplicity, give them the same subscript of the g functions. Then, the G functions obey (18) in a formal manner, and are defined by

$$\begin{aligned} G_{5/2}(\eta, z) &= \frac{2}{\sqrt{\pi}} I(2, \frac{\eta+5}{2}, 3), \\ G_{3/2}(\eta, z) &= \frac{2}{\sqrt{\pi}} [2I(2, \frac{\eta+5}{2}, 3) + 3I(3, \frac{\eta+5}{2}, 4)], \\ G_{1/2}(\eta, z) &= \frac{2}{\sqrt{\pi}} [4I(2, \frac{\eta+5}{2}, 3) + 15I(3, \frac{\eta+5}{2}, 4) + 12I(4, \frac{\eta+5}{2}, 5)] \end{aligned} \quad (20)$$

Chaichian *et al* [5] showed that the expansion in t , eq.(12), suffers a breakdown when $z \rightarrow 1$ and $x \rightarrow 0$, as the effective expansion parameter, t^2 times a positive power of Z_0 , grows large. It is not the case if we use eq.(15), making t a function of p , or, in terms of x , $t^2 = \epsilon x^{\eta/2} e^{-x}$. Doing so, we have a modified deformed system where the effective expansion parameter has a finite value when $z \rightarrow 1$ and $x \rightarrow 0$, if $\eta \geq 4$. In our calculations, we used $\eta = 4$.

The density of the system is

$$n = z \left(\frac{\partial(\beta p)}{\partial z} \right)_\beta, \quad (21)$$

and we get

$$n = \lambda^{-3} (g_{3/2}(z) - \epsilon G_{3/2}(\eta, z)). \quad (22)$$

The internal energy U and $(\partial z / \partial T)_V$ are, respectively, equal to

$$\frac{3}{2} \frac{T(g_{5/2}(z) - \epsilon G_{5/2}(\eta, z))}{g_{3/2}(z) - \epsilon G_{3/2}(\eta, z)} \quad (23)$$

and

$$-\frac{3}{2} \frac{z g_{3/2}(z)}{T g_{1/2}(z)} + \frac{3}{2} \frac{\epsilon z G_{3/2}(\eta, z)}{T g_{1/2}(z)}. \quad (24)$$

Using these equations, we find the specific heat of the system, for $T \geq T_c$:

$$\begin{aligned} \frac{C_V}{Nk} = & -\frac{15}{4} \frac{\epsilon G_{5/2}(\eta, z)}{g_{3/2}(z) - \epsilon G_{3/2}(\eta, z)} + \frac{9}{2} \frac{\epsilon g_{3/2}(z) G_{3/2}(\eta, z)}{g_{1/2}(z)(g_{3/2}(z) - \epsilon G_{3/2}(\eta, z))} \\ & - \frac{9}{4} \frac{\epsilon g_{5/2}(z) G_{1/2}(\eta, z)}{g_{1/2}(z)(g_{3/2}(z) - \epsilon G_{3/2}(\eta, z))} \\ & + \frac{15}{4} \frac{g_{5/2}(z)}{g_{3/2}(z) - \epsilon G_{3/2}(\eta, z)} - \frac{9}{4} \frac{g_{3/2}(z)^2}{g_{1/2}(z)(g_{3/2}(z) - \epsilon G_{3/2}(\eta, z))} \end{aligned} \quad (25)$$

For $z = 1$ or $T \leq T_c$, we have

$$\frac{C_V}{Nk} = \frac{15 g_{5/2}(1) - \epsilon G_{5/2}(\eta, 1)}{4 g_{3/2}(1) - \epsilon G_{3/2}(\eta, 1)} \left(\frac{T}{T_c} \right)^{3/2} \quad (26)$$

We note that, as $\epsilon \rightarrow 0$, we recover the ideal Bose-Einstein gas specific heat. In figure 1 we see that, as $z \rightarrow 0$ or $T \rightarrow \infty$, the specific heat limit is also $3/2$. The critical temperature, T_c , is

$$T_c = \frac{h^2}{2\pi m k} \left(\frac{n}{g_{3/2}(z) - \epsilon G_{3/2}(\eta, z)} \right)^{2/3}. \quad (27)$$

When $\epsilon \rightarrow 0$, we recover the critical temperature of the Bose-Einstein ideal gas.

Although our system dynamics is different from the ones in [5] and [7], there are some similarities. For example, our T_c is higher than that of the ideal Bose-Einstein gas, as in [7]. However, our specific heat does not show any discontinuity at $T = T_c$, as we see in the figure. The reason to that is the fact that our system averages the thermodynamic functions for values of q such that $1 \leq q \leq 1 + \epsilon$. Notice, however, that for increasing ϵ , the trend of the curves is compatible with the appearance of a λ - point transition at large q . Finally, we point out that the type of deformation proposed here makes the deformation parameter continuous. This fact enabled us to study BEC near $q = 1$, which has not been achieved for the usual deformed Bose-Einstein gas previously investigated.

Acknowledgements - The authors thank M.R-Monteiro for suggesting the present subject and for permanent interest. This work was supported by CNPq - Conselho Nacional de Desenvolvimento Científico e Tecnológico.




FIG.1. Specific heat of the ideal ($q = 1$) and deformed ($q=1+t(p,0.1)$) Bose - Einstein gas.

References

- [1] V.G.Drinfel'd, in Proc. of the Int. Congress of Math., Berkeley, 1986, reprinted in "Yang-Baxter Equation in Integrable Systems", Ed. Michio Jimbo, World Scientific, Singapore, 1989.
- [2] M.Jimbo, Lett.Math.Phys. 10, 63 (1985); 11,247 (1986).
- [3] A.J.MacFarlane, J.Phys.**A** 22, 4581 (1989).
- [4] L.C.Biedenharn, J.Phys.**A** 22, L873 (1989).
- [5] M.Chaichian, R.Gonzalez Felipe and C.Montonen, J.Phys.**A** 26, 4017 (1993).
- [6] M.R-Monteiro, I.Roditi and L.M.C.S.Rodrigues, Mod.Phys.Lett.**B** 7, 1897 (1993).
- [7] M.R-Monteiro, I.Roditi and L.M.C.S.Rodrigues, Int.J.Mod.Phys.**B** 8, 3281 (1994), and references therein.
- [8] A.P.Prudnikov, Yu.A.Brychkov, and O.I.Marichev, "Integrals and Series", vol.1, Gordon and Breach Science Publishers, New York, 1988.

NOTAS DE FÍSICA é uma pré-publicação de trabalho original em Física.
Pedidos de cópias desta publicação devem ser enviados aos autores ou ao:

Centro Brasileiro de Pesquisas Físicas
Área de Publicações
Rua Dr. Xavier Sigaud, 150 – 4^o andar
22290-180 – Rio de Janeiro, RJ
Brasil

NOTAS DE FÍSICA is a preprint of original unpublished works in Physics.
Requests for copies of these reports should be addressed to:

Centro Brasileiro de Pesquisas Físicas
Área de Publicações
Rua Dr. Xavier Sigaud, 150 – 4^o andar
22290-180 – Rio de Janeiro, RJ
Brazil

