

INSTITUTO ARGENTINO DE MATEMATICA - CONICET  
Y  
FACULTAD DE CIENCIAS EXACTAS Y NATURALES  
DEPARTAMENTO DE MATEMATICA - UBA

# Trabajos de Matemática

SCAN-9902051



CERN LIBRARIES, GENEVA

**PREPRINT 272**

SW9907

"The residues of the resolvent of the  
Laplacian on Damek-Ricci spaces".

**R.J. Miatello**  
**C.E. Will**

**1998**

## Publicaciones Previas

ISSN 0325-6677

---

## **Comité de Redacción**

Gustavo Corach

Norberto A. Fava

Angel R. Larotonda

Impreso por WorldPress Gráfica Digital (01)826-8976

# THE RESIDUES OF THE RESOLVENT OF THE LAPLACIAN ON DAMEK-RICCI SPACES.

R.J. MIATELLO\* AND C.E. WILL\*

Facultad de Matemática, Astronomía y Física,  
Universidad Nacional de Córdoba, 5000 Córdoba, Argentina.

## 1. INTRODUCTION.

Let  $N$  be an  $H$ -type group and let  $S$  be the canonical solvable extension, as defined in [3]. Harmonic analysis on  $S$  has strong similarities with that of symmetric spaces of negative curvature. In particular, it was proved by Damek-Ricci that  $S$  is a riemannian harmonic space.

In [7] the resolvent of the Laplacian  $R(\mu)$  was studied on symmetric and locally symmetric spaces of negative curvature and in particular it was shown that  $R(\mu)$  has a meromorphic continuation to  $\mathbf{C}$ .

In this paper we announce some recent results on the poles and residues of the resolvent  $R(\mu)$  on a general Damek-Ricci space  $S$ , namely all poles are simple and the residues are convolution operators which have finite rank (see Theorem 3.4). If  $S$  corresponds to a symmetric space of negative curvature, the image of each residue is a  $\mathfrak{g}_e$ -module with a specific highest weight, hence its dimension can be computed by Weyl dimension formula (see Theorem 3.5). We specialize our results at the end, by discussing the case of the real hyperbolic  $n$ -space, which was treated by Guillopé-Zworski in [4].

## 2. PRELIMINARIES.

In this section we will recall some basic notions on  $H$ -type groups and their canonical solvable extensions, following mainly [3] (see also [1]). We will also review some basic facts on symmetric spaces of negative curvature.

Let  $\mathfrak{n}$  be a two-step real nilpotent Lie algebra endowed with an inner product  $\langle \cdot, \cdot \rangle$ . Assume  $\mathfrak{n}$  has an orthogonal decomposition  $\mathfrak{n} = \mathfrak{z} \oplus \mathfrak{v}$ , where  $\mathfrak{z}$  is the center of  $\mathfrak{n}$  and  $[\mathfrak{v}, \mathfrak{v}] = \mathfrak{z}$ . Define a linear mapping  $J : \mathfrak{z} \rightarrow \text{End}(\mathfrak{v})$  by

$$(1) \quad \langle J_Z X, Y \rangle = \langle Z, [X, Y] \rangle$$

(note that  $J_Z$  is skew-symmetric). Now  $\mathfrak{n}$  is said to be an  $H$ -type algebra if for any  $Z_1, Z_2 \in \mathfrak{z}$

$$(2) \quad J_{Z_1} J_{Z_2} + J_{Z_2} J_{Z_1} = -2\langle Z_1, Z_2 \rangle$$

---

\* Partially supported by Conicet, Conicor, SecytUNC (Córdoba), and I.C.T.P. (Trieste).

The corresponding  $H$ -type group is the simply connected Lie group  $N$  with Lie algebra  $\mathfrak{n}$  endowed with the left invariant metric induced by the inner product  $\langle \cdot, \cdot \rangle$  on  $\mathfrak{n}$ .

Consider the solvable extension,  $S$ , of  $N$ , as defined in [3], that is,  $A = \mathbf{R}^+$  acts on  $N$  by the dilation  $(x, z) \rightarrow (t^{\frac{1}{2}}x, tz)$ , and  $S = AN$  is the semidirect product of  $A$  and  $N$ . If  $\mathfrak{s}$ ,  $\mathfrak{a}$ ,  $\mathfrak{n}$  denote respectively the Lie algebras of  $S$ ,  $A$  and  $N$ , we have that  $\mathfrak{s} = \mathfrak{a} \oplus \mathfrak{n}$ , where  $\mathfrak{a} = \mathbf{R}H$  acts on  $\mathfrak{n}$  via  $H$ , the derivation of  $\mathfrak{n}$  such that  $H|_{\mathfrak{v}} = \frac{1}{2}I$  and  $H|_{\mathfrak{z}} = I$ . We use the inner product on  $\mathfrak{s}$  extending the one on  $\mathfrak{n}$  and such that  $\|H\| = 1$ ,  $\langle H, \mathfrak{n} \rangle = 0$ ;  $S$  carries the induced left invariant riemannian structure. Also, we shall denote  $q = \dim \mathfrak{z}$ ,  $p = \dim \mathfrak{v}$ ,  $n = \dim \mathfrak{s} = p + q + 1$  and  $Q = \frac{1}{2}(p + 2q)$ . We point out that if  $\mathfrak{n}$  is abelian we shall use the convention that  $\mathfrak{v} = 0$ , and  $\mathfrak{n} = \mathfrak{z}$ .

Using coordinates form  $\mathfrak{v} \oplus \mathfrak{z} \oplus \mathbf{R}^+$ , the product on  $S$  is expressed as

$$(Z, Z, a)(X', Z', a') = (X + a^{\frac{1}{2}}X', Z + aZ' + \frac{1}{2}a^{\frac{1}{2}}[X, X'], aa').$$

Note that the volume element of the induced left invariant riemannian metric  $g$  on  $S$  is the left Haar measure

$$dm = a^{-Q-1}dXdZda.$$

We will need the fact that  $S$  can be realized as the unit ball in  $\mathfrak{s}$ :

$$B(\mathfrak{s}) = \{(X, S, u) : |X|^2 + |Z|^2 + u^2 = 1\}$$

via a Cayley type transform  $\tilde{C} : S \rightarrow B(\mathfrak{s})$ , where  $\tilde{C} = C^{-1}h$  in the notation of [3], Section 4. In  $B(\mathfrak{s})$  the geodesics through the origin are the diameters and the geodesic distance to the origin is given by  $r = d(\tilde{p}, 0) = \log \frac{1+|\tilde{p}|}{1-|\tilde{p}|}$ , thus  $|\tilde{p}| = \tanh(r/2)$ , where  $\tilde{p} = \tilde{C}(p)$  if  $p \in S$ . Since

$$|\tilde{C}^{-1}(X, Z, a)|^2 = 1 - \frac{4a}{(1 + a + \frac{1}{4}|X|^2)^2 + |Z|^2}$$

we have that

$$(3) \quad \cosh(r/2)^{-2} = \frac{4a}{(1 + a + \frac{1}{4}|X|^2)^2 + |Z|^2}$$

and furthermore, the image of the left Haar measure on  $S$  via  $\tilde{C}^{-1}$  is  $d\mu = J(r)d\sigma dr$  where  $r, \sigma$  are the radial coordinates on  $B$ ,  $r^2 = |X|^2 + |Z|^2 + u^2$  and  $J(r) = 2^p \sinh(r/2)^p \sinh(r)^q$  (see [3] Section 4).

The symmetric spaces of negative curvature are a main subclass of the Damek-Ricci spaces. Let  $G$  be a connected, non compact, semisimple Lie group of real rank one. Let  $K$  be a maximal compact subgroup of  $G$  and let  $\mathfrak{g}$  and  $\mathfrak{k}$  be the corresponding Lie algebras. If  $G = NAK$  is an Iwasawa decomposition of  $G$ , then  $N$  is an  $H$ -type group, and  $S = NA \approx G/K$  is a solvable group in the class introduced above. Indeed, if  $\mathfrak{a}$  and  $\mathfrak{n}$  denote the Lie algebras of  $A$  and  $N$  respectively, then  $\mathfrak{n}$  splits  $\mathfrak{n} = \mathfrak{g}_{\alpha/2} \oplus \mathfrak{g}_{\alpha}$ , where  $\mathfrak{g}_{j\alpha}$ ,  $j = 1/2, 1$ , denote the  $j\alpha$ -root spaces of  $\mathfrak{a}$ . In the notation above we have  $\mathfrak{n}_{\alpha} = \mathfrak{z}$ ,  $\mathfrak{n}_{\alpha/2} = \mathfrak{v}$ , and  $H = adH_o$  with  $H_o \in \mathfrak{a}$  such that  $\alpha(H_o) = 1$ . Then  $Q = 2\rho$ , where  $\rho = \frac{1}{4}(p + 2q)$ . If on  $S = NA$  we use the  $G$ -invariant metric induced by  $2(p + 4q)^{-1}B$  ( $B$  the Killing form of  $\mathfrak{g}$ ) then  $S$  is isometric to a Damek-Ricci space. We note that, because of our convention, if  $\mathfrak{n}$  is abelian then  $p = 0$ ,  $q = \dim \mathfrak{n}$ .

### 3. THE RESIDUES OF THE RESOLVENT OF THE LAPLACIAN ON $S$ .

Damek-Ricci spaces have similarities with symmetric spaces of negative curvature, in particular they are harmonic spaces. On  $S$  there is a radialization operator  $\pi$  which corresponds to the standard operator in the case of the ball model of  $S$  (see [3] p. 230). If  $f \in C_c^\infty(S)$ ,  $p \in S$  and  $\tilde{p} = \tilde{C}(p)$  then

$$\pi f(p) := \int_{S^{p+q}} \tilde{f}(\|\tilde{p}\|\sigma) d\sigma$$

where  $\tilde{f} := f \circ \tilde{C}^{-1}$ . This operator corresponds in the symmetric case to averaging over the action of  $K$  by left translation: if  $f \in C^\infty(NA)$ , then

$$(4) \quad \pi f(x) = \int_K \tilde{f}(kx) dk$$

where  $\tilde{f}$  denotes the right  $K$ -invariant extension of  $f$ .

If  $\{Z_i\}$ ,  $\{V_j\}$  are orthonormal bases of  $\mathfrak{z}$  and  $\mathfrak{v}$  respectively, the Laplacian-Beltrami operator is given by  $L = \sum_i Z_i^2 + \sum_j V_j^2 + H^2 - QH$  (see [2]). Since  $L$  is self-adjoint and  $S$  is a harmonic manifold, then  $L$  commutes with  $\pi$ . Moreover,  $L$  generates the algebra of left-invariant differential operators on  $S$  which commute with  $\pi$  (see [3] Theorem 5.2).

If  $f$  is smooth radial function on  $S - \{e\}$ , we will often abuse notation by writing  $f(r) = f(x)$ , where  $r = d(x, e)$ . The action of  $L$  on radial functions is thus given by (see [3])

$$(5) \quad Lf(r) = \frac{d^2}{dr^2} f(r) + \frac{1}{2}(p \coth(r/2) + 2q \coth(r)) \frac{d}{dr} f(r).$$

In the symmetric case, if  $\mathfrak{n}$  is not abelian and we set  $r = 2t$ , then  $Lf(t)$  corresponds to  $\frac{1}{4}Cf(a_t)$ , in the notation of [7] Section 1 (1). If  $\mathfrak{n}$  is abelian,  $L$  corresponds to  $C$ .

A spherical function  $\psi$  on  $S$  is a radial eigenfunction of the Laplace Beltrami operator such that  $\psi(e) = 1$ . This generalizes the corresponding notion in the symmetric case and has the following characterization ([3]):

**Proposition 3.1.** *Let  $\nu \in \mathbf{C}$ . The function  $\phi_\nu = \pi(a^{\nu+Q/2})$  is a spherical function with eigenvalue  $\lambda(\nu) = \nu^2 - Q^2/4$  and any spherical function on  $S$  is of this form. Furthermore  $\phi_\nu = \phi_{-\nu}$ , for each  $\nu \in \mathbf{C}$ .*

As in the symmetric case, we can express  $\phi_\nu$  by a hypergeometric function as follows. By letting  $z = -\sinh(r/2)$  equation

$$(6) \quad \left\{ \frac{d^2}{dr^2} + \frac{1}{2}(p \coth(r/2) + 2q \coth(r)) \frac{d}{dr} - \lambda(\nu) \right\} f_\nu(r) = 0$$

transforms into the hypergeometric equation with parameters  $a = Q/2 - \nu$ ,  $b = Q/2 + \nu$ , and  $c = n/2$ . Since  $\phi_\nu(e) = 1$ , it follows that

$$(7) \quad \phi_\nu(r) = F\left(-\nu + Q/2, \nu + Q/2, \frac{n}{2}, -\sinh(r/2)^2\right)$$

where  $F(a, b, c, z)$  denotes the hypergeometric function. Furthermore, if  $\text{Re } \nu > 0$ , the asymptotic behavior of  $\phi_\nu(r)$ , as  $r \rightarrow \infty$ , is given by (see [3], p. 239)

$$(8) \quad \phi_\nu(r) \sim c(\nu)e^{r(\nu+Q/2)}, \quad \text{where } c(\nu) = \frac{2^{-2\nu+Q}\Gamma(n/2)\Gamma(2\nu)}{\Gamma(\nu + Q/2)\Gamma(\nu + \frac{p+2}{4})}.$$

Here  $c(\nu)$  coincides with Harish Chandra's  $c$ -function in the symmetric case. The Plancherel measure,  $\mu(\nu) = (c(\nu)c(-\nu))^{-1}$ , can be written  $\mu(\nu) = c_o p(\nu)D(\nu)$ , where  $p(\nu)$  is the polynomial given by:

$$\begin{aligned} & \prod_{j=0}^{\frac{q}{2}-1} \left( -\nu^2 + \left( \frac{2j+1}{2} \right)^2 \right) \prod_{j=0}^{\frac{q}{2}-1} \left( -\nu^2 + \left( \frac{j}{2} \right)^2 \right), & q, \frac{p}{2} \text{ even.} \\ & - \prod_{j=1}^{p/4} (-\nu^2 + j^2) \nu^3, & q = 1, \frac{p}{2} \text{ odd.} \\ & - \prod_{j=0}^{\frac{q}{2}-1} \left( -\nu^2 + \left( \frac{2j+1}{2} \right)^2 \right) \prod_{j=0}^{\frac{q}{2}-1} \left( -\nu^2 + \left( \frac{2j+1}{2} \right)^2 \right) \nu, & q \text{ odd, } \frac{p}{2} \text{ even.} \end{aligned}$$

$c_o$  is a constant, and in each case  $D(\nu)$  is given respectively by 1,  $\cot(\pi\nu)$ , and  $\text{tg}(\pi\nu)$  ([ADY]).

**Remark 3.2.** We note that  $p$  is always even, since  $\mathfrak{v}$  is a module over the Clifford algebra of  $\mathfrak{z}$ . If  $p = 0$ , then  $X \approx H^{q+1}$ ,  $G \simeq SO(q+1, 1)$  and in this case  $D(\nu)$  equals 1 or  $\text{tg}(\pi\nu)$  depending on whether  $q$  is even or odd.

In [7], the resolvent of the Laplacian was studied on symmetric (and locally symmetric spaces) of negative curvature. In the symmetric case it was shown that it is given by convolution with a smooth radial function  $Q_\nu$  on  $S - \{e\}$  which is an eigenfunction of  $L$  with eigenvalue  $\lambda(\nu) = \nu^2 - \frac{Q^2}{4}$ . It was also shown that  $R(\lambda(\nu))$  has a meromorphic continuation to  $\mathbf{C}$ . These properties remain valid for any  $S$  as above and some arguments in [7], can be adapted. On the other hand, it is more convenient to give a different construction of  $Q_\nu$ , using a series solution.

Now for  $b \in \mathbf{R}$  and  $\delta > 0$ , let  $\mathcal{S}_{b,\delta} = \{\nu : \text{Re } \nu > b \text{ and } |\nu + j| > \delta \forall j \in -\mathbf{N} : b \leq j\}$ . That is  $\mathcal{S}_{b,\delta} = \{\nu : \text{Re } \nu > b\}$ , if  $b \geq 0$ , and  $\mathcal{S}_{b,\delta}$  is a half plane with finitely many discs removed, centered at  $-1, -2, \dots, -k$ , where  $-k \geq b$ , if  $b < 0$ . The following theorem gives the main properties of  $Q_\nu$ .

**Theorem 3.3.** *If  $\nu \in \mathbf{C}$ ,  $2\nu \notin -\mathbf{N}$ , then there exists a radial function  $Q_\nu \in C^\infty(S - \{e\})$  with the following properties*

(a)  $(L - \lambda(\nu))Q_\nu = 0$  and, for each  $s \in S$  the map  $\nu \mapsto Q_\nu(s)$  is holomorphic for  $2\nu \notin -\mathbf{N}$ . Furthermore, where defined,

$$(9) \quad \phi_\nu = c(-\nu)Q_\nu + c(\nu)Q_{-\nu}$$

(b) If  $f \in C^\infty(S - \{e\})$  is radial and  $Lf = \lambda(\nu)f$  with  $2\nu \notin -\mathbf{N}$  then  $f = a\phi_\nu + bQ_\nu$ , for some  $a, b \in \mathbf{C}$ .

(c) There exists a meromorphic function  $d(\nu)$  on  $\mathbf{C}$ , holomorphic if  $2\nu \notin -\mathbf{N}$ , such that, as  $r \rightarrow 0$ ,

$$Q_\nu(r) \sim d(\nu)r^{p+q-1} |\log r|^{\delta_{\nu+q,1}}.$$

(d)  $\lim_{r \rightarrow 0^+} J(r) \frac{d}{dt} Q_\nu(r) = -2\nu c(\nu)$ .

(e) If  $f \in C_c^\infty(S)$  and  $2\nu \notin -\mathbf{N}$  then

$$(10) \quad \int_S Q_\nu(x^{-1}y)(L - \lambda(\nu)I)f(y)dy = -2\nu c(\nu)f(x).$$

(f) For any  $b \in \mathbf{R}$ ,  $\delta, r_o > 0$ , there exists a constant  $K = K(b, \delta, r_o)$  such that  $|Q_\nu(r)| \leq K$  for any  $r \geq r_o$ ,  $\nu \in \mathcal{S}_{b,\delta}$ .

Let  $R(\mu)$  denote the resolvent of  $L$  at  $\mu$ . By (10), if  $\lambda(\nu) = \nu^2 - \frac{Q^2}{4}$ , then  $R(\lambda(\nu))$  is given by a kernel operator with kernel  $K_\nu(x, y) = -\frac{Q_\nu(x^{-1}y)}{2\nu c(\nu)}$ . The

following result gives the poles and residues of  $R(\lambda(\nu))$ , therefore the poles of  $\nu \mapsto -\frac{Q_\nu(x^{-1}y)}{2\nu c(\nu)}$ .

**Theorem 3.4.** *If  $p, q$  are both even then  $R(\lambda(\nu))$  is everywhere holomorphic. Otherwise, it has simple poles lying at  $\nu_k = -Q/2 - k$  with  $k \in \mathbf{N} \cup \{0\}$ . If  $\nu = \nu_k$ , then  $\text{Res}_{\nu=\nu_k} R(\lambda(\nu)) = (2\pi\nu_k)^{-1} p(\nu_k) f * \phi_\nu$  is a finite rank operator, for each  $k$ . Here  $p(\nu)$  is the polynomial part of the Plancherel measure  $\mu(\nu)$ .*

In the case when  $S$  is of symmetric type one can get more precise information on the operators  $T_{\nu_k}$ , by using representation theory. To do this we will first introduce some notation. The group of isometries  $G$  of  $S$  is a noncompact semisimple Lie group of real rank one.

Let  $\mathfrak{g}$ ,  $\mathfrak{k}$ ,  $N$ , and  $A$  be as in Section 1, let  $M$  be the centralizer of  $A$  in  $K$ ,  $P = MAN$  and let  $\mathfrak{p}$  be the Lie algebra of  $P$ . Extend  $\mathfrak{a}$  in the usual way to a Cartan subalgebra  $\mathfrak{h}_c = \mathfrak{a}_c + \mathfrak{h}_c^-$  of  $\mathfrak{g}$ , where  $\mathfrak{h}_c^-$  is a maximal abelian subalgebra of  $\mathfrak{m}$ , and introduce compatible orderings in the dual spaces of  $\mathfrak{a}$  and  $\mathfrak{a} + \sqrt{-1}\mathfrak{h}_c^-$ .

Let  $\Sigma^+(\Delta^+)$  denote the corresponding set of positive roots of the pair  $(\mathfrak{g}, \mathfrak{a})$  (respectively  $(\mathfrak{g}_c, \mathfrak{h}_c)$ ).

For  $\nu \in \mathbf{C}$ , let  $(\pi_\nu, H^\nu)$  be the spherical principal series representation of  $G$  (see [8], Section 3.6). The zonal spherical function  $\phi_\nu$  is given by

$$(11) \quad \phi_\nu(g) = \langle \pi_\nu(g) \mathbf{1}_\nu, \mathbf{1}_\nu \rangle$$

where  $\mathbf{1}_\nu \in H^\nu$  is the  $K$ -fixed vector given by  $\mathbf{1}_\nu(nak) = a^{(\nu+\rho)\alpha}$ ,  $n \in N$ ,  $a \in A$ ,  $k \in K$ , and  $\langle \cdot, \cdot \rangle$  is the standard inner product on  $H^\nu$ .

In the symmetric case, one can obtain more information on the dimension of  $\text{Res}_{\nu=\nu_k} R(\lambda(\nu)) = (2\pi\nu_k)^{-1} p(\nu_k) f * \tilde{\phi}_\nu$ .

**Theorem 3.5.** *Let  $S = G/K$  be a noncompact symmetric space of real rank one and let  $\nu_k = -\rho - k$  with  $k \in \mathbf{N} \cup \{0\}$ . Then  $\text{Im}(\text{Res}_{\nu=\nu_k} R(\lambda(\nu)))$  is an irreducible  $\mathfrak{g}_c$ -module of highest weight  $k\alpha$ . In particular, its dimension is given by the Weyl dimension formula.*

**Example 3.6.** *The real hyperbolic  $n$ -space.*

In [4], Section 2, Guillopé-Zworski consider the resolvent in the case when  $S \approx H^n$ , finding its poles and showing that the residues are operators of finite-rank. In this particular case one can obtain an explicit series expression for  $Q_\nu$ :

$$Q_\nu(r) = e^{-(\nu+\rho)r} \sum_{j=0}^{\infty} c_j(\nu) e^{-2jr}$$



and for the residues of  $R(\lambda(\nu))$ . If we choose  $c_o = 1$ , we have for  $j \geq 1$ :

$$(12) \quad c_j(\nu) = \frac{(\rho)_j(\nu + \rho)_j}{j! (\nu + 1)_j}.$$

Furthermore,  $c(\nu)$  can be written:

$$(13) \quad c(\nu) = \frac{2^{2\rho-1}\Gamma(n/2)}{\pi^{1/2}} \frac{\Gamma(\nu)}{\Gamma(\nu + \rho)}$$

and now, using the duplication formula for the Gamma function, we obtain:

$$(14) \quad \frac{Q_\nu(r)}{2\nu c(\nu)} = \frac{2^{-2n+3}}{(n-2)!} e^{-(\nu+\rho)r} \sum_{j=0}^{\infty} \frac{\Gamma(\rho+j)}{j!} \frac{\Gamma(\nu+\rho+j)}{\Gamma(\nu+j+1)} e^{-2jr}.$$

Also, if we take  $\mathcal{S}_{b,\delta}$  as in Theorem (3.3), it is not hard to see, by using Stirling's estimates, that there exists a constant  $K = K(b, \delta)$  such that the coefficients in the series are bounded by  $K j^{\rho-1} |\nu + j|^{\rho-1}$ , uniformly for  $\nu$  in  $\mathcal{S}_{b,\delta}$ .

If  $n$  is odd,  $\rho = \frac{n-1}{2} \in \mathbf{N}$ , hence the coefficients of the series in (14) are polynomial functions in  $\nu$  and  $R(\lambda(\nu))$  is everywhere holomorphic, in this case.

On the other hand, if  $n$  is even, (14) implies that the kernel of  $R(\lambda(\nu))$  is a meromorphic function with poles at  $\nu_k = -\rho - k$ ,  $k \in \mathbf{N} \cup \{0\}$ .

Since  $\Gamma(\nu + \rho + j)$  is holomorphic at  $\nu = \nu_k$  for  $j > k$ , in this case the residues are

$$\begin{aligned} \text{Res}_{\nu=\nu_k} \frac{Q_\nu(r)}{2\nu c(\nu)} &= \frac{2^{-2n+3}}{(n-2)!} \sum_{j=0}^k \frac{\Gamma(\rho+j)(-1)^{k-j}}{j!(k-j)!\Gamma(-k-\rho+j+1)} e^{-(2j-k)r} \\ &= \frac{2^{-2n+4}}{(n-2)!} \sum_{j=0}^{[n/2]} \frac{\Gamma(\rho+j)(-1)^{k-j}}{j!(k-j)!\Gamma(-k-\rho+j+1)} \cosh(2j-k)r \end{aligned}$$

since  $\frac{\Gamma(\rho+j)(-1)^k}{\Gamma(-k-\rho+j+1)} = \frac{\Gamma(\rho+k-j)}{\Gamma(-\rho-j+1)}$ , for  $0 \leq j \leq k$ .

We note finally that in this case  $G = SO(n, 1)$ ,  $K = SO(n-1)$  and one finds that the highest weight of the representation in Theorem 3.5 is  $k\lambda_1$  where  $\lambda_1$  is the first fundamental weight. Thus,  $V_k$  is isomorphic to the representation of  $G$  on  $\mathcal{H}_k$ , the space of homogeneous harmonic polynomials of degree  $k$  and the image of  $\text{Res}_{\nu=\nu_k} R(\lambda(\nu))$  has dimension  $\frac{(k+n-3)!(2k+n-2)}{k!(n-2)!}$  (see [6] Ch.4, (4.12)).

## REFERENCES

- [1] ANKER J.P., DAMEK E., YACOUB CH., *Spherical analysis on harmonic AN groups*, Annali della Scuola Normale Sup. di Pisa 23 (1996), 643-679.
- [2] E. DAMEK, *A Poisson kernel of Heisenberg type nilpotent Lie groups*, Colloquium Mathematicum 53 (1987), 239-247.
- [3] DAMEK E., RICCI F., *Harmonic analysis on solvable extensions of H-type groups*, Journal of Geometric Analysis 2 (1992), 213-248.
- [4] GUILLOPÉ L., ZWORSKI M., *Polynomial bounds for the number of resonances for some complete spaces of constant negative curvature near infinity*, Asymptotic Analysis 11 (1995), 1-22.
- [5] HELGASON S., *Groups and Geometric Analysis*, Pure and Applied Math. 113, Academic Press, 1984.
- [6] KNAPP A.W., *Representation Theory of Semisimple Groups*, Princeton Univ. Press, 1986.
- [7] MIATELLO R.J., WALLACH N.R., *The resolvent of the Laplacian on negatively curved locally symmetric spaces of finite volume*, Jour. Diff. Geometry 36 (1992), 663-698.
- [8] WALLACH N.R., *Real Reductive Groups I*, Pure and Applied Math. 132, Academic Press, 1988.

e-mail: [miatello@mate.uncor.edu](mailto:miatello@mate.uncor.edu), [cwill@mate.uncor.edu](mailto:cwill@mate.uncor.edu)

Received in July 1998.

## SERIE I - TRABAJOS DE MATEMATICA

PUBLICACIONES PREVIAS.

Director de Publicaciones: **Carlos Segovia Fernández**

- N° 260 “*Solutions to the mean curvature equation by fixed point methods*”, M.C. Mariani, D.F. Rial (1995).
- N° 261 “*The periodic parabolic eigenvalue problem with  $L^\infty$  weight*”, T. Godoy, E. Lami Dozo, S. Paczka, (1996).
- N° 262 “*Covariant non-linear non-equilibrium thermodynamics and the ergodic theory of stochastic and quantum flows*”, D. Rapoport, (1996).
- N° 263 “*A substitute of harmonic majorization*”, L. de Rosa y C. Segovia Fernández, (1996).
- N° 264 “*The integral representation of the elementary solution of the  $n$ -dimensional ultrahyperbolic Klein-Gordon operator iterated  $k$ -times*”, S.E. Trione, (1996).
- N° 265 “*On amenability and geometry of spaces of bounded representations*”, G. Corach y J.E. Galé, (1997).
- N° 266 “*Group conditional expectations of finite index*”, E. Andruchow y D. Stojanoff, (1997).
- N° 267 “*Geometry on state and weight orbits*”, A. Varela, (1998)
- N° 268 “*Approximation by ratios of bounded analytic functions*”, D. Suárez, (1998).
- N° 269 “*On the elementary  $(P \pm i0)^\lambda$  - ultrahyperbolic solution of the Klein-Gordon operator iterated  $k$ -times*”, S.E. Trione, (1998).
- N° 270 “*A generalization of a theorem by Calabi to the parabolic Monge-Ampère equation*”, C.E. Gutiérrez y Q. Huang, (1998).
- N° 271 “*Two basic lifting theorems in the continuous case*”, F. Peláez, (1998).

Toda correspondencia relativa a publicaciones debe dirigirse a:

Instituto Argentino de Matemática  
 Saavedra 15 - 3er. Piso  
 1083 - Buenos Aires  
 Argentina.

