

DIRECT PHOTON PRODUCTION IN 158 A GEV $^{208}\text{Pb}+^{208}\text{Pb}$ COLLISIONS

T. PEITZMANN

University of Münster, D-48149 Münster, Germany

FOR THE WA98 COLLABORATION

A measurement of direct photon production in $^{208}\text{Pb}+^{208}\text{Pb}$ collisions at 158 A GeV has been carried out in the CERN WA98 experiment. The invariant yield of direct photons in central collisions is extracted as a function of transverse momentum in the interval $0.5 < p_T < 4$ GeV/c. A significant direct photon signal, compared to statistical and systematical errors, is seen at $p_T > 1.5$ GeV/c. The results constitute the first observation of direct photons in ultrarelativistic heavy-ion collisions which could be significant for diagnosis of quark gluon plasma formation.

1 Introduction

The observation of a new phase of strongly interacting matter, the quark gluon plasma (QGP), is one of the most important goals of current nuclear physics research. To study QGP formation photons (both real and virtual) were one of the earliest proposed signatures^{3,4}. They are likely to escape from the system directly after production without further interaction, unlike hadrons. Thus, photons carry information on their emitting sources from throughout the entire collision history, including the initial hot and dense phase. Recently, it was shown by Aurenche et al.⁵ that photon production rates in the QGP when calculated up to two loop diagrams, are considerably greater than the earlier lowest order estimates⁶. Following this result, Srivastava⁷ has shown that at sufficiently high initial temperatures the photon yield from quark matter may significantly exceed the contribution from the hadronic matter to provide a direct probe of the quark matter phase.

A large number of measurements of prompt photon production at high transverse momentum ($p_T > 3$ GeV/c) exist for proton-proton, proton-antiproton, and proton-nucleus collisions (see e.g.⁸). First attempts to observe direct photon production in ultrarelativistic heavy-ion collisions with oxygen and sulphur beams found no significant excess^{9,10,11,12}. The WA80 collaboration¹² provided the most interesting result with a p_T dependent upper limit on the direct photon production in S+Au collisions at 200 A GeV. In this paper we report on the first observation of direct photon production in ultrarelativistic heavy-ion collisions.

2 Data Analysis

The results are from the CERN experiment WA98 ¹³ which consists of large acceptance photon and hadron spectrometers. Photons are measured with the WA98 lead-glass photon detector, LEDA, which consisted of 10,080 individual modules with photomultiplier readout. The detector was located at a distance of 21.5 m from the target and covered the pseudorapidity interval $2.35 < \eta < 2.95$ ($y_{cm} = 2.9$). The particle identification was supplemented by a charged particle veto detector in front of LEDA.

The results presented here were obtained from an analysis of the data taken with Pb beams in 1995 and 1996. The 20% most peripheral and the 10% most central reactions have been selected from the minimum bias cross section ($\sigma_{min.bias} \approx 6300$ mb) using the measured transverse energy E_T . In total, $\approx 6.7 \cdot 10^6$ central and $\approx 4.3 \cdot 10^6$ peripheral reactions have been analyzed.

The extraction of direct photons in the high multiplicity environment of heavy-ion collisions must be performed on a statistical basis by comparison of the measured inclusive photon spectra to the background expected from hadronic decays. Neutral pions and η mesons are reconstructed via their $\gamma\gamma$ decay branch. For a detailed description of the detectors and the analysis procedure see ¹⁴.

The final measured inclusive photon spectra are then compared to the calculated background photon spectra to check for a possible photon excess beyond that from long-lived radiative decays. The background calculation is based on the measured π^0 spectra and the measured η/π^0 -ratio. The spectral shapes of other hadrons having radiative decays are calculated assuming m_T -scaling ¹⁵ with yields relative to π^0 's taken from the literature. It should be noted that the measured contribution (from π^0 and η) amounts to $\approx 97\%$ of the total photon background.

3 Results

Fig. 1 shows the fully corrected inclusive photon spectra for peripheral and central collisions. The spectra cover the p_T range of 0.3 – 4.0 GeV/c (slightly less for peripheral collisions) and extend over six orders of magnitude. Fig. 1 also shows the distributions of neutral pions which extend over a similar momentum range with slightly larger statistical errors.

The ratio of measured photons to calculated background photons is displayed in Fig. 2 as a function of transverse momentum. The upper plot shows the ratio for peripheral collisions which is seen to be compatible with one, i.e. no indication of a direct photon excess is observed. The lower plot shows the




Figure 1. The inclusive photon (circles) and π^0 (squares) transverse momentum distributions for peripheral (open points) and central (solid points) 158 A GeV $^{208}\text{Pb}+^{208}\text{Pb}$ collisions. The data have been corrected for efficiency and acceptance. Only statistical errors are shown.

same ratio for central collisions. It rises from a value of ≈ 1 at low p_T to exhibit an excess of about 20% at high p_T .

A careful study of possible systematical errors is crucial for the direct photon analysis. The largest contributions are from the γ and π^0 identification efficiencies and the uncertainties related to the η measurement. It should be emphasized that the inclusive photon and neutral meson (the basis for the background calculation) yields have been extracted from the same detector for exactly the same data sample. This decreases the sensitivity to many detector related errors and eliminates all errors associated with trigger bias or absolute yield normalization. Full details on the systematical error estimates are given in ¹⁴. The total p_T -dependent systematical errors are shown by the shaded regions in Fig. 2. A significant photon excess is clearly observed in




Figure 2. The $\gamma_{\text{Meas}}/\gamma_{\text{Bkgd}}$ ratio as a function of transverse momentum for peripheral (part a)) and central (part b)) 158 A GeV $^{208}\text{Pb}+^{208}\text{Pb}$ collisions. The errors on the data points indicate the statistical errors only. The p_T -dependent systematical errors are indicated by the shaded bands.

central collisions for $p_T > 1.5 \text{ GeV}/c$.

The final invariant direct photon yield per central collision is presented in Fig. 3. The statistical and asymmetric systematical errors of Fig. 2 are added in quadrature to obtain the total upper and lower errors shown in Fig. 3. An additional p_T -dependent error is included to account for that portion of the uncertainty in the energy scale which cancels in the ratios. In the case that the lower error is less than zero a downward arrow is shown with the tail of the arrow indicating the 90% confidence level upper limit ($\gamma_{\text{Excess}} + 1.28 \sigma_{\text{Upper}}$).

No published prompt photon results exist for proton-induced reactions at the \sqrt{s} of the present measurement. Instead, prompt photon yields for proton-induced reactions on fixed targets at 200 GeV are shown in Fig. 3 for comparison^{16,17,18}. These results have been scaled for comparison with the present measurements according to the calculated average number of nucleon-




Figure 3. The invariant direct photon multiplicity for central 158 A GeV $^{208}\text{Pb}+^{208}\text{Pb}$ collisions. The error bars indicate the combined statistical and systematical errors. Data points with downward arrows indicate unbounded 90% CL upper limits. Results of several direct photon measurements for proton-induced reactions have been scaled to central $^{208}\text{Pb}+^{208}\text{Pb}$ collisions for comparison.

nucleon collisions (660) for the central Pb+Pb event selection and according to the beam energy under the assumption that $E d^3 \sigma_{\gamma} / dp^3 = f(x_T) / s^2$, where $x_T = 2p_T / \sqrt{s}$ ¹⁹. This comparison indicates that the observed direct photon production in central $^{208}\text{Pb}+^{208}\text{Pb}$ collisions has a shape similar to that expected for proton-induced reactions at the same \sqrt{s} but a yield which is enhanced.

4 Summary

The first observation of direct photons in ultrarelativistic heavy-ion collisions has been presented. While peripheral Pb+Pb collisions exhibit no significant

photon excess, the 10% most central reactions show a clear excess of direct photons in the range of p_T greater than about 1.5 GeV/ c . The invariant direct photon multiplicity as a function of transverse momentum was presented for central $^{208}\text{Pb}+^{208}\text{Pb}$ collisions and compared to proton-induced results at similar incident energy. The comparison indicates excess direct photon production in central $^{208}\text{Pb}+^{208}\text{Pb}$ collisions beyond that expected from proton-induced reactions. The result suggests modification of the prompt photon production in nucleus-nucleus collisions, or additional contributions from pre-equilibrium or thermal photon emission. The result should provide a stringent test for different reaction scenarios, including those with quark gluon plasma formation, and may provide information on the initial temperature attained in these collisions.

References

1. M.C. Abreu et al., Phys. Lett. B **410**, 337 (1997).
2. E. Andersen et al., Phys. Lett. B **449**, 401 (1999).
3. E.L. Feinberg, Nuovo Cimento **34 A**, 391 (1976).
4. E. Shuryak, Phys. Lett. B **78**, 150 (1978).
5. P. Aurenche, F. Gelis, H. Zaraket, and R. Kobes, Phys. Rev. D **58**, 085003 (1998).
6. J. Kapusta, P. Lichard, and D. Seibert, Phys. Rev. D **44**, 2774 (1991).
7. D.K. Srivastava, Eur. Phys. J. C **10**, 487 (1999).
8. W. Vogelsang and M.R. Whalley, J. Phys. G: Nucl. Part. Phys. **23**, A1 (1997).
9. HELIOS Collaboration, T. Åkesson et al., Z. Phys. C **46**, 369 (1990).
10. WA80 Collaboration, R. Albrecht et al., Z. Phys. C **51**, 1 (1991).
11. CERES Collaboration, R. Baur et al, Z. Phys. C **71**, 571 (1996).
12. WA80 Collaboration, R. Albrecht et al., Phys. Rev. Lett. **76**, 3506 (1996).
13. WA98 Collaboration, *Proposal for a large acceptance hadron and photon spectrometer*, 1991, Preprint CERN/SPSLC 91-17, SPSLC/P260
14. WA98 Collaboration, M.M. Aggarwal et al, nucl-ex/0006007, submitted to Phys. Rev. C.
15. M. Bourquin and J.-M. Gaillard, Nucl. Phys. B **114**, 334 (1976).
16. E704 Collaboration, D.L. Adams et al., Phys. Lett. B **345**, 569 (1995).
17. E629 Collaboration, M. McLaughlin et al., Phys. Rev. Lett. **51**, 971 (1983).
18. NA3 Collaboration, J. Badier et al., Z. Phys. C **31**, 341 (1986).
19. J.F. Owens, Rev. Mod. Phys. **59**, 465 (1987).