



|                  |   |
|------------------|---|
| Title            | Effect of chemical fertilizer and manure application on N <sub>2</sub> O emission from reed canary grassland in Hokkaido, Japan                                 |
| Author(s)        | Jin, Tao; Shimizu, Mariko; Marutani, Satoru; Desyatkin, Alexey R.; Iizuka, Natsumi; Hata, Hiroshi; Hatano, Ryusuke  |
| Citation         | Soil Science & Plant Nutrition, 56(1), 53-65<br><a href="https://doi.org/10.1111/j.1747-0765.2010.00447.x">https://doi.org/10.1111/j.1747-0765.2010.00447.x</a> |
| Issue Date       | 2010-02   |
| Doc URL          | <a href="http://hdl.handle.net/2115/44778">http://hdl.handle.net/2115/44778</a>   |
| Rights           | The definitive version is available at <a href="http://www.blackwellsynergy.com">www.blackwellsynergy.com</a>   |
| Type             | article (author version)  |
| File Information | SSPN56-1_53-65.pdf  |



[Instructions for use](#)

1  
2  
3  
4  
5 **1 Title**  
6

7  
8 2 Effect of chemical fertilizer and manure application on N<sub>2</sub>O emission from reed canary  
9  
10 3 grassland in Hokkaido, Japan  
11

12  
13 **4 Names of authors**  
14

15 5 Tao Jin<sup>1</sup>, Mariko Shimizu<sup>1</sup>, Satoru Marutani<sup>1</sup>, Alexey R. Desyatkin<sup>1</sup>, Natsumi Iizuka<sup>1</sup>,  
16

17  
18 6 Hiroshi Hata<sup>2</sup>, Ryusuke Hatano<sup>1</sup>  
19

20  
21 7  
22

23 **8 Affiliations**  
24

25  
26 9 <sup>1</sup>Soil Science Laboratory, Graduate School of Agriculture, Hokkaido University, Kita 9  
27

28  
29 10 Nishi 9, Kita-ku, Sapporo, Hokkaido 060-8589, Japan  
30

31 11 <sup>2</sup>Field Science Center for Northern Biosphere, Agro-ecosystem Research Station,  
32

33  
34 12 Shizunai Livestock Farm, Hokkaido University, Kita 11 Nishi 10, Kita-ku, Sapporo,  
35

36 13 Hokkaido 060-0811, Japan  
37  
38

39 14  
40

41 **15 Corresponding author**  
42

43  
44 16 Tao Jin  
45

46  
47 17 Tel.: +81 11 706 2503  
48

49 18 Fax.: +81 11 706 2494  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

19 E-mail address: jintao@chem.agr.hokudai.ac.jp

20 Soil Science Laboratory, Graduate School of Agriculture, Hokkaido University, Kita 9

21 Nishi 9, Kita-ku, Sapporo, Hokkaido 060-8589, Japan

For review

22

23 **Abstract**

24 We evaluated the effect of chemical fertilizer and manure applications on N<sub>2</sub>O emission  
25 from a managed grassland by establishing three treatment plots of chemical N fertilizer  
26 (chemical fertilizer), manure combined chemical N fertilizer (manure), and no N fertilizer  
27 (control) at the Shizunai Experimental Livestock Farm in southern Hokkaido, Japan. N<sub>2</sub>O  
28 fluxes from soils were measured by a closed-chamber method from May 2005 to April  
29 2008. Soil denitrifying enzyme activity (DEA) in root-mat layer (0-2.5cm) and mineral  
30 soil layer (2.5-5cm) of each treatment plot was measured by an acetylene inhibition  
31 method after treatment with NO<sub>3</sub><sup>-</sup>-N addition, glucose addition, both NO<sub>3</sub><sup>-</sup>-N and glucose  
32 addition and neither NO<sub>3</sub><sup>-</sup>-N nor glucose addition, respectively. Annual N<sub>2</sub>O emission  
33 ranged from 0.6 4.9 kg N<sub>2</sub>O-N ha<sup>-1</sup>yr<sup>-1</sup>, with the highest observed in manure plot and  
34 lowest in control plot. Chemical fertilizer-induced emission factor (EF) (range: 0.85 -  
35 1.32%) was significantly higher than manure-induced EF (range 0.35 - 0.85%).  
36 Denitrification potential of soil horizons was measured with addition of both NO<sub>3</sub><sup>-</sup>-N and  
37 glucose, which was significantly higher in root-mat soil than that in mineral soil. Soil  
38 DEA in root-mat in NO<sub>3</sub><sup>-</sup>-N addition with and without addition of glucose had a  
39 significantly positive correlation with soil pH (P < 0.05). Soil pH was significantly

40 influenced by N source, which was significantly lower in chemical fertilizer plot than that  
 41 in control and manure plot. For a fixed quantity of available N, application of manure  
 42 could result in higher N<sub>2</sub>O emission compared to chemical fertilizer owing to high soil pH  
 43 values under manure application than under chemical fertilizer application.

44 **Key words:** chemical fertilizer, grassland, manure, N<sub>2</sub>O emission, soil DEA.

## 46 INTRODUCTION

47 N<sub>2</sub>O is one of the most important radiatively active trace gases in the atmosphere that  
 48 contributes at least 5% to the observed global warming at present (Myhre *et al.* 1998).

49 The atmospheric concentration of N<sub>2</sub>O has increased from a pre-industrial value of about  
 50 270 ppb to 319 ppb in 2005 and continues to increase as a result of human activities  
 51 (IPCC 2007). Agriculture as a whole (i.e. animal excreta, denitrification of leached  
 52 nitrate, etc.) contributes about 80% of the anthropogenic N<sub>2</sub>O emissions (Brown *et al.*  
 53 2001). Direct and indirect emissions from agricultural systems are now thought to  
 54 contribute approximately 6.2 Tg N<sub>2</sub>O-N yr<sup>-1</sup> to the total global source strength of 17.7 Tg  
 55 N<sub>2</sub>O-N yr<sup>-1</sup> (Kroeze *et al.* 1999). About 57% of the global atmospheric sources of N<sub>2</sub>O are  
 56 estimated to be related to emissions from soils (Mosier *et al.* 1998).

57 N<sub>2</sub>O is produced in soils primarily by microbial processes of nitrification and

**Deleted:** We evaluated the effect of chemical fertilizer and manure applications on N<sub>2</sub>O emission from a managed grassland by establishing three treatment plots of chemical fertilizer, manure, and control at the Shizunai Experimental Livestock Farm in southern Hokkaido, Japan. Seasonal N<sub>2</sub>O and NO fluxes were measured by a closed-chamber method at 4 to 6 replications in each treatment plot from May 2005 to April 2008. Soil samples were collected from a 0-5 cm top soil layer at three replications on each gas sampling date for measuring pH, NO<sub>3</sub><sup>-</sup>-N, NH<sub>4</sub><sup>+</sup>-N, and DOC. Soil samples were collected from the root-mat layer (0-2.5cm) and the mineral soil layer (2.5-5cm) of each treatment plot in April, June, and August 2007, which was followed by measuring soil denitrifying enzyme activity (DEA). The soil DEA was measured by an acetylene inhibition method under the four treatments with and without the addition of NO<sub>3</sub><sup>-</sup>-N and glucose. The cumulative N<sub>2</sub>O emission in control, chemical fertilizer, and manure plots ranged from 0.6 to 0.7, 1.4 to 3.0, and 2.1 to 4.9 kg N<sub>2</sub>O-N ha<sup>-1</sup>yr<sup>-1</sup>, respectively. The application of both

**Formatted:** Font: Times New Roman

**Deleted:** contributing

1  
2  
3  
4  
5 58 denitrification (Tiedje 1988; Conrad 1996). Nitrification is the biological oxidation of  
6  
7  
8 59 ammonium to nitrite or nitrate under aerobic conditions, while denitrification is the  
9  
10 60 reduction of nitrate to  $N_2O$  and  $N_2$  when the supply of oxygen is limited. Increasing soil N  
11  
12 61 availability as a result of increased N inputs by the application of chemical fertilizer and  
13  
14 62 manure and atmospheric deposition have greatly enhanced  $N_2O$  emissions from soils  
15  
16 63 (Kroeze *et al.* 1999) by influencing nitrifying and denitrifying enzyme activity. Chemical  
17  
18  
19  
20  
21 64 fertilizer and animal wastes are the two most important sources of direct  $N_2O$  emissions  
22  
23 65 from agricultural soils (Mosier *et al.* 1998). The default IPCC emission factor, i.e. the  
24  
25 66 percentage of applied N emitted as  $N_2O$ , is 1.0%, regardless of the fertilizer type (IPCC  
26  
27  
28  
29 67 2006). However, the type of N inputs to the fields may affect the  $N_2O$  emission rate in  
30  
31 68 different ways, leading to different patterns of  $N_2O$  emissions from inorganic and organic  
32  
33 69 N fertilizers. Addition of inorganic N increases  $N_2O$  emission through affecting the  
34  
35 70 process of nitrification and denitrification by increasing the available  $NH_4^+$ -N and  
36  
37  
38  
39 71  $NO_3^-$ -N substrates. Organic fertilizers with a high and easily mineralizable organic C  
40  
41 72 content stimulate microbial activity and thus  $N_2O$  emissions (Chadwick *et al.* 2000). The  
42  
43 73 application of chemical N fertilizer resulted in short-lived  $N_2O$  peaks (Dobbie and Smith  
44  
45 74 2003). However,  $N_2O$  losses from manure plots extended over a longer period of time and  
46  
47  
48  
49 75 were greater in magnitude than from chemical N fertilization (Jones *et al.* 2007). Higher  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Deleted: agricultural

Deleted: through affecting

Deleted: activities

Deleted: type of fertilizer

Deleted: the

Deleted: is reported to

Deleted: the

Deleted: by

Deleted: providing

Deleted: additional

Deleted: .

1  
2  
3  
4  
5 76 N<sub>2</sub>O fluxes from manure and sewage applications compared to that from chemical N  
6  
7  
8 77 fertilizers were also observed in other studies (Scott *et al.* 2000). High N<sub>2</sub>O fluxes from  
9  
10 78 manure treatments can be partly explained by the higher total N input than chemical  
11  
12 fertilizer treatments, providing more available N that was mineralized over a longer  
13  
14 79  
15  
16 80 period of time. Another reason for the increased N<sub>2</sub>O losses could be an addition of  
17  
18 81 organic C by the manures, which is known to stimulate denitrification. McTaggart *et al.*  
19  
20 82 (1997) reported that C supply from slurry spread onto grasslands stimulated N<sub>2</sub>O  
21  
22 83 production, resulting in a four times larger N<sub>2</sub>O loss compared to the application of  
23  
24 84 NH<sub>4</sub>NO<sub>3</sub> although the total rate of N application was similar in both treatments.  
25  
26  
27

Deleted: partly

Deleted: additional

28  
29 85 Grassland is an important ecosystem to support the production of herbivorous livestock  
30  
31 86 (Soussana *et al.* 2007). Application of chemical fertilizer and animal manure to  
32  
33 87 grasslands has been conducted to increase grass production, especially in developed  
34  
35 88 countries where grassland-based livestock production is important (Bouwman *et al.*  
36  
37 89 2002). However, N application to grasslands also poses a risk of N loss to the  
38  
39 90 environment in the form of N<sub>2</sub>O emission. The objective of this study is to clarify the  
40  
41 91 effect of fertilizer and manure application on N<sub>2</sub>O emission and to identify the factors  
42  
43  
44  
45  
46  
47 92 controlling N<sub>2</sub>O emission from the grassland.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Deleted: clarify

## 94 MATERIALS AND METHODS

## 95 Study site

96 This study was conducted on a managed grassland located at the Shizunai  
97 Experimental Livestock Farm, Field Science Center for Northern Biosphere of Hokkaido  
98 University in Southern Hokkaido, Japan (42°26'N, 142°29'E). The site is characterized  
99 by a humid continental climate with cold winters and cool summers. During 1979 to 2000,  
100 the mean annual precipitation and air temperature for this region were 1365 mm and  
101 7.9°C, respectively. The soil is derived from Tarumae (b) volcanic ash, and is classified as  
102 Thaptic Melanudands (Soil Survey Staff, 2006; Mollic Andosol (IUSS Working Group,  
103 WRB, 2006)). A layer of three cm thick root-mat was found on the top, and a 21 cm thick  
104 Ap-layer was found under the root-mat in a survey conducted in August 2004 (Shimizu *et*  
105 *al.* 2009). The C and N contents in the Ap-layer were 3.7% and 0.33%, respectively, and  
106 the C:N ratio was 11.1. Dominant grass species at this site were reed canary grass  
107 (*Phalaris arundinacea L.*) and foxtail grass (*Alopecurus pratensis L.*). The harvesting of  
108 grass was carried out twice a year (21<sup>st</sup> June and 11<sup>th</sup> August in 2005, 27<sup>th</sup> June and 23<sup>rd</sup>  
109 August in 2006 and 18<sup>th</sup> June and 18<sup>th</sup> August in 2007) in accordance with the local  
110 practice.

Formatted: Justified

Deleted: in

Formatted: Superscript

Formatted: Superscript



## 112 Experimental setup

113 Three experimental plots were set up on the study site; one for treatment with chemical  
114 fertilizer (chemical fertilizer plot), another with beef cattle manure and chemical fertilizer  
115 (manure plot), and the other with no N fertilizer or manure (control plot). Setting of the  
116 treatments was initiated in the spring of 2005. Eighteen subplots (5 m × 4 m) were  
117 established for the chemical fertilizer, manure, and control plots with six and four  
118 replications from May 2005 to April 2007 and from May 2007 to April 2008,  
119 respectively.

120 Table 1 shows the information on the date of application and the application rates of  
121 chemical fertilizer and manure. The N application rates in the chemical fertilizer plot  
122 were at the recommended level for this site on the basis of soil tests, and were 164 kg N  
123 ha<sup>-1</sup> year<sup>-1</sup> in 2005 and 184 kg N ha<sup>-1</sup> year<sup>-1</sup> in 2006 as ammonium sulfate and ammonium  
124 phosphate (Table 1). For learning the N<sub>2</sub>O emission from grassland soil under the local N  
125 fertilization level, N fertilization decreased to 74 kg N ha<sup>-1</sup> year<sup>-1</sup> in chemical fertilizer  
126 plot according to the recommends of Shizunai Experimental Livestock Farm staffs. The  
127 pH value of manure that used in 2005, 2006 and 2007 were 8.3, 8.8 and 9.1, respectively.

128 The manure application rates were the optimum rates used by farmers in the region, and  
129 were based on adequate amounts of potassium (K) application to the fields. Beef cattle

Deleted: ,

Deleted: in

Formatted: Font color: Auto

Formatted: Subscript

Formatted: Font color: Auto

1  
2  
3  
4  
5 130 manure with bedding litter (bark) was applied to the manure plot, and the application  
6  
7  
8 131 rates were 44 Mg FM ha<sup>-1</sup> (236 kg N ha<sup>-1</sup> and 5.8 Mg C ha<sup>-1</sup>) in May 2005, 43 Mg FM ha<sup>-1</sup>  
9  
10 132 (310 kg N ha<sup>-1</sup> and 6.0 Mg C ha<sup>-1</sup>) in May 2006, and 43 Mg FM ha<sup>-1</sup> (331 kg N ha<sup>-1</sup> and 7.7  
11  
12 133 Mg C ha<sup>-1</sup>) in May 2007 (Table 1). In the manure plot, the N supply rates from manure  
14  
15  
16 134 were estimated by multiplying the application rates by the N mineralization rate, and the  
17  
18 135 differences between the supply rates in manure and the application rates in the fertilizer  
19  
20  
21 136 plot were supplied by chemical fertilizer. The N mineralization rates were estimated  
22  
23 137 based on Uchida's model (Shiga *et al.* 1985) which was developed in Japan and were  
24  
25  
26 138 13.2%, 7.0% and 5.5%, respectively in the first, second and third years after application.  
27  
28  
29 139 The mineralization rates of P and K from the manure were estimated based on the  
30  
31 140 handbook of animal waste management and utilization in Hokkaido 2004 (Anon. 2004).  
32  
33  
34 141 The P mineralization rate was 20%, 10%, and 0% and the K mineralization rate was 70%,  
35  
36 142 10%, and 0% in the first year, second year, and third year after application, respectively.  
37  
38  
39 143

#### 144 **N<sub>2</sub>O and NO fluxes**

145 We defined the crop growing season as a 7-day moving average of daily air temperature  
146 above 5 °C and the non-growing season as the rest (Shimizu *et al.* 2009). The growing  
147 season was 215 days in 2005 (From 10<sup>th</sup> April 2005 to 10<sup>th</sup> November 2005), 218 days in

Deleted: nutrient

1  
2  
3  
4  
5 148 2006 (From 15<sup>th</sup> April 2006 to 18<sup>th</sup> November 2006) and 220 days in 2007 (From 13<sup>th</sup>  
6  
7  
8 149 April 2007 to 18<sup>th</sup> November 2007. N<sub>2</sub>O and NO fluxes from the soil to the atmosphere  
9  
10 150 were measured by the static closed chamber method on the control, fertilizer, and manure  
11  
12  
13 151 plots (Shimizu *et al.* 2009). The flux measurements were conducted in 2 - 28 day intervals  
14  
15  
16 152 during the crop growing season and 10 - 30 day intervals during the non growing season  
17  
18  
19 153 and between 8:00 and 11:00 h in each measuring day to minimize the effect of diurnal  
20  
21 154 temperature variation. The stainless steel chambers were 40 cm in diameter and 30 cm  
22  
23  
24 155 high in the chemical fertilizer and manure plots, and 20 cm in diameter and 25 cm high in  
25  
26 156 the control plots. The chambers were placed directly into the soil to a depth of about 3 cm,  
27  
28  
29 157 12 hours before the measurement of each subplot, and contained no aboveground biomass  
30  
31 158 in the chemical fertilizer, manure, and control plots. Before closing the chamber, a 250 ml  
32  
33  
34 159 gas sample from the headspace of each chamber was extracted into a Tedlar bag for NO  
35  
36  
37 160 analysis, and a 20 ml gas sample was injected into an evacuated vial (10 ml) for N<sub>2</sub>O  
38  
39 161 analysis. This measurement was regarded as time 0 min. After 20 min or 30 min under a  
40  
41  
42 162 closed-chamber condition, 250 ml of the headspace gas sample was extracted from each  
43  
44  
45 163 chamber into a bag, and 20 ml was injected into a vial. From these bag samples, NO gas  
46  
47  
48 164 concentrations were determined in a laboratory within 16 hours using a  
49  
50 165 Chemiluminescence N Oxide Analyzer (Model 265P, Kimoto Electric, Osaka, Japan).  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5 166 N<sub>2</sub>O gas concentrations were determined in a laboratory within 1 month using an ECD  
6  
7  
8 167 (Electron capture detector) gas chromatograph (model GC-14B, Shimadzu, Kyoto,  
9  
10 168 Japan) from the samples of vials.

11  
12  
13 169 Gas fluxes were calculated from the change in gas concentration in the chamber against  
14  
15  
16 170 closure time:

17  
18 171 
$$F = \rho \times h \times (\Delta c / \Delta t) \times [273 / (273 + T)]$$

19  
20  
21 172 where F is the gas flux ( $\mu\text{g N m}^{-2} \text{ h}^{-1}$  for N<sub>2</sub>O),  $\rho$  is the gas density (N<sub>2</sub>O-N =  $1.26 \times 10^9$   
22  
23 173  $\mu\text{g m}^{-3}$ ), h is the height of the chamber from the soil surface (m),  $\Delta c / \Delta t$  is the change in  
24  
25  
26 174 gas concentration inside the chamber during the sampling period ( $\text{m}^3 \text{ m}^{-3} \text{ h}^{-1}$ ), and T is the  
27  
28  
29 175 average air temperature during the sampling period ( $^{\circ}\text{C}$ ). A positive flux denotes the  
30  
31 176 emission from the soil, whereas a negative flux denotes the uptake from the atmosphere.

32  
33  
34 177 The cumulative gas flux was calculated as follow:

35  
36  
37 178 Cumulative gas flux = 
$$\sum_{i=1}^n (R_i \times 24 \times D_i)$$

38  
39  
40 179 where R<sub>i</sub> is the mean gas flux ( $\text{mg m}^{-2} \text{ hr}^{-1}$ ) of the two successive sampling dates, D<sub>i</sub> is the

41  
42  
43 180 number of days in the sampling interval, and n is the number of sampling times. The

44  
45  
46 181 cumulative period of 2005, 2006 and 2007 were calculated from 10<sup>th</sup> April 2005 to 14<sup>th</sup>

47  
48  
49 182 April, from 15<sup>th</sup> April 2006 to 18<sup>th</sup> April 2007 and from 19<sup>th</sup> April 2007 to 4<sup>th</sup> April 2008,

50  
51 183 respectively.

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

184 **Emission factor**

185 N<sub>2</sub>O Emission factor (EF) for chemical fertilizer and manure (kg N<sub>2</sub>O–N (kg N input)<sup>-1</sup>)

186 was calculated as follows:

187 Chemical fertilizer-induced EF = {[N<sub>2</sub>O emission (chemical fertilizer plot)] – [N<sub>2</sub>O  
188 emission (control plot)]} / [chemical fertilizer N application rate (chemical fertilizer  
189 plot)]

190 Manure-induced EF = {[N<sub>2</sub>O emission (manure plot)] – [chemical fertilizer N application  
191 rate (manure plot)] × chemical fertilizer-induced EF – [N<sub>2</sub>O emission (control plot)]} /  
192 [manure N application rate (manure plot)]

193 **Environmental variables**

194 Daily precipitation was obtained at the Sasayama AMeDAS (Automated Meteorological  
195 Data Acquisition System) station by the Japan Meteorological Agency. Air temperature  
196 and soil temperature at a 5 cm depth were measured at the same time with the flux  
197 measurements using a thermistor thermometer (CT220, CUSTOM, Tokyo, Japan), and  
198 soil moisture content at a 0 - 6 cm depth was measured using the Frequency Domain  
199 Reflectometry (FDR) method (DIK-311A, Daiki, Saitama, Japan). Soil core samples (14  
200 cm diameter, 13 cm height) were collected in April 2007, and calibration curves were  
201 made to calculate water-filled pore space (WFPS) from the FDR device reading (m<sup>3</sup> m<sup>-3</sup>)

**Deleted:** and daily maximum depths  
of snow were

**Deleted:**

**Deleted:** closest

1  
2  
3  
4  
5 202 and percent total porosity (Linn and Doran 1984). The percent total porosity was  
6  
7  
8 203 measured using a 100 ml soil core collected in April 2007 and was regarded as constant  
9  
10 204 throughout the study period because of no tillage.

#### 11 12 13 205 **Soil chemical analyses**

14  
15 206 Soil samples at a depth of 0 to 5 cm from ground surface were collected during a period  
16  
17  
18 207 from April to November at three replicates in all treatment plots. Within 48 hours of soil  
19  
20 208 sampling, soil samples were sieved through the 2 mm sieve and stones and roots were  
21  
22 209 removed. Soil samples were then immediately extracted in deionized water (1:5) and in 2  
23  
24 210 M KCl (1:10), and the extracts were stored at 4 °C until analysis for dissolved nutrients  
25  
26 211 after filtered through 0.2-µm membrane filters. Water-soluble organic carbon (DOC)  
27  
28 212 content in the deionized-water-extract solution was analyzed using a TOC analyzer (TOC  
29  
30 213 5000A, Shimadzu). The concentration of NO<sub>2</sub>-N and NO<sub>3</sub>-N in the  
31  
32 214 deionized-water-extract solution was analyzed by ion chromatography (Dionex QIC  
33  
34 215 Analyzer, Dionex Japan, Osaka, Japan). The concentration of NH<sub>4</sub><sup>+</sup>-N in the 2 M KCl  
35  
36 216 extracted solution was determined by the indophenol-blue method (UV mini 1240,  
37  
38 217 Shimadzu, Kyoto, Japan). Soil pH was measured in the deionized-water-extract solution  
39  
40  
41  
42 218 with a combined electrode pH meter (F-8 pH meter, Horiba, Japan).

#### 43 44 45 46 47 48 49 219 **Measurement of soil denitrifying enzyme activity (DEA)**

Deleted: -----Page Break-----

Deleted: under the root-mat

1  
2  
3  
4  
5 220 For measuring soil denitrifying enzyme activity (DEA), soil samples were taken from all  
6  
7  
8 221 treatment plots in the root-mat layer (0 - 2.5cm depth) and mineral soils layer (2.5 - 5cm  
9  
10 222 depth) with 3 replications in April, June, and August 2007. The root-mat soil samples  
11  
12  
13 223 were cut into small pieces with 1cm diameter, and stones or roots were removed from the  
14  
15 224 mineral soil samples by passing through the 2mm sieve within 48 h after sampling. Then  
16  
17  
18 225 we mixed three replicates of soil samples and kept in refrigerator at 4 °C until analysis.  
19  
20  
21 226 The DEA was determined by an acetylene block technique, which inhibits the final  
22  
23 227 conversion of N<sub>2</sub>O to N<sub>2</sub> gas (Tiedje, 1994). Soil samples were incubated under anaerobic  
24  
25  
26 228 condition at 25°C with a solution treated with 1) chloramphenicol (1g L<sup>-1</sup>) (Chl), 2) with  
27  
28 229 chloramphenicol (1g L<sup>-1</sup>) and NO<sub>3</sub><sup>-</sup>-N (200 mg N L<sup>-1</sup> as KNO<sub>3</sub>) (Chl+N), 3) with  
29  
30 230 chloramphenicol (1g L<sup>-1</sup>) and organic-C (2 g C L<sup>-1</sup> as glucose) (Chl+C), and 4) with  
31  
32 231 chloramphenicol (1g L<sup>-1</sup>), NO<sub>3</sub><sup>-</sup>-N (200 mg N L<sup>-1</sup> as KNO<sub>3</sub>) and organic-C (2 g C L<sup>-1</sup> as  
33  
34 232 glucose) (Chl+N+C). Fresh soil of 15g was placed into a 100 conical flask, and 15 ml  
35  
36  
37 233 treated solution was added to the flask. The flasks were evacuated and flushed four times  
38  
39  
40 234 with N<sub>2</sub> to ensure anaerobic conditions, and acetylene (C<sub>2</sub>H<sub>2</sub>) gas was added to a final  
41  
42  
43 235 concentration of 10% (10 kPa) in the headspace. The headspace gas was sampled by a  
44  
45  
46 236 syringe at 2 and 4 h and denitrification rates were calculated from the linear increment of  
47  
48  
49 237 N<sub>2</sub>O production against time. Denitrification potential of the soil horizons was measured

238 with the addition of both NO<sub>3</sub><sup>-</sup>-N and soluble C source as proposed by D'Haene *et al.*

239 (2003).

## 240 **Statistical analyses**

241 Analysis of variance (ANOVA) and Pearson correlation analysis were performed using

242 SPSS 13.0. Linear regression and other statistical analyses were carried out by using

243 Excel 2003. Two-way ANOVA and Tukey test were used to compare the mean difference

244 (P < 0.05) of a given variable between treatment plots and years. Three-way ANOVA and

245 Tukey test were used to compare the mean difference (P < 0.05) in soil N<sub>2</sub>O fluxes among

246 the seasons, treatment plots of field experiment, and years; and in soil DEA among the

247 treatment plots of field experiment, soil layers, and incubation treatments.

248

## 249 **RESULTS**

### 250 **Soil temperature and moisture**

251 Daily precipitation is shown in Fig. 1a. Annual precipitations were 1176 mm from the

252 mid April 2005 to the mid April 2006, 1047 mm from the mid April 2006 to the mid April

253 2007, and 879 mm from the mid April 2007 to the beginning of April 2008. These values

254 are smaller than that the mean annual precipitation (1365 ± 215 mm) from 1989 to 2000.

255 Soil moisture content expressed as WFPS in a 0 to 6 cm depth is shown in Fig. 1b. The

Deleted: ,

Deleted: plots

Deleted: These values are smaller than the long-term average (1365 ± 215 mm).¶



1  
2  
3  
4  
5 256 WFPS from April to November was influenced by precipitation and the low soil moisture  
6  
7  
8 257 was observed with low precipitation in August 2006 and June 2007. From December to  
9  
10 258 March, soil moisture was not observed because of soil freezing. In the winter of  
11  
12  
13 259 2006/2007, soil freezing began from beginning of December 2006, reaching the  
14  
15 260 maximum depth of 17.75 cm in 9<sup>th</sup> March 2007, and then thawed in early April in 2007.  
16  
17  
18 261 In the winter of 2007/2008, soil freezing also began from beginning of December 2007,  
19  
20 262 but reaching the maximum 29 cm in 12<sup>th</sup> March 2008, and then thawed in early April in  
21  
22  
23 263 2008. The soil freezing depth was not observed in the winter of 2005/2006.

24  
25  
26 264 Soil temperature at a 5 cm depth is shown in Fig. 1c. The soil temperature increased  
27  
28  
29 265 from April, reaching its maximum from July through August, and then decreased  
30  
31 266 gradually. The soil temperature was around 0 °C from December to March. There was no  
32  
33  
34 267 difference in soil temperature between the chemical fertilizer and manure plots, but soil  
35  
36 268 temperature was higher in the control plot than in the chemical fertilizer and manure plots  
37  
38  
39 269 (P < 0.05).

#### 270 **N<sub>2</sub>O fluxes**

271 The seasonal patterns of N<sub>2</sub>O fluxes were mainly driven by a seasonal variation as  
272 varieties of air and soil temperature which were higher in summer and lower in winter,  
273 and influenced by fertilization (Fig. 2a). N<sub>2</sub>O fluxes in the chemical fertilizer and manure

Deleted: ¶

Deleted: pattern of N<sub>2</sub>O fluxes was mainly driven by a seasonal variation

plots increased after application of manure or chemical fertilizer. These remained at a higher level than in the control plot until the beginning of September (Fig. 2a). Three-way ANOVA showed that there was a significant difference in N<sub>2</sub>O fluxes between the non-growing and growing seasons (P < 0.001) and between each treatment (P < 0.05). In the meantime, an interaction between the season and treatment in the N<sub>2</sub>O fluxes was observed (P < 0.05) (Table 2). In the growing season, the mean N<sub>2</sub>O fluxes in 2005, 2006, and 2007 were 12.3, 12.9, and 16.7 ug N<sub>2</sub>O-N m<sup>-2</sup>h<sup>-1</sup> for control plot, 85.3, 83.9, and 36.5 ug N<sub>2</sub>O-N m<sup>-2</sup>h<sup>-1</sup> for chemical fertilizer plot, and 101.9, 187.6, and 50.6 ug N<sub>2</sub>O-N m<sup>-2</sup>h<sup>-1</sup> for manure plot, respectively (Table 3). The N<sub>2</sub>O fluxes in the growing season were significantly higher in the chemical fertilizer and manure plots than that in the control plot (P < 0.01), but there was no significant difference between the chemical fertilizer and manure plots. In the non-growing season, the N<sub>2</sub>O fluxes were lower and stable with the mean values of 0.3, 2.6 and 4.7 ug N<sub>2</sub>O-N m<sup>-2</sup>h<sup>-1</sup> in the control plot, 2.7, 6.0 and 7.4 ug N<sub>2</sub>O-N m<sup>-2</sup>h<sup>-1</sup> in the chemical fertilizer plot, and 2.7, 3.6 and 3.5 ug N<sub>2</sub>O-N m<sup>-2</sup>h<sup>-1</sup> in the manure plot in 2005, 2006, and 2007, respectively (Table 3). There was no significant difference in the mean of N<sub>2</sub>O fluxes between each treatment plot.

The annual N<sub>2</sub>O emission in control, chemical fertilizer, and manure plots ranged from 0.6 to 0.7, 1.4 to 3.0 and 2.1 to 4.9 kg N<sub>2</sub>O-N ha<sup>-1</sup>yr<sup>-1</sup> during 2005 to 2007, respectively

**Deleted:** N<sub>2</sub>O fluxes in the chemical fertilizer and manure plots increased after the application of N fertilizer.

**Deleted:** 8

**Deleted:** 4

**Deleted:** cumulative

**Deleted:** in

**Deleted:** , 2006, and

1  
2  
3  
4  
5 292 (Table 4). Application of both chemical fertilizer and manure stimulated the annual  
6  
7  
8 293 cumulative N<sub>2</sub>O emissions, and the significantly highest annual cumulative N<sub>2</sub>O emission  
9  
10 294 was observed in the manure plot, followed by the chemical fertilizer plot. Application of  
11  
12  
13 295 chemical fertilizer contributed to 76.9, 79.2 and 47.2% of the total N<sub>2</sub>O emission from the  
14  
15 296 chemical fertilizer plot in 2005, 2006, and 2007, respectively. In the manure plot, N<sub>2</sub>O  
16  
17  
18 297 emission from the applied chemical fertilizer and manure contributed to 81.8, 87.4, and  
19  
20  
21 298 67.6% of the total N<sub>2</sub>O emission in 2005, 2006, and 2007, respectively. Chemical  
22  
23  
24 299 fertilizer-induced EF was 1.32, 1.30, and 0.85 % in 2005, 2006, and 2007, respectively.  
25  
26 300 Manure-induced EF was significantly lower than the fertilizer-induced EF (P < 0.001),  
27  
28  
29 301 and was 0.51, 0.85, and 0.35 % in 2005, 2006, and 2007, respectively (Table 5).

30  
31 302 NO fluxes showed a seasonal variation that similar with the seasonal pattern of N<sub>2</sub>O  
32  
33  
34 303 fluxes, which were higher in summer and lower in winter and also influenced by  
35  
36 304 fertilization (Fig. 2b). The NO fluxes ranged from -1.2 to 91.3 ug NO-N m<sup>-2</sup>h<sup>-1</sup> which was  
37  
38  
39 305 smaller than the N<sub>2</sub>O fluxes (-3.6 to 1290.7ug N<sub>2</sub>O-N m<sup>-2</sup>h<sup>-1</sup>). Large NO fluxes were  
40  
41  
42 306 observed mainly after the manure and chemical fertilizer application (Fig. 2b). Most of  
43  
44 307 the values of N<sub>2</sub>O/NO ratio were distributed from 1 to 100, and a significant positive  
45  
46  
47 308 correlation was found (P < 0.01) between the N<sub>2</sub>O/NO ratio and the N<sub>2</sub>O fluxes (Fig. 3).  
48  
49  
50 309

Deleted: variability in a

Deleted:

Deleted: seasonal variation and

Deleted: 249.0

Deleted: of fertilizer

Deleted:

Deleted: ly

310 **Soil chemical properties**

311 ~~During 2005 to 2007, soil pH in chemical fertilizer plot was obviously lower than that in~~  
 312 ~~control and manure plot (Fig.4a). Chemical fertilizer application could lead to the soil pH~~  
 313 ~~decreasing not only in chemical fertilizer plot but also in manure plot (Fig.4a). The mean~~  
 314 soil pH ~~during 2005 to 2007~~, in control, chemical fertilizer and manure plots were 5.2, 4.6,  
 315 and 5.1, respectively. The soil pH in the chemical fertilizer plot was significantly lower  
 316 than that in the manure and control plots ( $P < 0.001$ ). There was no significant difference  
 317 in soil pH between the manure and control plots.

318 The mean  $\text{NH}_4^+\text{-N}$  concentrations, ~~during 2005 to 2007~~, in control, chemical fertilizer,  
 319 and manure plots were 4.4, 23.7, and 17.5  $\text{mg kg}^{-1}$ , respectively. Soil  $\text{NH}_4^+\text{-N}$   
 320 concentration in chemical fertilizer and manure plots exhibited a wide range of 0.4 - 245  
 321  $\text{mg kg}^{-1}$  (Fig. 4b). In contrast, soil  $\text{NH}_4^+\text{-N}$  concentration in the control plot was stable,  
 322 and was always below 12  $\text{mg kg}^{-1}$ . ~~The pattern of soil  $\text{NH}_4^+\text{-N}$  concentration was not~~  
 323 ~~influenced by manure application only; however, it was influenced by the chemical~~  
 324 ~~fertilizer~~ application (Fig. 4b). Soil  $\text{NH}_4^+\text{-N}$  concentration in chemical fertilizer and  
 325 manure plots increased rapidly right after ~~chemical fertilizer~~ application, but then  
 326 decreased within a few ~~days (Fig. 4b)~~. In 2005 and 2006, peak concentrations of soil  
 327  $\text{NH}_4^+\text{-N}$  were ~~always~~ observed ~~in both~~ chemical fertilizer ~~and Manure plot after chemical~~

Deleted: From

Deleted: soil pH from the 0 - 5 cm top soil layer in control, chemical fertilizer and manure plots showed a similar seasonal pattern (Fig. 4a).

Formatted: Font color: Aqua

Deleted: However

Deleted: , t

Deleted: in 2005, 2006, and 2007,

Deleted:

Deleted: in 2005, 2006, and 2007

Deleted: T

Deleted: he pattern of soil  $\text{NH}_4^+\text{-N}$  concentration

Deleted: N

Deleted: the N

Deleted: days

Deleted: twice

1  
2  
3  
4  
5 328 fertilizer application. But in 2007, only one small peak was observed in the chemical  
6  
7  
8 329 fertilizer plot following the base fertilizer application in May. After chemical fertilizer  
9  
10 330 application in July 2007, soil  $\text{NH}_4^+$ -N concentration peak was not observed may caused  
11  
12  
13 331 by low precipitation and low soil moisture that limited the additional  $\text{NH}_4^+$ -N by chemical  
14  
15  
16 332 fertilizer application go into more deep soil layer by water dynamics before it was  
17  
18 333 assumed on the soil surface and in root mat through  $\text{NH}_4^+$ -N volatilization or microbial,  
19  
20  
21 334 chemical and physical reaction.

Deleted: and manure plots

22  
23 335 The mean  $\text{NO}_3^-$ -N concentrations, during 2005 to 2007, were 1.4, 2.7, and 2.2  $\text{mg kg}^{-1}$  in  
24  
25  
26 336 the control, chemical fertilizer, and manure plot, respectively. The pattern of soil  $\text{NO}_3^-$ -N  
27  
28  
29 337 concentration was also influenced by N application, and the peaks were observed slightly  
30  
31 338 later than those of soil  $\text{NH}_4^+$ -N concentration (Fig. 4c).

Deleted:

Deleted: in 2005, 2006, and 2007

32  
33 339 During the study period, the soil DOC concentration ranged from 48 to 121  $\text{mg kg}^{-1}$  in  
34  
35  
36 340 the control plot, from 23 to 116  $\text{mg kg}^{-1}$  in the chemical fertilizer plot, and from 43 to 199  
37  
38  
39 341  $\text{mg kg}^{-1}$  in the manure plot (Fig. 4d). The mean soil DOC concentrations in control,  
40  
41  
42 342 chemical fertilizer, and manure plots were 73.3, 59.4, and 97.8  $\text{mg kg}^{-1}$ , respectively. The  
43  
44  
45 343 soil DOC concentration in the manure plot was significantly higher than that in the  
46  
47  
48 344 control and chemical fertilizer plots ( $P < 0.01$ ), but application of chemical fertilizer had  
49  
50 345 no significant influence on the soil DOC concentration compared with that in the control

Deleted: 156

1  
2  
3  
4  
5 346 plot. Continuous 3 years manure application significantly increased the soil DOC  
6  
7  
8 347 concentration, which was significantly higher in 2007 than in 2005 and 2006 (P<0.01).  
9

10 348 Pearson correlation analysis showed that instantaneous N<sub>2</sub>O flux had a strong positive  
11  
12  
13 349 correlation with soil temperature (P < 0.01), soil NO<sub>3</sub><sup>-</sup>-N concentration (P < 0.01), and  
14  
15 350 soil NH<sub>4</sub><sup>+</sup>-N concentration (P < 0.01) (Table 6).  
16  
17

### 18 351 **Soil denitrifying enzyme activity (DEA)**

19  
20 352 Table 7 shows the result of DEA. Result of a 3-way ANOVA shows that there was a  
21  
22  
23 353 significant difference in soil DEA among soil layers (root-mat and mineral) (P < 0.001)  
24  
25  
26 354 and treatments (with and without NO<sub>3</sub> and glucose) (P < 0.001), but there was no  
27  
28  
29 355 significant difference in soil DEA among plots (control, chemical fertilizer, and manure)  
30  
31 356 (P = 0.058) (Table 8). However, there was a significant interaction between the soil layer  
32  
33  
34 357 and treatments (P < 0.001). In the root-mat layer, soil DEA was significantly increased by  
35  
36 358 the addition of NO<sub>3</sub>-N with (P < 0.001) or without (P < 0.001) the addition of glucose. But  
37  
38  
39 359 there was no significant effect on soil DEA by the addition of only glucose. There was no  
40  
41  
42 360 significant difference in soil DEA between the treatments Chl+N and Chl+N+C. On the  
43  
44  
45 361 other hand, there was no significant effect of single addition of NO<sub>3</sub>-N (Chl+N) or  
46  
47 362 glucose (Chl+C) on soil DEA in the mineral soil. However, a combination of NO<sub>3</sub>-N and  
48  
49  
50 363 glucose addition (Chl+N+C) increased the soil DEA significantly (P < 0.05). The soil  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5 364 DEA with the addition of both  $\text{NO}_3\text{-N}$  and glucose in the root-mat soil was significantly  
6  
7  
8 365 higher than that in the mineral soil ( $P < 0.001$ ). The soil DEA in the root-mat soil with the  
9  
10 366 addition of  $\text{NO}_3\text{-N}$  and both  $\text{NO}_3\text{-N}$  and glucose had a significantly positive correlation  
11  
12  
13 367 with soil pH ( $P < 0.05$ ) (Fig. 5).

14 368

17  
18 369 **DISCUSSION**19  
20  
21 370 **Seasonal pattern of  $\text{N}_2\text{O}$  emission**

22  
23 371 Soil  $\text{N}_2\text{O}$  fluxes were significantly higher in the growing season than that in the  
24  
25  
26 372 non-growing season. This is attributed to the high soil temperature (Table 6) in the  
27  
28  
29 373 growing season. Granli and Bøckman (1994) found an increased rate of  $\text{N}_2\text{O}$  production  
30  
31 374 with an increase in soil temperature up to 20-40 °C. High peaks of  $\text{N}_2\text{O}$  fluxes were  
32  
33  
34 375 usually observed in both chemical fertilizer and manure plots within a few weeks after the  
35  
36 376 application of manure or chemical fertilizer in our study (Fig. 2a). This can be attributed  
37  
38  
39 377 to the result of rapid increase in soil  $\text{NH}_4^+\text{-N}$  and  $\text{NO}_3^-\text{-N}$  concentrations immediately  
40  
41  
42 378 after the application of fertilizer, which decreased within a few days (Fig 4). It is well  
43  
44  
45 379 established that the rate of  $\text{N}_2\text{O}$  emission usually increases with an increase in soil  
46  
47 380 available N (Skiba and Smith 2000; Sehy *et al.* 2003). Several studies reported that  $\text{N}_2\text{O}$   
48  
49  
50 381 fluxes significantly increased after the application of N fertilizers. Mu *et al.* (2008)

Deleted: fertilizer

1  
2  
3  
4  
5 382 reported that N<sub>2</sub>O fluxes increased rapidly to higher emission levels in soils cultivated  
6  
7  
8 383 with wheat (from 242 to 433 μg N m<sup>-2</sup> h<sup>-1</sup>) and onion (from 47.2 to 157 μg N m<sup>-2</sup> h<sup>-1</sup>) after  
9  
10 384 N fertilization and that the fluxes lasted for about three weeks. Schils *et al.* (2008) also  
11  
12  
13 385 reported high N<sub>2</sub>O fluxes occurred in the first week after the application of chemical  
14  
15  
16 386 fertilizer or cattle slurry.

17  
18 387 The bacterial processes of nitrification and denitrification are the most important  
19  
20  
21 388 sources of N<sub>2</sub>O in soil (Granli and Bøckmann 1994). According to Davidson (1992) and  
22  
23 389 Skiba *et al.* (1993), nitrification produces more NO than N<sub>2</sub>O; conversely, denitrification  
24  
25  
26 390 produces more N<sub>2</sub>O than NO. The ratio of N<sub>2</sub>O-N/NO-N is the index of N<sub>2</sub>O production  
27  
28  
29 391 from nitrification or denitrification (Lipschultz *et al.* 1981). Lipschultz *et al.* (1981)  
30  
31 392 reported that the ratio of production of N<sub>2</sub>O-N / NO-N ranged from 0.2 to 1.0 in  
32  
33  
34 393 nitrification and 100 in denitrification. A significant positive correlation between the N<sub>2</sub>O  
35  
36  
37 394 fluxes and the ratio of N<sub>2</sub>O-N / NO-N and was found (Fig. 3) in our study (P < 0.01),  
38  
39 395 indicating that the high N<sub>2</sub>O emissions are primarily due to denitrification.

#### 40 41 42 396 **Cumulative N<sub>2</sub>O emission**

43  
44 397 Chemical fertilizer and animal wastes are [the](#) two most important sources of direct N<sub>2</sub>O  
45  
46  
47 398 emissions from agricultural soils (Mosier *et al.* 1998). Increasing soil N availability  
48  
49  
50 399 associated with application of N by chemical fertilizer and manure has greatly enhanced



1  
2  
3  
4  
5 400 N<sub>2</sub>O emissions from agricultural soils (Kroeze *et al.* 1999). Meng *et al.* (2005) found that  
6  
7  
8 401 chemical fertilizer and manure contributed to 74–82% of the total N<sub>2</sub>O emissions. Mori *et*  
9  
10 402 *al.* (2008) also reported the N<sub>2</sub>O emission predominantly derived from the manure and  
11  
12 403 the chemical fertilizer N application on a volcanic grassland soil in Nasu, Japan.  
13  
14  
15 404 Generally, emissions of N<sub>2</sub>O increase with an increase in N application rates (Granli and  
16  
17 405 Bøckman 1994; MacKenzie *et al.* 1997). In our study, the N<sub>2</sub>O emission from applied  
18  
19 406 chemical fertilizer and manure contributed to 77-85% of the total N<sub>2</sub>O emission in 2005  
20  
21 407 and 2006. The contribution of chemical fertilizer and manure to N<sub>2</sub>O emission in 2007  
22  
23 408 decreased to 47-65% due to the lower application rates than that in 2005 and 2006. The  
24  
25  
26  
27  
28 409 chemical fertilizer-induced EF ranged from 0.85 to 1.32%, which was comparable to the  
29  
30  
31 410 IPCC default value 1% (IPCC 2006), but was higher than that reported by Akiyama and  
32  
33 411 Tsuruta (2003) from the Japanese Andisols amended with chemical fertilizer (ranging  
34  
35 412 from 0.06% to 0.29%). The manure-induced EF of our study ranged from 0.35 to 0.85%,  
36  
37  
38 413 which was significantly lower than the chemical fertilizer-induced EF and the IPCC  
39  
40  
41 414 default value, but close to that reported by Akiyama and Tsuruta (2003), which was  
42  
43  
44 415 0.55%.

Deleted: average

Deleted: was 1.17%

Deleted: average

Deleted: was

Deleted: 57

#### 416 Soil DEA

417 The soil DEA with an addition of NO<sub>3</sub><sup>-</sup>-N and glucose in the root-mat soil was

Deleted: Denitrification potential of the soil horizons was measured with the addition of both NO<sub>3</sub><sup>-</sup>-N and soluble C source as proposed by D'Haene *et al.* (2003).

1  
2  
3  
4  
5 418 significantly higher than that in the mineral soil, indicating that the soil denitrification  
6  
7  
8 419 potential in the root-mat soil was significantly higher than that in the mineral soil.  
9  
10 420 Microbial activities in the surface soil are reported to be higher than in the deeper soil  
11  
12  
13 421 (Speir *et al.* 1984; Higashida and Takao, 1985). Parkin and Meisinger (1989) reported  
14  
15 422 that total viable bacteria and numbers of denitrifying bacteria were found to decrease  
16  
17  
18 423 exponentially with an increase in soil depth on a well-drained silt loam soil.

19  
20  
21 424 Soil DEA in the root-mat soil significantly increased by the addition of NO<sub>3</sub>-N with (P <  
22  
23 425 0.001) or without (P < 0.001) the addition of glucose, indicating that the availability of  
24  
25  
26 426 soil NO<sub>3</sub><sup>-</sup>-N could be the major limiting factor for soil DEA in our study grassland. In the  
27  
28 427 mineral soil, only addition NO<sub>3</sub>-N or glucose could not increase the soil DEA, but  
29  
30  
31 428 addition NO<sub>3</sub>-N and glucose together increased the soil DEA that means not only the  
32  
33  
34 429 NO<sub>3</sub>-N but also the carbon is the limited factor for soil DEA in mineral soil. The soil

35  
36 430 DEA in the root-mat soil with NO<sub>3</sub>-N addition and both NO<sub>3</sub>-N and glucose addition had  
37  
38  
39 431 a significantly positive correlation with soil pH (P < 0.05, Fig. 5). The soil pH is supposed  
40  
41  
42 432 to be a major variable of soil, controlling the microbial community in general and the  
43  
44 433 community of denitrifiers in particular (Simek and Hopkins 1999). Simek and Hopkins  
45  
46  
47 434 (1999) detected an optimum pH value for denitrification in soils as a range from 7 to 8.  
48  
49  
50 435 Simek and Cooper (2002) reported that both the overall rates of denitrification under field

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

1  
2  
3  
4  
5 436 conditions (i.e. the formation of  $N_2O$ ,  $N_2$  and  $NO$  and their subsequent emission) and  
6  
7  
8 437 DEA were influenced by soil pH, and they were less in acidic soils than in neutral or  
9  
10 438 slightly alkaline soils. Ellis *et al.* (1998) observed that the production of  $N_2O$  decreased  
11  
12  
13 439 with decreasing pH under anaerobic conditions through an incubation experiment. These  
14  
15  
16 440 results suggest that the highest  $N_2O$  emission in the manure plot in our study was resulted  
17  
18 441 from the soil DEA that could have been controlled by the soil pH.

19  
20  
21 442 The application of chemical fertilizer significantly decreased the soil pH in the chemical  
22  
23 443 fertilizer plot than in the control plot. However, the soil pH in manure plot was not  
24  
25  
26 444 significantly different with control plot. That maybe because higher pH of manure  
27  
28  
29 445 (8.3-9.1) input decrease the effect of chemical fertilizer application on the soil pH. Soil  
30  
31 446 acidity is controlled by the amount of  $H^+$  and  $Al^{3+}$  which is either contained in or  
32  
33  
34 447 generated by the soil and soil components. According to Kirikae *et al.* (2001),  
35  
36  
37 448 nitrification is a source of  $H^+$  through two nitrification pathways of  $NH_4^+$  origin and  
38  
39 449 organic N origin. They reported that the ratio of  $H^+$  to  $NO_3^-$  was 2 in the pathway of  $NH_4^+$   
40  
41  
42 450 origin and 1 in the pathway of organic N origin. In the meanwhile,  $NO_3^-$  uptake by  
43  
44  
45 451 vegetation was the sink of  $H^+$ . Therefore, the organic N has less effect on  $H^+$  production  
46  
47 452 than  $NH_4^+$ -N. On the other hand, application of manure increased the value of cation  
48  
49  
50 453 exchange capacity (CEC) compared to that of chemical fertilizer (Bulluck *et al.* 2002).

Deleted: manure application had no significant effect on t

454 The soils with a high CEC have a greater capacity to contain or generate sources of acidity.

455 The soil pH is higher in soils with manure than that with chemical fertilizer as reported by

456 several studies (Bulluck *et al.* 2002 and Gil *et al.* 2008).

### 457 **Conclusions**

458 The application of both chemical fertilizer and manure to grassland stimulated the annual

459 N<sub>2</sub>O emission. The chemical fertilizer-induced EF (range: 0,85 - 1,32%) was

460 significantly higher than the manure-induced EF (range 0.35 - 0.85%). However, annual

461 N<sub>2</sub>O emission was significantly higher in the manure plot than that in the chemical

462 fertilizer plot. The soil DEA in the NO<sub>3</sub>-abundant root-mat layer significantly decreased

463 with a decrease in soil pH. Moreover, application of chemical fertilizer could

464 significantly decrease soil pH, but the manure application had no significant effect on soil

465 pH. Therefore, for the a fixed quantity of available N, application of manure could result

466 in higher N<sub>2</sub>O emission compared to chemical fertilizer owing to high soil pH values

467 under manure application than under chemical fertilizer application.

468

### 469 **ACKNOWLEDGMENTS**

470 We would like to thank the technical staff of Shizunai Livestock Farm for their help in the

471 field measurements. This study was partly supported by a research grant provided by the

Deleted: cumulative

Deleted: 89

Deleted: 33

Deleted: similar

Deleted: application of manure could result in higher N<sub>2</sub>O emission compared to chemical fertilizer owing to high pH values in manures.¶

1  
2  
3  
4  
5 472 Project entitled 'Establishment of good practices to mitigate Greenhouse Gas emissions  
6  
7  
8 473 from Japanese grasslands' funded by Racing and Livestock Association.

9  
10 474

11  
12  
13 475 **REFERENCES**

14  
15 476 Akiyama H, Tsuruta H 2003: Effect of organic matter application on N<sub>2</sub>O, NO, and NO<sub>2</sub>  
16  
17  
18 477 fluxes from an Andisol field. *Global Biogeochem. Cycles*, **17**, 1-16.

19  
20 478 Anon., 2004: Handbook of animal waste management and utilization in Hokkaido.

21  
22  
23 479 Hokkaido Prefectural Experiment Stations and Hokkaido Animal Research Center,  
24  
25  
26 480 Sapporo, 64-67 (in Japanese).

27  
28 481 Bouwman AF, Boumans LJM, Batjes NH 2002: Modeling global annual N<sub>2</sub>O and NO  
29  
30  
31 482 emissions from fertilized fields. *Global Biogeochem. Cycles*, **16**, 1080.

32  
33  
34 483 Brown CD, Fryer CJ, Walker A 2001: Influence of topsoil till and soil moisture status on  
35  
36 484 losses of pesticide to drains from a heavy clay soil. *Pest Management Science*, **57**,  
37  
38  
39 485 1127-1134.

40  
41 486 Bulluck III LR, Brosius M, Evanylo GK, Ristaino JB 2002: Organic and synthetic  
42  
43  
44 487 fertility amendments influence soil microbial, physical and chemical properties on  
45  
46  
47 488 organic and conventional farms. *Applied Soil Ecology*, **19**, 147-160.

- 1  
2  
3  
4  
5 489 Chadwick DR, Pain BF, Brookman SKE 2000: Nitrous oxide and methane emissions  
6  
7  
8 490 following application of animal manures to grassland. *J. Environ. Qual.*, **29**, 277-287.  
9  
10 491 Conrad R 1996: Soil microorganisms as controllers of atmospheric trace gases (H<sub>2</sub>, CO,  
11  
12 492 CH<sub>4</sub>, OCS, N<sub>2</sub>O, and NO). *Microbiol. Rev.*, **60**, 609 - 640.  
13  
14  
15 493 D'Haene K, Moreels E, Neve S, Daguilar BC, Boeckx P, Hofman G, Van Cleemput O  
16  
17  
18 494 2003: Soil properties influencing the denitrification potential of Flemish agricultural  
19  
20 495 soils. *Biol. Fertil. Soils.*, **38**, 358-366.  
21  
22  
23 496 Davidson EA 1992: Source of nitric oxide and nitrous oxide following wetting of dry soil.  
24  
25  
26 497 *Soil Sci. Soc. Am. J.*, **56**, 95-102.  
27  
28  
29 498 Dobbie, KE, Smith KA 2003: Nitrous oxide emission factors for agricultural soil in Great  
30  
31 499 Brita: the impact of soil water-filled pore space and other controlling variables.  
32  
33 500 *Global Change Biol.*, **9**, 204-218.  
34  
35  
36 501 Ellis S, Howe MT, Goulding WT, Muggleston M.A, Dendooven, L 1998: Carbon and  
37  
38 502 nitrogen dynamics in a grassland soil with varying pH: effect of pH on the  
39  
40 503 denitrification potential and dynamics of the reduction enzymes. *Soil Biol. Biochem.*,  
41  
42 504 **30**, 359-367.  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5 505 Gil MV, Carballo MT and Calvo LF 2008: Fertilization of maize with compost from  
6  
7  
8 506 cattle manure supplemented with additional mineral nutrients. *Waste Management*,  
9  
10 507 **28**, 1432-1440.  
11  
12 508 Granli T and Bøckman OC 1994: Nitrous oxide from agriculture. *Nor. J. Agric. Sci.*,  
13  
14  
15 509 *Suppl.*, **12**, 7-128.  
16  
17  
18 510 Higashida S, and Takao K 1985: Seasonal fluctuation patterns of microbial numbers in  
19  
20  
21 511 the surface of a grassland. *Soil Sci. plant Nutr.*, **31**, 113-121.  
22  
23 512 IPCC 2006: Guidelines for National Greenhouse Gas Inventories, prepared by the  
24  
25  
26 513 National Greenhouse Gas Inventories Programme. Intergovernmental Panel on  
27  
28  
29 514 Climate Change.  
30  
31 515 IPCC 2007: Climate changes 2007: The Physical Science Basis. Cambridge University  
32  
33  
34 516 Press, Cambridge.  
35  
36 517 IUSS Working Group WRB 2006: World Reference Base for Soil Resources 2006.  
37  
38  
39 518 World Soil Resources Reports 103, FAO, Rome.  
40  
41 519 Jones SK, Rees RM, Skiba UM, Ball BC 2007: Influence of organic and mineral N  
42  
43  
44 520 fertiliser on N<sub>2</sub>O fluxes from a temperate grassland. *Agr. Ecosyst. Environ.*, **121**,  
45  
46  
47 521 74–83.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5 522 Kirikae M, Hatano R, Shibata H, Tanaka Y 2001: Analysis of proton generation and  
6  
7  
8 523 consumption of forest surface soils in Hokkaido, northern Japan. *Water, Air and Soil*  
9  
10 524 *Pollution*, **130**, 697-702.
- 11  
12  
13 525 Kroeze C, Mosier AR, Bouwman AF 1999: Closing the global N<sub>2</sub>O budget: A  
14  
15 526 retrospective analysis 1500 - 1994. *Global Biogeochem. Cycles*, **13**, 1-8.
- 17  
18 527 Linn DM, Doran JW 1984: Effect of water-filled pore-space on carbon-dioxide and  
19  
20 528 nitrous-oxide production in tilled and nontilled soils. *Soil Sci. Soc. Am. J.*, **48**,  
21  
22 529 1267-1272.
- 23  
24  
25  
26 530 Lipschultz F, Zafiriou OC, Wofsy SC, McElroy MB, Valois FW, Watson SW 1981:  
27  
28 531 Production of NO and N<sub>2</sub>O by soil nitrifying bacteria. *Nature*, **294**, 641-643.
- 30  
31 532 MacKenzie AF, Fan MX, Cadrin F 1997: Nitrous oxide emission as affected by tillage,  
32  
33 533 corn-soybean-alfalfa rotations and nitrogen fertilization. *Can. J. Soil Sci.*, **77**,  
34  
35 534 145-152.
- 36  
37  
38  
39 535 McTaggart IP, Douglas JT, Clayton H, Smith KA 1997: Nitrous oxide emissions from  
40  
41 536 slurry and mineral nitrogen applied to grassland. In: Jarvis, S.C., Pain, B.F. (Eds.),  
42  
43 537 Gaseous Nitrogen Emission from Grasslands. CAB International, Wallingford, UK,  
44  
45 538 209-210.
- 46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



- 1  
2  
3  
4  
5 539 Meng L, Ding W, Cai Z 2005: Long-term application of organic manure and nitrogen  
6  
7  
8 540 fertilizer on N<sub>2</sub>O emissions, soil quality and crop production in a sandy loam soil. *Soil*  
9  
10 541 *Biol. Biochem.*, **37**, 2037-2045  
11  
12  
13 542 Mogge B, Kaiser EA, Munch JC 1999: Nitrous oxide emissions and denitrification  
14  
15 543 N-losses from agricultural soils in the Bornhoved Lake region: influence of organic  
16  
17  
18 544 fertilizers and land-use. *Soil Biol. Biochem.*, **31**, 1245-1252.  
19  
20  
21 545 Mori A, Hojito M, Shimizu M, Matsuura S, Miyaji T, Hatano R 2008: N<sub>2</sub>O and CH<sub>4</sub>  
22  
23 546 fluxes from a volcanic grassland soil in Nasu, Japan: Comparison between manure plus  
24  
25  
26 547 fertilizer plot and fertilizer-only plot. *Soil Sci. Plant Nutr.*, **54**, 606-617.  
27  
28  
29 548 Mosier A, Kroeze C, Nevison C, Oenema O, Seitsinger S, van Cleemput O 1998: Closing  
30  
31 549 the global N<sub>2</sub>O budget: nitrous oxide emissions through the agricultural nitrogen  
32  
33  
34 550 cycle. *Nutr. Cycling Agroecosyst.*, **52**, 225-248.  
35  
36  
37 551 Mu Z, Kimura SD, Toma Y, Hatano R 2008: Nitrous oxide fluxes from upland soils in  
38  
39 552 central Hokkaido, Japan. *J. Environ. Qual.*, **20**, 1312-1322.  
40  
41  
42 553 Myhre G, Highwood EK, Shine KP, Stordal F 1998: New estimates of radiative forcing  
43  
44 554 due to well mixed greenhouse gases. *Geophys. Res. Lett.*, **25**, 2715-2718.  
45  
46  
47 555 Parkin TB, Meisinger JJ 1989: Denitrification below the crop rooting zone as influenced  
48  
49 556 by surface tillage. *J. Environ. Qual.*, **18**, 12-16.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5 557 Schils RLM, van Groenigen JW, Velthof GL, Kuikman PJ 2008: Nitrous oxide emissions  
6  
7  
8 558 from multiple combined applications of fertiliser and cattle slurry to grassland. *Plant*  
9  
10 559 *and Soil*, **310**, 89-101.
- 11  
12  
13 560 Scott A, Ball BC, Crichton IJ, Aitken MN 2000: Nitrous oxide and carbon dioxide  
14  
15 561 emissions from grassland amended with sewage sludge. *Soil Use Manage.*, **16**,  
16  
17 562 36-41.
- 18  
19  
20  
21 563 Sehy U, Ruser R, Munch JC 2003: Nitrous oxide fluxes from maize fields: relationship to  
22  
23 564 yield, site-specific fertilization, and soil conditions. *Agric. Ecosyst. Environ.*, **99**,  
24  
25 565 97-111.
- 26  
27  
28 566 Shiga H, Ohyama N, Maeda K, Suzuki M 1985: An evaluation of different organic  
29  
30 567 materials based on their decomposition pattern in paddy soils. *Res. Bull. Natl. Agric.*  
31  
32 *Res. Cetr.*, **5**, 1-19 (in Japanese with English summary).  
33  
34 568
- 35  
36 569 Shimizu M, Marutani S, Desyatkin AR, Jin T, Hata H, Hatano R 2009: The effect of  
37  
38  
39 570 manure application on carbon dynamics and budgets in a managed grassland of  
40  
41 571 Southern Hokkaido, Japan. *Agric. Ecosyst. Environ.*, **130**, 31-40.
- 42  
43  
44 572 Simek M, Cooper JE 2002: The influence of soil pH on denitrification-progress towards  
45  
46 573 the understanding of this interaction over the last 50 years. *European Journal of Soil*  
47  
48 574 *Science*, **53**, 345-354.

Formatted: Font: Italic

Deleted: Title:

Formatted: Font: Italic

- 1  
2  
3  
4  
5 575 Simek M, Hopkins DW 1999: Regulation of potential denitrification by soil pH in  
6  
7  
8 576 long-term fertilized arable soils. *Biol. Fertil. Soils*, **30**, 41-47.  
9  
10 577 Skiba U, Smith K.A 2000: The control of nitrous oxide emissions from agricultural and  
11  
12 578 natural soils. *Chemosphere Global Change Sci.*, **2**, 379-386.  
13  
14  
15 579 Skiba U, Smith KA, Fowler D 1993: Nitrification and denitrification as source of nitric  
16  
17 580 oxide and nitrous oxide in a sandy loam soils. *Soil Biol. Biochem.*, **25**, 1527-1536.  
18  
19  
20 581 Soil Survey staff 2006: Keys to Soil Taxonomy, 10th ed. USDA-Natural Resources  
21  
22 582 Conservation Service, Washington DC.  
23  
24  
25 583 Soussana JF, Allard V, Pilegaard K 2007: Full accounting of the greenhouse gas (CO<sub>2</sub>,  
26  
27 584 N<sub>2</sub>O, CH<sub>4</sub>) budget of nine European grassland sites. *Agric. Ecosyst. Environ.*, **121**,  
28  
29 585 121-134.  
30  
31  
32 586 Speir T, Ross DJ, Orchard VA 1984: Spatial Variability of biochemical properties in a  
33  
34 587 taxonomically-uniform soil under grazed pasture. *Soil Biol Biochem.*, **16**, 153-160.  
35  
36  
37 588 Tiedje JM 1988: Ecology of denitrification and dissimilatory nitrate reduction to  
38  
39 589 ammonium. In: Zehnder, A.J.B., (Ed.), *Biology of Anaerobic Microorganisms*,  
40  
41 590 Wiley, New York, 179-244.  
42  
43  
44 591 Tiedje JM, 1994: Denitrifiers. *Methods of Soil Analysis, Part 2. Microbiological and*  
45  
46  
47  
48  
49  
50 592 *Biochemical Properties*. Soil Sci. Soc. Amer., Madison, Wisconsin, 245-267.

Deleted: .

Deleted: .

Deleted: .

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

593

For review

594

595 **Figure captions**

596 **Figure 1** Seasonal patterns of precipitation (a), WFPS at 6 cm depth (b), and soil  
 597 temperature at 5 cm depth (c). Data of WFPS and soil temperature represent means  $\pm$   
 598 SD (n = 4 to 6).

599 **Figure 2** Seasonal patterns of soil N<sub>2</sub>O fluxes (a) and soil NO fluxes (b). Data represent  
 600 means  $\pm$  SD (n = 4 to 6). Full arrow indicates date of the chemical fertilizer  
 601 application, and dotted arrow indicates date of manure application. The growing  
 602 season was 215 days in 2005 (From 10<sup>th</sup> April 2005 to 10<sup>th</sup> November 2005), 218  
 603 days in 2006 (From 15<sup>th</sup> April 2006 to 18<sup>th</sup> November 2006) and 220 days in 2007  
 604 (From 13<sup>th</sup> April 2007 to 18<sup>th</sup> November 2007).

**Deleted:** The arrows indicate the date of fertilizer application.

605 **Figure 3** Relationships between N<sub>2</sub>O fluxes and the ratio of N<sub>2</sub>O to NO.

606 **Figure 4** Seasonal patterns of Soil pH (a), soil NH<sub>4</sub><sup>+</sup>-N (b), soil NO<sub>3</sub><sup>-</sup>-N (c), and soil DOC  
 607 (d) at a depth of 0 - 5 cm. Data represent means  $\pm$  SD (n = 3). Full arrow indicates  
 608 date of the chemical fertilizer application, and dotted arrow indicates date of manure  
 609 application. The arrows indicate the date of fertilizer application.

610 **Figure 5** Relationship between the soil DEA and the soil pH in root-mat soil.

Page 4: [1] Deleted

Jin

11/25/2009 6:49:00 PM

We evaluated the effect of chemical fertilizer and manure applications on N<sub>2</sub>O emission from a managed grassland by establishing three treatment plots of chemical fertilizer, manure, and control at the Shizunai Experimental Livestock Farm in southern Hokkaido, Japan. Seasonal N<sub>2</sub>O and NO fluxes were measured by a closed-chamber method at 4 to 6 replications in each treatment plot from May 2005 to April 2008. Soil samples were collected from a 0-5 cm top soil layer at three replications on each gas sampling date for measuring pH, NO<sub>3</sub><sup>-</sup>-N, NH<sub>4</sub><sup>+</sup>-N, and DOC. Soil samples were collected from the root-mat layer (0-2.5cm) and the mineral soil layer (2.5-5cm) of each treatment plot in April, June, and August 2007, which was followed by measuring soil denitrifying enzyme activity (DEA). The soil DEA was measured by an acetylene inhibition method under the four treatments with and without the addition of NO<sub>3</sub><sup>-</sup>-N and glucose. The cumulative N<sub>2</sub>O emission in control, chemical fertilizer, and manure plots ranged from 0.6 to 0.7, 1.4 to 3.0, and 2.1 to 4.9 kg N<sub>2</sub>O-N ha<sup>-1</sup>yr<sup>-1</sup>, respectively. The application of both chemical fertilizer and manure significantly increased the cumulative N<sub>2</sub>O emission, and the highest cumulative N<sub>2</sub>O emission was observed in the manure plot. N<sub>2</sub>O fluxes were positively correlated with the N<sub>2</sub>O:NO ratio (P < 0.01), indicating high N<sub>2</sub>O fluxes resulting from the increased denitrification activities. The denitrification potential of the soil horizons was significantly higher in the root-mat soil than that in the mineral soil. The soil DEA in the root-mat soil in NO<sub>3</sub><sup>-</sup>-N addition with and without the addition of glucose had a significantly positive correlation with soil pH (P < 0.05). The soil pH was significantly influenced by N fertilization, which was significantly lower in the chemical fertilizer plot than that in the control and manure plots. For the similar quantity of available N,

1  
2  
3 application of manure could result in higher N<sub>2</sub>O emission compared to chemical fertilizer  
4  
5  
6 owing to high pH values in manures.  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For review

**Table 1** The applied date and the application rates of chemical fertilizer and manure during the study period.

| Treatment           | Date      | Fertilizer type                  | Application rates (kg ha <sup>-1</sup> ) |     |                               |                  |
|---------------------|-----------|----------------------------------|--|-----|-------------------------------|------------------|
|                     |           |                                  | C  | N   | P <sub>2</sub> O <sub>5</sub> | K <sub>2</sub> O |
| Control             | 2005/5/11 | Chemical fertilizer <sup>a</sup> | 0  | 0   | 0                             | 0                |
|                     | 2005/7/4  | Chemical fertilizer <sup>a</sup> | 0  | 0   | 0                             | 0                |
|                     | 2006/5/9  | Chemical fertilizer <sup>a</sup> | 0  | 0   | 0                             | 0                |
|                     | 2006/7/10 | Chemical fertilizer <sup>a</sup> | 0  | 0   | 0                             | 0                |
|                     | 2007/5/12 | Chemical fertilizer <sup>a</sup> | 0  | 0   | 14                            | 73               |
|                     | 2007/7/5  | Chemical fertilizer <sup>a</sup> | 0  | 0   | 7                             | 37               |
| Chemical fertilizer | 2005/5/11 | Chemical fertilizer <sup>a</sup> | 0  | 103 | 23                            | 168              |
|                     | 2005/7/4  | Chemical fertilizer <sup>a</sup> | 0  | 61  | 23                            | 97               |
|                     | 2006/5/9  | Chemical fertilizer <sup>a</sup> | 0  | 124 | 50                            | 177              |
|                     | 2006/7/10 | Chemical fertilizer <sup>a</sup> | 0  | 59  | 18                            | 97               |
|                     | 2007/5/12 | Chemical fertilizer <sup>a</sup> | 0  | 49  | 14                            | 73               |
|                     | 2007/7/5  | Chemical fertilizer <sup>a</sup> | 0  | 25  | 7                             | 37               |
| Manure              | 2005/5/11 | Manure <sup>b</sup>              | 5833                                     | 236 | 191                           | 266              |
|                     | 2005/7/4  | Chemical fertilizer <sup>a</sup> | 0  | 133 | 7                             | 70               |
|                     | 2006/5/9  | Manure <sup>b</sup>              | 5958                                     | 310 | 212                           | 167              |
|                     | 2006/5/9  | Chemical fertilizer <sup>a</sup> | 0  | 71  | 0                             | 33               |
|                     | 2006/7/10 | Chemical fertilizer <sup>a</sup> | 0  | 59  | 6                             | 97               |
|                     | 2007/5/12 | Manure <sup>b</sup>              | 7714                                     | 331 | 342                           | 336              |
|                     | 2007/7/5  | Chemical fertilizer <sup>a</sup> | 0  | 21  | 0                             | 0                |

<sup>a</sup> Chemical fertilizer is comprised of ammonium sulfate, ammonium phosphate, potassium sulfate, and potassium magnesium sulfate.

<sup>b</sup> Beef cattle manure with bedding litter was applied in the manure plot.



**Table 2** ANOVA results for mean N<sub>2</sub>O fluxes

| Source                    | df  | Mean Square | F      | P value |
|---------------------------|-----|-------------|--------|---------|
| Season                    | 1   | 179,974.04  | 17.521 | 0       |
| Year                      | 2   | 13,990.75   | 1.362  | 0.258   |
| Treatment                 | 2   | 39,922.53   | 3.887  | 0.022   |
| Season * Year             | 2   | 15,307.66   | 1.49   | 0.228   |
| Season * Treatment        | 2   | 38,549.47   | 3.753  | 0.025   |
| Year * Treatment          | 4   | 7,659.25    | 0.746  | 0.562   |
| Season * Year * Treatment | 4   | 7,454.57    | 0.726  | 0.575   |
| Error                     | 210 | 10,271.71   |        |         |

Deleted: Sig

For review

**Table 3** Mean N<sub>2</sub>O fluxes from the control, chemical fertilizer, and manure plots.

| Season             | Treatment           | Mean N <sub>2</sub> O fluxes (ug N <sub>2</sub> O-N m <sup>-2</sup> h <sup>-1</sup> ) |               |             |
|--------------------|---------------------|---|---------------|-------------|
|                    |                     | 2005  | 2006          | 2007        |
| Growing season     | Control             | 12.3 (12.7)   | 12.9 (15.23)  | 16.7 (27.6) |
|                    | Chemical fertilizer | 85.3 (94.1)   | 83.9 (66.3)   | 36.5 (53.2) |
|                    | Manure              | 101.9 (102.1)   | 187.6 (301.6) | 50.6 (48.9) |
| Non-growing season | Control             | 0.3 (0.6)   | 2.6 (6.0)     | 4.7 (3.1)   |
|                    | Chemical fertilizer | 2.7 (2.3)   | 6.0 (11.4)    | 7.4 (13.7)  |
|                    | Manure              | 2.7 (3.9)   | 3.6 (5.1)     | 3.5 (8.0)   |

Data represent means (SD), n=4 to 6. We defined the crop growing season as a 7-day moving average of daily air temperature above 5 °C and the non-growing season as the rest (Shimizu *et al.* 2009). The growing season was 215 days in 2005 (From 10<sup>th</sup> April 2005 to 10<sup>th</sup> November 2005), 218 days in 2006 (From 15<sup>th</sup> April 2006 to 18<sup>th</sup> November 2006) and 220 days in 2007 (From 13<sup>th</sup> April 2007 to 18<sup>th</sup> November 2007).

Formatted: Font: 12 pt

Deleted: .

Formatted: Font: 12 pt

**Table 4** Cumulative N<sub>2</sub>O emissions from the control, chemical fertilizer, and manure

plots

Deleted: .

| Treatment                        | Cumulative N <sub>2</sub> O emissions (kg N <sub>2</sub> O-N ha <sup>-1</sup> yr <sup>-1</sup> ) |           |           |
|----------------------------------|--|-----------|-----------|
|                                  | 2005   | 2006      | 2007      |
| Control <sup>a</sup>             | 0.7 (0.4)  | 0.6 (0.3) | 0.7 (0.5) |
| Chemical fertilizer <sup>b</sup> | 2.8 (0.7)  | 3.0 (0.8) | 1.4 (0.5) |
| Manure <sup>c</sup>              | 3.6 (1.2)  | 4.9 (2.8) | 2.1 (0.6) |

Deleted: .

Data represent means (SD), n=4 to 6, and different small letters denote the significant difference at the 0.05 level between each treatment.

For review

**Table 5** N<sub>2</sub>O Emission factor for chemical fertilizer and manure (kg N<sub>2</sub>O–N (kg N input)<sup>-1</sup>)

|                     | N <sub>2</sub> O emission factor (%) |             |             |
|---------------------|--------------------------------------|-------------|-------------|
|                     | 2005                                 | 2006        | 2007        |
| Chemical fertilizer | 1.32 (0.43)                          | 1.30 (0.44) | 0.85 (0.97) |
| Manure              | 0.51 (0.42)                          | 0.85 (0.89) | 0.35 (0.23) |

Data represent means (SD), n=4 to 6.

For review

**Table 6** Relationships (Pearson correlation coefficient, r) between instantaneous N<sub>2</sub>O fluxes and environmental factors using the whole data

|                              | N <sub>2</sub> O | Soil T.   | WFPS     | pH        | NO <sub>3</sub> <sup>-</sup> | NH <sub>4</sub> <sup>+</sup> | DOC |
|------------------------------|------------------|-----------|----------|-----------|------------------------------|------------------------------|-----|
| N <sub>2</sub> O             | 1                |           |          |           |                              |                              |     |
| Soil T.                      | 0.330 **         | 1         |          |           |                              |                              |     |
| WFPS                         | 0.020            | -0.288 ** | 1        |           |                              |                              |     |
| pH                           | -0.117           | -0.275 ** | 0.378 ** | 1         |                              |                              |     |
| NO <sub>3</sub> <sup>-</sup> | 0.307 **         | 0.378 **  | -0.061   | -0.201 *  | 1                            |                              |     |
| NH <sub>4</sub> <sup>+</sup> | 0.329 **         | 0.156 *   | -0.048   | -0.273 ** | 0.380 **                     | 1                            |     |
| DOC                          | -0.104           | -0.173 *  | -0.204 * | 0.341 **  | -0.282 **                    | -0.040                       | 1   |

\*p<0.05; \*\*p<0.01

Deleted: .

Deleted: 1 ... [1]

Deleted: 1 ... [2]

Deleted: 1 ... [3]

Deleted: 1 ... [4]

Deleted: 1 ... [5]

Deleted: 1 ... [6]

Deleted: 1

For review

**Table 7** Denitrification enzyme activity (DEA) of soil samples from three treatment plots of control, chemical fertilizer, and manure.

| Date      | Soil layer | Treatment | DEA (mgN <sub>2</sub> O-N kg <sup>-1</sup> h <sup>-1</sup> ) |        |                          |         |             |        |
|-----------|------------|-----------|--|--------|--------------------------|---------|-------------|--------|
|           |            |           | Control plot   |        | Chemical fertilizer plot |         | Manure plot |        |
| 2007/4/29 | Root-mat   | Chl       | 0.04   | (0.04) | 0.04                     | (0.01)  | 0.04        | (0.02) |
|           |            | Chl+N     | 9.99   | (1.07) | 3.66                     | (0.50)  | 16.30       | (2.98) |
|           |            | Chl+C     | 0.06   | (0.01) | 0.07                     | (0.02)  | 0.14        | (0.07) |
|           | Mineral    | Chl+N+C   | 10.98  | (6.71) | 3.08                     | (0.49)  | 13.56       | (3.16) |
|           |            | Chl       | 2.68   | (3.84) | 0.16                     | (0.11)  | 0.58        | (0.34) |
|           |            | Chl+N     | 0.52   | (0.64) | 0.16                     | (0.03)  | 1.56        | (0.25) |
|           |            | Chl+C     | 0.02   | (0.12) | 0.22                     | (0.01)  | 0.07        | (0.14) |
| 2007/6/11 | Root-mat   | Chl+N+C   | 1.00   | (0.20) | 2.21                     | (1.50)  | 1.97        | (0.36) |
|           |            | Chl       | 0.15   | (0.02) | 0.01                     | (0.00)  | 0.10        | (0.05) |
|           |            | Chl+N     | 14.05  | (5.03) | 2.63                     | (0.68)  | 15.05       | (2.28) |
|           | Mineral    | Chl+C     | 0.08   | (0.02) | 0.02                     | (0.01)  | 0.13        | (0.03) |
|           |            | Chl+N+C   | 9.47   | (0.33) | 4.66                     | (0.40)  | 17.35       | (0.81) |
|           |            | Chl       | 0.09   | (0.03) | 0.07                     | (0.02)  | 0.71        | (0.09) |
|           |            | Chl+N     | 1.43   | (0.00) | 0.35                     | (0.10)  | 1.24        | (0.55) |
| 2007/8/20 | Root-mat   | Chl+C     | 0.03   | (0.00) | 0.04                     | (0.01)  | 0.05        | (0.14) |
|           |            | Chl+N+C   | 2.14   | (0.21) | 0.54                     | (0.15)  | 1.71        | (2.15) |
|           |            | Chl       | 0.12   | (0.05) | 0.20                     | (0.06)  | 0.04        | (0.01) |
|           | Mineral    | Chl+N     | 27.84  | (3.87) | 9.80                     | (0.18)  | 13.58       | (1.83) |
|           |            | Chl+C     | 0.13   | (0.06) | 0.14                     | (0.05)  | 0.03        | (0.02) |
|           |            | Chl+N+C   | 21.33  | (0.23) | 35.19                    | (37.29) | 19.91       | (1.73) |
|           |            | Chl       | 1.14   | (0.31) | 0.76                     | (0.37)  | 3.01        | (0.20) |
| Mineral   | Chl+N      | 1.34      | (0.17)   | 0.55   | (0.09)                   | 2.49    | (1.83)      |        |
|           | Chl+C      | 0.37      | (0.03)   | 0.64   | (0.33)                   | 1.03    | (1.22)      |        |
|           | Chl+N+C    | 2.92      | (0.23)   | 0.83   | (0.33)                   | 7.65    | (0.87)      |        |

Data represent means (SD), n=3.

**Table 8** ANOVA results for soil denitrification enzyme activity (DEA)

| Source                        | df | Mean Square | F      | <u>P value</u> |
|-------------------------------|----|-------------|--------|----------------|
| Plot                          | 2  | 94,796,698  | 3.351  | 0.058          |
| Soil layer                    | 1  | 991,237,940 | 35.040 | 0.000          |
| Treatment                     | 3  | 832,401,859 | 29.426 | 0.000          |
| Plot * Soil layer             | 2  | 9,030,352   | 0.319  | 0.390          |
| Plot * Treatment              | 6  | 53,775,775  | 1.901  | 0.174          |
| Soil layer * Treatment        | 3  | 451,061,076 | 15.945 | 0.000          |
| Plot * Soil layer * Treatment | 5  | 25,358,833  | 0.896  | 0.258          |
| Error                         | 37 | 28,288,420  |        |                |

Deleted: .

Deleted: Sig.

For review

|    |                     |     |                       |
|----|---------------------|-----|-----------------------|
| 1  |                     |     |                       |
| 2  |                     |     |                       |
| 3  | Page 6: [1] Deleted | Jin | 11/24/2009 7:25:00 AM |
| 4  | 1                   |     |                       |
| 5  |                     |     |                       |
| 6  | Page 6: [1] Deleted | Jin | 11/24/2009 7:24:00 AM |
| 7  | 0.330 **            |     |                       |
| 8  |                     |     |                       |
| 9  | Page 6: [1] Deleted | Jin | 11/24/2009 7:24:00 AM |
| 10 | 0.020               |     |                       |
| 11 |                     |     |                       |
| 12 | Page 6: [1] Deleted | Jin | 11/24/2009 7:24:00 AM |
| 13 | -0.117              |     |                       |
| 14 |                     |     |                       |
| 15 | Page 6: [1] Deleted | Jin | 11/24/2009 7:24:00 AM |
| 16 | 0.307 **            |     |                       |
| 17 |                     |     |                       |
| 18 | Page 6: [1] Deleted | Jin | 11/24/2009 7:24:00 AM |
| 19 | 0.329 **            |     |                       |
| 20 |                     |     |                       |
| 21 | Page 6: [1] Deleted | Jin | 11/24/2009 7:24:00 AM |
| 22 | -0.104              |     |                       |
| 23 |                     |     |                       |
| 24 | Page 6: [2] Deleted | Jin | 11/24/2009 7:24:00 AM |
| 25 | 1                   |     |                       |
| 26 |                     |     |                       |
| 27 | Page 6: [2] Deleted | Jin | 11/24/2009 7:24:00 AM |
| 28 | -0.288 **           |     |                       |
| 29 |                     |     |                       |
| 30 | Page 6: [2] Deleted | Jin | 11/24/2009 7:24:00 AM |
| 31 | -0.275              |     |                       |
| 32 |                     |     |                       |
| 33 | Page 6: [2] Deleted | Jin | 11/24/2009 7:24:00 AM |
| 34 | 0.378 **            |     |                       |
| 35 |                     |     |                       |
| 36 | Page 6: [2] Deleted | Jin | 11/24/2009 7:24:00 AM |
| 37 | 0.156 *             |     |                       |
| 38 |                     |     |                       |
| 39 | Page 6: [2] Deleted | Jin | 11/24/2009 7:24:00 AM |
| 40 | -0.173 *            |     |                       |
| 41 |                     |     |                       |
| 42 | Page 6: [3] Deleted | Jin | 11/24/2009 7:25:00 AM |
| 43 | 1                   |     |                       |
| 44 |                     |     |                       |
| 45 | Page 6: [3] Deleted | Jin | 11/24/2009 7:24:00 AM |
| 46 | 0.378 **            |     |                       |
| 47 |                     |     |                       |
| 48 | Page 6: [3] Deleted | Jin | 11/24/2009 7:24:00 AM |
| 49 | -0.061              |     |                       |
| 50 |                     |     |                       |
| 51 | Page 6: [3] Deleted | Jin | 11/24/2009 7:24:00 AM |
| 52 | -0.048              |     |                       |
| 53 |                     |     |                       |
| 54 | Page 6: [3] Deleted | Jin | 11/24/2009 7:24:00 AM |
| 55 | -0.204 *            |     |                       |
| 56 |                     |     |                       |
| 57 | Page 6: [4] Deleted | Jin | 11/24/2009 7:25:00 AM |
| 58 | 1                   |     |                       |
| 59 |                     |     |                       |
| 60 | Page 6: [4] Deleted | Jin | 11/24/2009 7:25:00 AM |
|    | -0.201 *            |     |                       |
|    |                     |     |                       |
|    | Page 6: [4] Deleted | Jin | 11/24/2009 7:25:00 AM |



1  
2  
3 -0.273 \*\*

4  
5 **Page 6: [4] Deleted** **Jin** **11/24/2009 7:25:00 AM**

6 0.341 \*\*

7  
8 **Page 6: [5] Deleted** **Jin** **11/24/2009 7:25:00 AM**

9 1

10 **Page 6: [5] Deleted** **Jin** **11/24/2009 7:25:00 AM**

11 0.380 \*\*

12  
13 **Page 6: [5] Deleted** **Jin** **11/24/2009 7:25:00 AM**

14 -0.282 \*\*

15  
16 **Page 6: [6] Deleted** **Jin** **11/24/2009 7:25:00 AM**


17 1

18 **Page 6: [6] Deleted** **Jin** **11/24/2009 7:25:00 AM**


19 -0.040


20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For review



Follow the 'Figure captions' in the manuscript.





1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

