

FR 8400 B 14

BEHAVIOR OF THE WILSON PARAMETER IN U(1) LATTICE GAUGE THEORY  
WITH LONG RANGE GAUGE INVARIANT INTERACTIONS

Jean RUIZ \*

Centre de Physique Théorique  
C.N.R.S. - Luminy - Case 907  
13288 MARSEILLE CEDEX 9  
France

ABSTRACT :

The U(1) lattice gauge model with fermions can be expressed after integration over the fermionic variables as a "long range gauge model" : the effective action is a sum over all possible gauge field loops with corresponding weight factors. Different behaviors of the Wilson parameter are shown according to the hypothesis on the weight factors.

APRIL 1983

**CNRS** - CPT-83/P.1500

\* Phymat - Département de Mathématique - Université de Toulon et du Var.

## 1. INTRODUCTION

The purpose of this paper is to study the behavior of the Wilson parameter [1] in U(1) lattice gauge theory with long range gauge invariant interactions occurring in particular in lattice gauge theories with fermions [2]. These theories have been intensively studied analytically and recently also numerically by the Monte-Carlo method. The usual groups considered in a gauge invariant field theory are U(1), SU(N). One way to study the models consists in doing the "integration out" over the fermionic variables proposed by Matthews and Salam [3], [4], [5]; this "integration out" leads to an effective action which can be expressed as a sum over all possible gauge field loops affected with weight factors [2]. In the U(1) case the result is simple. For example in two space-time dimension and for Suskind fermions [6], the lattice fermionic action coupled to a gauge field is given by (see [7]):

$$S = S_F + S_G$$

$$\begin{aligned} (1) \quad S_F &\equiv (\bar{\Psi}, G(u) \Psi) \\ &\equiv \sum_{i,j} \left\{ \bar{\Psi}_{ij} U_{ij,i+1}^x \Psi_{i+1j} - \bar{\Psi}_{ij} U_{ij,i-1}^x \Psi_{i-1j} \right. \\ &\quad - i (-1)^{i+j} (\bar{\Psi}_{ij} U_{ij,i,j+1}^y \Psi_{i,j+1} - \bar{\Psi}_{ij} U_{ij,i,j-1}^y \Psi_{i,j-1}) \\ &\quad \left. + m \bar{\Psi}_{ij} \Psi_{ij} \right\} \end{aligned}$$

$\Psi$  and  $\bar{\Psi}$  are Grassman variables representing the fermion field. The couple  $(i,j)$  of integers represents the sites of the lattice. The one component variable  $\Psi_{ij}$  with  $i+j$  even or odd can be taken to represent respectively the field  $\psi^+$  or  $\psi^-$ .  $U_{ij,i+1}^x$ ,  $U_{ij,i,j+1}^y$  are the gauge field variables belonging to U(1) and indexed by links. They verify  $U_{a,b} = \bar{U}_{b,a}$ .  $S_G$  is the usual Wilson's lattice action.

$$(2) \quad S_G = \beta \sum_P \text{Re} [\text{tr} U(\rho)]$$

$P$  represents an elementary square (plaquette) of the lattice and  $U(\rho)$

is the product of the link variables associated to the plaquette  $P$ . To integrate out over the Grassman variables one uses the well known formulae (see [5])

$$\int d\psi d\bar{\psi} \exp(\bar{\psi} Q \psi) = \det Q$$

Expanding  $\det G(u) = \exp \text{Tr} \log G(u)$  by random walk techniques [8], [9], one obtains an effective action of the form

$$(3) \quad S_{\text{eff}} = \sum_{\gamma} \int_{\gamma} (m) R_{\epsilon} [t \cdot U_{\gamma}]$$

where  $U_{\gamma}$  is the product of the link variables associated to the closed path  $\gamma$ . The corresponding weight factors  $\int_{\gamma} (m)$  depend on  $m$  and on  $N$ :  $\int_{\gamma} (m) = \mathcal{E}(N)^{|\gamma|} m^{-|\gamma|}$ ,  $|\gamma|$  representing the length of the path and  $\mathcal{E}(N) = \pm 1$  according to the geometry of  $N$ . For "naive" fermions the result is similar.

The purpose of this paper is to study the behavior of the Wilson parameter: for this kind of action according to different hypothesis on the interaction  $\int_{\gamma} (m)$  in particular the interaction obtained from the Matthew-Salam expansion. The pure lattice gauge theory with action given by (2) is known to have a linear confinement in two dimension [10] a logarithmic confinement in three dimension [11] and is not confining at low temperature in four dimension [12]. We shall show that if the interaction does not decrease sufficiently with  $|\gamma|$  the model can have a confining behavior at all temperature: this occurs for ferromagnetic interactions, where  $\int_{\gamma} \geq 0$  for all  $\gamma$ . In the converse case we show that if the interaction decreases rapidly enough with  $|\gamma|$  then the model has a confining behavior at all temperature in dimensions two and three. These results are stated precisely in Section II, the proofs are given in Section III.

## II. DEFINITIONS AND RESULTS

We consider an infinite  $d$ -dimensional hypercubic lattice of unit spacing  $\Lambda \cong \mathbb{Z}^d$  ( $d \geq 2$ ). The basic objects on the lattice are the sites  $x \equiv \{x^1, x^2, \dots, x^d\} \in \mathbb{Z}^d$ , the links  $\langle x, x' \rangle$  where  $x$  and  $x'$  are nearest neighbours and the plaquettes  $p$  (elementary squares).

A walk on the lattice is an ordered set of oriented links

$$\omega \equiv \{ \langle x_1, x_2 \rangle, \langle x_2, x_3 \rangle, \dots, \langle x_{n-1}, x_n \rangle \}$$

A closed walk is a walk such that  $x_n = x_1$ . We divide the set of closed walks into equivalent classes by letting  $\omega_1, \omega_2$  be equivalent whenever  $\omega_1, \omega_2$  have the same links and the order of the links in  $\omega_2$  is a cyclic permutation of the order of the links in  $\omega_1$ . We call the equivalent classes "loops" and denote by  $\Lambda(\mathcal{L})$  the set of the loops.

To a loop  $\mathcal{L}$  we associate a loop  $\gamma(\mathcal{L})$  obtained from  $\mathcal{L}$  by eliminating two by two the terms  $\langle x_n, x_{n+1} \rangle, \langle x_n, x_{n+1} \rangle$  such that:  $x_n = x_{n+1}$  and  $x_{n+1} = x_n$ . We denote by  $\Lambda(\gamma)$  the set of these loops.  $|\mathcal{L}|$  (resp.  $|\gamma|$ ) denotes the number of links of  $\mathcal{L}$  (resp.  $\gamma$ ).

A connected surface  $S$  is a connected set of plaquettes.  $|S|$  denotes the number of surface of  $S$  and  $\Lambda(S)$  the set of connected surfaces.

Let  $\mathcal{L}$  be the set of links of  $\Lambda$ . To each link  $\ell = \langle x, x' \rangle$  of  $\mathcal{L}$  we associate a random variable  $H(\ell)$  with value in  $[-\pi, \pi]$  and such that  $H(x, x') = -H(x', x)$ . We denote by  $H_{\mathcal{L}}$  the sum of the link variables of the loop  $\mathcal{L}$  and by  $B_S$  the sum over the plaquettes  $p$  of  $S$  of  $B(p)$  where  $B(p) = A \partial_p$ ,  $\partial$  being the boundary operator.

We now consider the following actions

$$(4) \quad H_{\Lambda}^1 = - \sum_{\gamma \in \Lambda(N)} J_{\gamma} \cos A_{\gamma}$$

$$(5) \quad H_{\Lambda}^2 = - \sum_{S \in \Lambda(S)} K_S \cos B_S$$

where  $J_{\gamma}$  and  $B_S$  are real parameters.

Remark :  $H^1$  and  $H^2$  can be rewritten as

$$(6) \quad H_{\Lambda} = - \sum_{\gamma \in \Lambda(\Gamma)} I_{\gamma} \cos H_{\gamma}$$

with

$$I_{\gamma} = \sum_{\substack{\gamma' \in \Lambda(N) \\ \gamma' \text{ is a loop} \\ \text{associated to } \gamma}} J_{\gamma'} \quad \text{for } H^1$$

$$I_{\gamma} = \sum_{\substack{S \in \Lambda(S) \\ \partial S = \gamma}} K_S \quad \text{for } H^2$$

The Wilson parameter is given by

$$(7) \quad W_{\gamma}(c) = \langle e^{i A_{\gamma}} \rangle (\beta) = Z^{-1}(\beta) \int_{\mathcal{L}} \prod_{\ell \in \mathcal{L}} \frac{dA(\ell)}{2\pi} e^{i A_{\gamma} - \beta H_{\Lambda}}$$

$$Z^{-1}(\beta) = \int_{\mathcal{L}} \prod_{\ell \in \mathcal{L}} \frac{dA(\ell)}{2\pi} e^{-\beta H}$$

where  $dA/2\pi$  is the invariant measure on  $S(U)$ . The formulae (7) are to be interpreted as the thermodynamic limit  $N \rightarrow \infty$  of the corresponding finite volume quantities  $\langle e^{i A_{\gamma}} \rangle_{\Lambda'}(\beta)$  defined by the same expressions but with links restricted to a finite box  $\Lambda'$ . Let  $C$  be a rectangular loop of sides of length  $L$  and  $T$ , for pure gauge model given by (2) we consider  $E(L) = \lim_{T \rightarrow \infty} -\frac{1}{T} \log W_C(L)$  as the energy between static quarks separated by a distance  $L$ .

We denote by  $n(\ell)$  the number of loops of length  $\ell$  containing a given link. It is known that  $n(\ell) \leq (2d)^\ell$ . If  $N(s)$  denotes the number of connected surfaces of area  $s$  containing a given plaquettes then  $N(s) \leq \mu_d^s$ , where  $\mu_d$  is a positive number depending on the dimension  $d$  of the lattice. This follows by drawing the graphs whose edges connect the centers of the plaquettes containing a same link and by using the following fact : on every connected graph there is a path that passes through every edge at most twice [13].

We will now consider the following conditions.

Condition 1 : at large  $|V|$ ,  $J_V \sim |V| e^{-\nu_1 |V|}$  with  $\nu_1 > \log 2d$

Condition 2 : at large  $|V|$ ,  $J_V \sim |V| e^{-\nu_2 |V| \log |V|}$  with  $\nu_2 > 0$

Condition 3 : at large  $|S|$ ,  $K_S \sim e^{-\nu_3 |S|}$  with  $\nu_3 > \log \nu_d$

The condition 3 implies that  $I_Y$  decreases as  $\exp\{-cste \text{ minimal area with boundary } Y\}$ .

The conditions 1, 2, 3 imply the existence of the thermodynamic limit and give sufficient conditions of the Matthew-Salam expansion. The condition  $\nu_1 > 2d$  is a sufficient condition for the existence of the Matthew Salam expansion.

### Theorem 1

Let  $C$  be any loop. Consider the action given by (4) and assume that  $J_V$  verifies the condition 1, then :

a)  $\langle e^{iA_C} \rangle (\beta) \leq e^{-k_1 |Y(C)|}$  for any positive  $\beta$   
 $k_1$  is a positive constant and at large  $\beta$ ,  $k_1 \sim k'_1/\beta$  ( $k'_1$  being a positive constant).

b) If moreover :  $J_V \geq 0$  for all  $V$  then,

$$\beta_{1/2} |Y(C)| e^{-\nu_2 |Y(C)|} \leq \langle e^{iA_C} \rangle (\beta)$$

Theorem 2.

Let  $C$  be a rectangular loop of sides of length  $L$  and  $T$ . Consider the action given by (4) and assume that  $J_{\Psi}$  verifies the condition 2, then for any positive  $\beta$

- a) if  $d=2$   $\langle e^{iAc} \rangle (\beta) \leq e^{-k_2 T (\log L + \cot \alpha)}$   
 b) if  $d=3$   $\langle e^{iAc} \rangle (\beta) \leq e^{-k_3 T (\log L + \cot \alpha)}$   
 c) if  $d=4$   $\langle e^{iAc} \rangle (\beta) \leq e^{-k_4 (T+L)}$

$k_2$ ,  $k_3$  and  $k_4$  are positive constants and at large  $\beta$   $k_i \sim k'_i / \beta$ ,  $k'_i$  being positive constants.

d) if moreover :  $J_{\Psi} \geq 0$  for all  $\Psi$  then

$$\beta |T+L| e^{-2\mu_2 |T+L| \log |T+L|} \leq \langle e^{iHc} \rangle (\beta)$$

Theorem 3

Let  $C'$  be a rectangular loop of sides of length  $L$  and  $T$ . Consider the action given by (5) and assume that  $K_S$  verifies the condition 3. Then for any positive  $\beta$ ,

- a) if  $d=2$   $\langle e^{iAc} \rangle (\beta) \leq e^{-k_5 T L}$   
 b) if  $d=3$   $\langle e^{iAc} \rangle (\beta) \leq e^{-k_6 (\log L + \cot \alpha)}$   
 c) if  $d=4$   $\langle e^{iAc} \rangle (\beta) \leq e^{-k_7 (T+L)}$

$k_5$ ,  $k_6$  and  $k_7$  are positive constants and at large  $\beta$ ,  $k_i \sim k'_i / \beta$ ,  $k'_i$  being positive constants

d) if moreover :  $K_S \geq 0$  for all  $S$  then

$$\beta^{1/2} e^{-\mu_2 T \cdot L} \leq \langle e^{iAc} \rangle (\beta)$$

REMARKS

We can see that the upper bounds obtained in Theorem 1 for  $d = 4$ , in part b and c of Theorem 2 and in part a, b, c of Theorem 3 are of the same kind than those obtained for the  $U(1)$  pure lattice gauge theory with action given by (2).

If the interaction is ferromagnetic and in the 4-dimensional case one can obtain better lower bounds ( $\exp\{-c_1(\Gamma+L)\}$ ) than those obtained under the conditions 2 and 3 by using Ginzburg inequality [14] and Guth's lower bound [12].

The inequality a of Theorem 1 can be applied to the lattice gauge theory with fermions since the weight factors are given by  $E(\Gamma)/\Gamma^{|\Gamma|}$ . Nevertheless the lower bounds are only obtained in the ferromagnetic case and cannot be applied to this theory.



### III. PROOF OF THEOREMS

In the proof of upper bounds the idea consists in a comparison with Gaussian process. So we first use the method of complex translation of Mac Bryan and Spencer [15]. Our starting point is the following estimate due to Mac Bryan and Spencer (see also Glimm and Jaffe [11] for Gauge model).

#### Lemma 1

Let  $\{a(p)\}_{p \in \Lambda}$  be some configuration of links. Then

$$a) \langle e^{iA_c} \rangle(\beta) \leq \exp\{-a_c\} \exp\left\{\beta \sum_{\Gamma \in \Lambda(\Gamma)} J_\Gamma (ch a_\Gamma - 1)\right\}$$

$$b) \langle e^{iA_c} \rangle(\beta) \leq \exp\{-a_c\} \exp\left\{\beta \sum_{S \in \Lambda(S)} K_S (ch b_S - 1)\right\}$$

where  $b_S = \sum_{p \in S} b(p)$ ,  $b(p) = a_{op}$

We refer the reader to [15], [11] for the proof of this lemma.

For the proof of the lower bounds one uses Ginibre's inequality [14], [16]. In terms of gauge model it can be rewritten as follows :

$$(8) \quad \langle \cos H_\Gamma \rangle_{\mathcal{F}} \leq \langle \cos A_\Gamma \rangle_{\mathcal{F}} \quad \text{if } |J_p| \leq J_\Gamma \text{ for all } \Gamma$$

#### III.1 Proofs of the Lower Bounds in Theorems 1, 2, 3

In formula (7), let  $J_p = 0$  for all  $\Gamma$  excepted for  $\Gamma = \gamma(c)$ . Then by using inequality (8), we obtain if the interaction is ferromagnetic

$$(9) \quad \langle e^{iA_c} \rangle(\beta) \geq \frac{\int_{-\pi}^{\pi} \frac{dA(p)}{2\pi} e^{iA_c} e^{\beta J_\gamma(c) \cos A_\gamma(c)}}{\int_{-\pi}^{\pi} \frac{dA(p)}{2\pi} e^{\beta J_\gamma(c) \cos A_\gamma(c)}}$$

The right hand side of inequality (9) is equal to  $\frac{I_1(\beta J_\gamma(c))}{I_0(\beta J_\gamma(c))}$  where

$I_k(x)$  is the modified Bessel function.

Then one can show that

$$\frac{I_0(\beta J_{\gamma(c)})}{I_0(\beta J_{\gamma(c)})} \geq \beta^{1/2} J_{\gamma(c)}$$

According to the different hypothesis on  $J_{\gamma}$  we obtain the statement b of Theorem 1 and the statement d of Theorem 2. The statement c of Theorem 3 is obtained in the same way.

### III.2 Proof part a) of Theorem 1 and part c) of Theorem 2

Let  $C$  be an oriented loop. We consider a configuration  $\{a(\ell)\}_{\ell \in \mathcal{L}}$  verifying the following condition.

$$(10) \begin{cases} a(\ell) = \frac{1}{\beta k} & \text{for all } \ell \text{ in } C, \ell \text{ is oriented in the sense of } C \\ a(\ell) = 0 & \text{if } \ell \notin C \end{cases}$$

$k$  is a positive constant chosen later.

Let  $\ell$  be some link such that  $\gamma(c)$  contains  $\ell$ . By using part a) of Lemma 1 we obtain

$$\langle e^{iA_c} \rangle(\beta) \leq \exp \left\{ -\frac{|\gamma(c)|}{\beta k} \right\} \exp \left\{ \beta |\gamma(c)| \sum_{\substack{\gamma \in \Lambda(\beta) \\ \gamma \supset \ell}} J_{\gamma}(c a_{\gamma-1}) \right\}$$

$$\text{Let } P = \sum_{\substack{\gamma \in \Lambda(\beta) \\ \gamma \supset \ell}} J_{\gamma}(c a_{\gamma-1})$$

For  $\beta k$  large enough (we take  $\beta > \beta_0$  with  $\beta_0 \gg \frac{1}{k}$ ) we can write

$$P = \sum_{\substack{\gamma: |\gamma| < \beta k \\ \gamma \supset \ell}} J_{\gamma}(c a_{\gamma-1}) + \sum_{\substack{\gamma: |\gamma| \geq \beta k \\ \gamma \supset \ell}} J_{\gamma}(c a_{\gamma-1})$$

Since  $|a_{\gamma}| \leq |\gamma|/\beta k$ , we can use for  $|\gamma| < \beta k$  the estimate  $c a_{\gamma-1} \leq (|\gamma|/\beta k)^2$ . For  $|\gamma| \geq \beta k$  we use the estimate

$$c a_{\gamma-1} \leq \exp \left\{ \frac{|\gamma|}{\beta k} \right\}$$

Then under condition 1 we have

$$P \leq \sum_{\substack{4 \leq l < \beta k \\ l \in \mathcal{N}}} n(l) e^{-\nu_1 l} \frac{l^3}{\beta^2 k^2} + \sum_{\substack{l \geq \beta k \\ l \in \mathcal{N}}} n(l) l e^{-\nu_1 l} e^{l/\beta k}$$

where  $\nu_1 \geq \log 2/d$ , with  $d > 0$ . Since  $n(l) \leq (2d)^l$  we have

$$P \leq \sum_{l < \beta k} e^{-\alpha l} l^3 \beta^{-2} k^{-2} + \sum_{l \geq \beta k} l e^{-\alpha l} e^{l/\beta k}$$

Let  $\beta_2$  such that  $\beta_2 k > \frac{1}{\alpha}$ . Then for  $\beta \geq \sup\{\beta_0, \beta_1\}$  we obtain

$$P \leq A \beta^{-2} k^{-2} + A' e^{-\alpha \beta k}$$

where  $A$  and  $A'$  are positive constants. Therefore

$$\langle e^{iA_0} \rangle(\beta) \leq \exp\{-18(c) \beta^{-2} k^{-2} (1 - A k^{-2} - \beta^2 k A' e^{-\alpha \beta k})\}$$

we choose  $k > 2A$ . Let  $\beta_2$  such that  $\beta_2^2 k A e^{-\alpha \beta_2 k} < 1/2$ .

Then for  $\beta \geq \sup\{\beta_0, \beta_1, \beta_2\}$  we obtain statement A of Theorem 2 for large

$\beta$ . By using inequality (8) one extends the proof to any positive  $\beta$ .

The same method is applied to prove statement c of Theorem 2.

### III.3 Proof of part a) of Theorem 3

Let  $d = 2$ , and  $S_1$  be the rectangle of vertices  $O \equiv (0,0)$ ,  $x_1 \equiv (\tau, 0)$ ,  $x_2 \equiv (\tau, L)$ ,  $x_3 \equiv (0, L)$ . Let  $S_2$  be the symmetric of  $S_1$  with respect to  $Ox^1$  axis and and  $S_0 = S_1 \cup S_2$ .




Figure 1

We now choose a configuration  $\{a(\ell)\}_{\ell \in d_0}$  verifying the following conditions.

$$(11) \left\{ \begin{array}{l} \text{for the links } \ell \text{ such that } \ell \in \Lambda/S_0 \text{ we take } a(\ell) = 0 \\ \text{for the links } \ell \text{ such that } \ell \in \partial S_0 \text{ we take } a(\ell) = 0 \\ \text{for the links } \ell \text{ parallel to the direction } O x^1 \text{ we take } a(\ell) = 0 \\ \text{for the links } \ell \text{ parallel to the direction } O x^2 \text{ we take} \\ \text{if } x^1 \geq 0 \quad a[\{x^1, x^2\}, \{x^1+1, x^2\}] - a[\{x^1, x^2+1\}, \{x^1+1, x^2+1\}] = \frac{1}{\beta k} \\ \text{if } x^1 < 0 \quad a[\{x^1, x^2\}, \{x^1+1, x^2\}] = a[\{x^1, -x^2\}, \{x^1+1, -x^2\}] \end{array} \right.$$

$k$  is a positive constant chosen later.

Under these conditions, for the  $b(p)$  variables we have

$$|b(p)| = \beta^{-1} k^{-1} \text{ if } p \in S_0, b(p) = 0 \text{ otherwise.}$$

Let  $p$  be some plaquettes of  $S_0$ . By using part b) of Lemma 1 we obtain

$$\langle e^{i A \phi_S}, \rangle (\beta) \leq \exp \left\{ -L T \beta^{-1} k^{-1} \right\} \exp \left\{ \beta \beta L T \sum_{S \supset p} \kappa_S (ch b_S - 1) \right\}$$

If  $\beta c$  is large enough ( $\beta > \beta_0$ , with  $\beta_0 \rightarrow \frac{1}{k}$ ) we can write

$$Q = \sum_{S \supset p} \kappa_S (ch b_S - 1) = \sum_{\substack{S \supset p \\ |S| < \beta k}} \kappa_S (ch b_S - 1) + \sum_{S \supset p} \kappa_S (ch b_S - 1)$$

For  $|S| < \beta k$  we use the estimate

$$ch b_S - 1 \leq (|S| \beta^{-1} k^{-1})^2$$

For  $|S| \geq \beta k$  we use  $ch b_S - 1 \leq e^{|S| \beta^{-1} k^{-1}}$

Then under condition 3 we have:

$$Q \leq \sum_{\substack{\lambda < \beta k \\ \lambda \in \mathbb{N}}} \mu_d^\lambda e^{-\mu_d^\lambda} \lambda^2 \beta^2 k^2 + \sum_{\substack{\lambda \geq \beta k \\ \lambda \in \mathbb{N}}} \mu_d^\lambda e^{-\mu_d^\lambda} e^{\lambda \beta^{-1} k^2}$$

where  $\mu_d \geq \log \mu_d + d$ ,  $d > 0$ . Let  $\beta_2$  be such that  $\beta_2 k > \frac{1}{\alpha}$ . For  $\beta \geq \sup(\beta_1, \beta_2)$  we obtain:

$$Q \leq A \beta^{-2} k^{-2} + A' e^{-\alpha \beta k}$$

$A$  and  $A'$  are positive constants. The proof of inequality a) of Theorem 3 ends analogously to III.2. To prove statement c) of Theorem 2, we use the same method but in choosing the configuration given by (10).

We now consider the 3-dimensional case. The idea of the proof consists in choosing a configuration  $\{a(\ell)\}_{\ell \in \mathcal{L}}$  to reduce it to a bidimensional problem. We first introduce some notations.

#### III.4 Notations

Let  $x \equiv \{x^1, x^2, x^3\}$  be a site of  $\Lambda$

We denote by  $d(x)$  the distance of  $x$  to the  $Ox^3$  axis

$$d(x) = \text{dist}(x, Ox^3) = \sup\{|x^1|, |x^2|\}$$

We define the projection of  $x$  on the half-plane  $\{x^3 = 0, x^2 \geq 0\}$

$$P_{20j}[\{x^1, x^2, x^3\}] = \{y^1, y^2, y^3\}$$

where  $y^1 = x^1$ ,  $y^2 = d(x)$ ,  $y^3 = 0$

Let  $\ell = \langle x, y \rangle$  be a link. We define the projection of the link  $\ell$  on the half-plane  $\{x^3 = 0, x^2 \geq 0\}$

$$P_{20j}[\ell] = \langle P_{20j}[x], P_{20j}[y] \rangle$$

We consider the links  $\ell = \langle x, y \rangle$  parallel to  $Ox^1$  and introduce the distance of  $\ell$  to  $Ox^1$

$$d(\ell) = d(x) = d(y)$$

Let  $p = (x_1, x_2, x_3, x_4)$  be some plaquettes such that

$$P_{\alpha\beta j} [x_i] \neq P_{\alpha\beta j} [x_j] \quad \forall i, \forall j \quad i \neq j$$

We define the projection of the plaquette  $p$  as

$$P_{\alpha\beta j} [p] = (P_{\alpha\beta j} [x_1], P_{\alpha\beta j} [x_2], P_{\alpha\beta j} [x_3], P_{\alpha\beta j} [x_4])$$




Figure 2

Let  $p$  be a plaquette on the half-plane  $\{x^2 = 0, x^3 \geq 0\}$

We define the "tube"  $\mathcal{T}_p$  associated to the plaquette  $p$  by

$$\mathcal{T}_p = \{ \text{set of plaquettes } q \text{ such that } \{P_{\alpha\beta j} [q]\} = p \}$$

We define the distances of the plaquette  $p = (x_1, x_2, x_3, x_4)$  to  $Ox^1$

$$d(p) = \min_{x_i \in p} d(x)$$

The distances of the tube  $\zeta_p$  to  $Ox^1$  are given by

$$d(\zeta_p) = d(p)$$




Figure 3

### III.5 Proof of statement b of Theorem 3

We consider the rectangle  $S_2$  of vertices  $O \equiv \{0, 0, 0\}$ ,  $x_2 \equiv \{\tau, 0, 0\}$ ,  $x_3 \equiv \{\tau, L, 0\}$ ,  $x_4 \equiv \{0, L, 0\}$ , and the box

$$\Lambda_{LT} : \left\{ 0 \leq x^1 \leq \tau, -L \leq x^2 \leq L, -L \leq x^3 \leq L \right\}$$

We choose a configuration  $\{a(\ell)\}_{\ell \in \mathcal{L}}$  verifying the following conditions :

$$(12) \quad \left\{ \begin{array}{l} \text{for all links } \ell \text{ perpendicular to } O x^1 \text{ direction we take } a(\ell) = 0 \\ \text{for all links of } \partial \Lambda_{LT} \text{ and } \Lambda / \Lambda_{LT} \text{ we take } a(\ell) = 0 \\ \text{for the links in } \Lambda_{LT} \text{ parallel to } O x^1 \text{ and oriented in} \\ \text{the } O x^1 \text{ direction we take} \\ a(\ell) = \frac{1}{\beta k} \sum_{m=d(\ell)}^{L-2} \frac{1}{m+1} \end{array} \right.$$

$k$  is a positive constant chosen later.

With this choice, for the  $b(p)$  variables we have

$$\left\{ \begin{array}{l} \forall p \in S_2, \forall q \in \mathcal{L}_p \quad |b(q)| = \frac{1}{\beta k (d(p)+1)} \\ b(p) = 0 \quad \text{otherwise} \end{array} \right.$$

Using part b of Lemma 1 and assuming that the configuration verifies the condition (12) we obtain

$$(13) \quad \langle e^{L A_{\partial S_2}} \rangle (\beta) \leq \exp \{-a_{\partial S_2}\} \exp \left\{ \beta \sum_{p \in S_2} \sum_{q \in \mathcal{L}_p} \sum_{S \supset q} K_S (c^{\beta} b_{S-1}) \right\}$$

with

$$(14) \quad \exp \{-a_{\partial S_2}\} = \exp \left\{ -\tau \beta^{-1} k^{-2} \sum_{j=1}^L \frac{1}{j} \right\}$$



We can write

$$Q' = \beta \sum_{p \in S} \sum_{q \in \bar{p}} \sum_{s > q} K_s(ch b_s - 1) \leq \beta T \sum_{j=1}^h 4(2j-1) \sum_{\substack{s > p \\ d(p)=j}} K_s(ch b_s - 1)$$

We can decompose the sum  $Q'$  as follows

$$(15) \quad Q' \leq \beta T \sum_{j=1}^h 4(2j-1) \sum_{\substack{s > p: d(p)=j \\ |s| < d/2}} K_s(ch b_s - 1) \\ + \beta T \sum_{j=1}^h 4(2j-1) \sum_{s > p: d(p)=j, |s| > d/2} K_s(ch b_s - 1)$$

In the first term of the R.H.S. of (15) we use the estimate

$$ch b_s - 1 \leq \left( \frac{2|s|}{(j+1)\beta k} \right)^2$$

In the second term of R.H.S. of (15) we use the estimate

$$ch b_s - 1 \leq e^{|s|} \beta^{-1} k^{-1}$$

Then under condition 3 on  $K_s$

$$Q' \leq \beta T \sum_{j=1}^h 4(2j-1) \sum_{\substack{\lambda < d/2 \\ \Delta \in \mathcal{N}}} \frac{4\lambda^2 e^{-\lambda^2} \mu_d^\lambda}{\beta^2 k^2 (j+1)^2} + \beta T \sum_{j=1}^h 4(2j-1) \sum_{\substack{\lambda > d/2 \\ \Delta \in \mathcal{N}}} e^{-\lambda^2} \mu_d^\lambda e^{\lambda} \beta^{-1} k^{-1}$$

where  $\mu_s \geq \log \mu_d + \alpha$ , with  $\alpha > 0$ . For  $\beta > d^{-1} k^{-1}$  we obtain

$$(16) \quad Q' \leq \beta T \left\{ A \beta^{-2} k^{-2} \sum_{j=1}^h \frac{1}{j} + A' \right\}$$

where  $A, A'$  are positive constants. By choosing  $k > A$  a statement b of Theorem 3 follows from (13), (14) and (16).

### III.6 Proof of Statement a of Theorem 2

We keep the notation of Sections III.3 and III.4. We consider a configuration  $\{a(\ell)\}_{\ell \in \mathcal{L}}$  verifying the following conditions

$$(17) \quad \left\{ \begin{array}{l} \text{for all links of } \mathcal{D}S_0 \text{ and } \Lambda/S_0 \text{ we take } a(\ell) = 0 \\ \text{for all links parallel to } O_2^2 \text{ we take } a(\ell) = 0 \\ \text{for all links } \ell \text{ in } S_0 \text{ parallel to } O_2^4 \text{ and oriented} \\ \text{in the } O_2^4 \text{ direction we take} \\ a(\ell) = \frac{1}{\beta^k} \sum_{m=d(\ell)}^{b-1} \frac{1}{m+1} \end{array} \right.$$

We shall assume  $k = 1$ . Under these conditions for the  $b(p)$  variables we have

$$|b(p)| = \beta^{-1} k^{-1} (d(p)+1)^{-1} \text{ if } p \in S_0, \quad b(p) = 0 \text{ otherwise}$$

Using part a) of Lemma 1 for a configuration verifying the conditions (11) we obtain

$$(18) \quad \langle e^{iA\mathcal{D}S_0} \rangle (\beta) \leq \exp \left\{ -\frac{T}{\beta^k} \sum_{j=1}^b \frac{1}{j} \right\} \exp \left\{ \beta \sum_{r \in \Lambda(r)} J_r (d a_r - 1) \right\}$$

$a_r \neq 0$

Let

$$(19) \quad R = \sum_{r: a_r \neq 0} J_r (d a_r - 1)$$

we can write

$$R \leq \sum_{p \in S_0} \sum_{\substack{r \in \Lambda(r) \\ \gamma(r) \text{ contains a link of } p}} J_r |r|^2 (d a_r - 1)$$

It is clear that

$$R \leq 2T \sum_{j=1}^L \sum_{\substack{M: \\ \{\delta(M) \text{ contains } a \\ \text{link of } p; d(p)=j\}}} |M|^2 J_M(\text{ch } a_{p-1})$$

Let  $\epsilon$  a some positive constant larger than 3. For  $\beta k$  large enough we make the following decomposition of  $R$ .

$$(20) \quad R \leq 2\epsilon T \sum_{\substack{M: \\ \left\{ \begin{array}{l} \delta(M) \text{ contains } a \\ \text{given link} \\ |M| < \sqrt{\beta k} \end{array} \right\}}} |M|^2 J_M(\text{ch } a_{p-1}) + 2\epsilon T \sum_{\substack{M: \\ \left\{ \begin{array}{l} \delta(M) \text{ contains } a \\ \text{given link} \\ |M| \geq \sqrt{\beta k} \end{array} \right\}}} |M|^2 J_M(\text{ch } a_{p-1}) \\ + 2T \sum_{j=\epsilon}^L \sum_{\substack{M: \\ \left\{ \begin{array}{l} \delta(M) \text{ contains} \\ \text{a link of } p \\ d(p)=j, |M| < \sqrt{\beta k} \end{array} \right\}}} |M|^2 J_M(\text{ch } a_{p-1}) + 2T \sum_{j=\epsilon}^L \sum_{\substack{M: \\ \left\{ \begin{array}{l} \delta(M) \text{ contains} \\ \text{a link of } p \\ d(p)=j, |M| \geq 4\sqrt{\beta k} \end{array} \right\}}} |M|^2 J_M(\text{ch } a_{p-1})$$

Let  $R_1, R_2, R_3, R_4$  the first second third and fourth terms of the R.H.S. of the inequality (20). We now use the estimates :

$$\text{ch } a_{p-1} \leq \left( \frac{|M|}{\beta k} \right)^2 \quad \text{in } R_1$$

$$\text{ch } a_{p-1} \leq \text{sup} \left\{ \frac{|M|}{2\beta k}, \log(|M|/2) \right\} \quad \text{in } R_2 \quad \text{and } R_4$$

$$\text{ch } a_{p-1} \leq \left( \frac{|M|}{\beta k} \right)^2 \quad \text{in } R_3$$

Under the condition 2 on  $J_M$  we obtain for large  $\beta$ .

$$R_1 \leq 2\epsilon T \sum_{\substack{\ell_1=1 \\ \ell_1 \leq \sqrt{\beta k}}}^{\sqrt{\beta k}-1} e^{-\mu_2 \ell \log \ell} \ell^2 \beta^{-2} k^{-2} e^{\ell \log 2d} \leq H_3 T \beta^{-2}$$

$$R_2 \leq 2\epsilon T \sum_{\substack{\ell_2 \geq \sqrt{\beta k}}} \ell^2 e^{-\mu_2 \ell \log \ell} e^{\ell \beta^{-1} k^{-1} \log \frac{\ell}{2}} e^{\ell \log 2d} \leq A_2 \epsilon T$$

$$R_3 \leq 2T \sum_{j=c}^L \sum_{\frac{d}{2}=z}^{2\sqrt{j-1}} e^{-A_2 \ell \log \ell} \ell^2 (j \beta^4)^{-z} e^{\ell \log^2 d} \leq A_3 T \beta^{-z}$$

$$R_4 \leq 2T \sum_{j=c}^L \sum_{\ell \geq 4\sqrt{j}} \ell^2 e^{-A_2 \ell \log \ell} \ell^2 \beta^{k' \log \frac{1}{2} \ell} e^{\ell \log^2 d} \leq A_4 T$$

$A_1, A_2, A_3$  and  $A_4$  are positive constants. From these four inequalities and from (18), (19), (20) follows the proof of statement a) of Theorem 2 at large  $\beta$ . Ginibre inequality extends the proof to any positive  $\beta$ .

### III.7 Proof of Part b of Theorem 2

In this case we choose a configuration  $\{a(\ell)\}_{\ell \in d}$  verifying the condition (12) as in III.5. Using part a) of Lemma 1 for this configuration we obtain

$$(21) \quad \langle e^{iA_0 S_0} \rangle (\beta) \leq \exp \left\{ -\beta^{-k'} \sum_{j=1}^L \frac{1}{j} \right\} \exp \left\{ \beta \sum_{\substack{r \in \Lambda(r) \\ a_r \neq 0}} J_r (c a_r - 1) \right\}$$

$$\text{Let } R' = \sum_{r: a_r \neq 0} J_r (c a_r - 1)$$

We can write

$$R' \leq \sum_{p \in S_0} \sum_{q \in \mathcal{C}_p} \sum_{\substack{r: \\ \{ \gamma(r) \text{ contains} \\ \text{a link of } q \}}} |r|^\ell J_r (c a_r - 1)$$

It is clear that

$$R' \leq T \cdot \sum_{j=1}^L 4(2j-1) \sum_{\substack{P: \\ \left\{ \begin{array}{l} \gamma(P) \text{ contains a} \\ \text{link } i; p, d(p)=j \end{array} \right\}}} |P|^2 J_P (c h a_{P-1})$$

We remark that  $R'$  differs from  $R$  only by the factor  $2(2j-1)$ . By using the same decomposition and estimates as in Section III.6 we obtain

$$R' = R'_1 + R'_2 + R'_3 + R'_4 \quad \text{with}$$

$$R'_1 \leq T A'_1 \beta^{-1} ; \quad R'_2 \leq T A'_2 ; \quad R'_3 \leq T A'_3 (\beta k)^2 \sum_{j=c}^L \frac{1}{j} ; \quad R'_4 \leq A'_4 T$$

where  $A'_1, A'_2, A'_3$  and  $A'_4$  are positive constants. By choosing  $k > A'_3$  we obtain part b) of Theorem 2.

#### ACKNOWLEDGEMENTS

The author is very indebted to A. Messenger and S. Miracle-Sole for helpful comments and discussions. He would like to thank R. Senor for encouragements and for pointing out to him the relation between long range and fermions. He would like to thank also C.P. Korthals-Altes, P. Picco and R. Kotecky for useful discussions.

## REFERENCES

-0-0-

- [1] K.G. WILSON, Phys. Rev. D10, 2245 (1974).
- [2] G.B. LANG, H. NICOLAI, Nucl. Phys. B200 FS4 , 135 (1982).
- [3] T. MATTHEWS, A.SALAM, Nuov. Cim. 12, 563 (1954).
- [4] F.A. BEREZIN, "The method of second quantization", Academic Press New York (1966).
- [5] E. SEILER, in "Gauge theories : fundamental interactions and rigorous results", Lectures given at the 1981 International Summer School of Theoretical Physics, Poiana Brasov, Romania, Progress in Physics, Vol.5, edited by A. Jaffe and D. Ruelle.
- [6] L. SUSSKIND, Phys. Rev. D16, 3031 (1977).
- [7] E. MARIANI, G. PARISI, C. REBBI, Nucl. Phys. B190 (FS3), 734 (1981).
- [8] P.W. KASTELEYN, in "Graph theory and theoretical physics", edited by Frank Harari (1967), Academic Press (London and New York).
- [9] D. BRYDGES, J. FROLICH, T. SPENCER, Comm. math. Phys. 83, 123 (1982).
- [10] R. BALIAN, J.M. DROUFFE, C. ITZYKSON, Phys. Rev. D10, 3376 (1974).
- [11] J. GLIMM, A. JAFFE, Phys. Letters B66, 67 (1977).
- [12] A.M. GUTH, Phys. Rev. D21, 2291 (1981).

- [13] R. KOTECKY, S.B. SHLOSMAN. *Comm. math. Phys.* 83, 493 (1982).
- [14] J. GINIBRE, *Comm. math. Phys.* 16, 310 (1970).
- [15] J. MAC BRYAN, T. SPENCER, *Comm. math. Phys.* 52, 299 (1977).
- [16] A. MESSEGER, S. MIRACLE-SOLE, C.E. PFISTER, *Comm. math. Phys.* 58, 19 (1978).