

Gamma-rays from Dark Matter Annihilation in the Central Region of the Galaxy

Pasquale Dario Serpico

Physics Division, Theory Group, CERN, CH-1211 Geneva 23, Switzerland

E-mail: serpico@cern.ch

Dan Hooper

Theoretical Astrophysics, Fermi National Accelerator Laboratory, Batavia, USA

Department of Astronomy and Astrophysics, The University of Chicago, USA

E-mail: dhooper@fnal.gov

Abstract. In this article, we review the prospects for the FERMI satellite (formerly known as GLAST) to detect gamma-rays from dark matter annihilations in the Central Region of the Milky Way, in light of the recent observations and discoveries of Imaging Atmospheric Cherenkov Telescopes. While the existence of significant astrophysical backgrounds in this part of the sky limits FERMI's discovery potential to some degree, this can be mitigated by exploiting the peculiar energy spectrum and angular distribution of the dark matter annihilation signal relative to those of astrophysical backgrounds.

Keywords: Dark Matter, Gamma Rays, Galactic Center

1. Introduction

Despite the numerous cosmological and astrophysical indications of the presence of non-baryonic dark matter (DM), the particle nature of this substance remains unknown. If the DM consists of weakly interacting massive particles (WIMPs), an important tool for inferring their properties could be “indirect detection”, i.e. astrophysical observations of the annihilation (or decay) products of DM in our Galaxy or beyond. For WIMPs with masses at or around the electroweak scale, $m_X \sim \mathcal{O}(0.1-1)$ TeV, the annihilation products are typically injected at GeV-TeV energies, the domain of high-energy astrophysics. Of the different annihilation products, gamma-rays and neutrinos have the important advantage of retaining directional information while not suffering energy losses. The very small cross sections of neutrinos, however, make their flux from the region of the Galactic Center very difficult to detect. On the contrary, the gamma-ray spectrum from DM annihilation or decay may be detectable with sufficient statistics, energy resolution, and over an extended angular distribution, to provide a very distinctive set of information related to both the particle identity of the WIMP and its astrophysical distribution. A major challenge is in separating the DM signal from any astrophysical backgrounds, whose energy spectrum and angular distribution are not well known.

A major change in the prospects for DM detection has occurred in the last few years, following the discovery of a bright astrophysical source of TeV gamma-rays from the Galactic Center. As a result of this discovery, we now know that DM emission from the Galactic Center will not be detectable in a (quasi) background-free regime, and—unless one turns the attention to other targets—the peculiar spectral shape and angular distribution of the signal must be used to extract it from this and other backgrounds. This topic is the main subject of this review. The remainder of this paper is structured as follows: In Sec. 2 we briefly describe the capabilities of current high energy gamma-ray telescopes. In Sec. 3 we review the features of the DM annihilation signal, both in its energy (Sec. 3.1) and angular dependence (Sec. 3.2). In Sec. 4 we review the prospects for detecting or constraining DM annihilations in the Galactic Center region, while in Sec. 5 we briefly discuss some interesting aspects of the diffuse signal from the inner halo. In Sec. 6 we report our conclusions. Appendix A briefly describes the qualitative changes to the considerations we develop in the main text when decaying (as opposed to annihilating) DM is considered. Appendix B describes a subtlety regarding the angular shape of the signal arising for a velocity-dependent annihilation cross section.

2. The Instruments

The ability of gamma-ray experiments to identify DM annihilation radiation from the Galactic Center region relies on the effective area, angular and energy resolution of the existing telescopes, as well as the rejection of other (mostly hadronic) cosmic ray events contaminating the gamma-ray sample. Since the mean free path of gamma-

rays is much shorter than the atmospheric slant depth, direct observations in the GeV region and above can only be done from space—which is the strategy pursued by the FERMI gamma-ray space telescope (formerly GLAST) [1]—or indirectly by ground-based Imaging Atmospheric Cerenkov Telescopes (IACTs) such as HESS [2], MAGIC [3], VERITAS [4] and CANGAROO-III [5]. In the latter category, the direction and energy of the primary particle hitting the atmosphere is reconstructed from the Cherenkov emission of the secondary charged particles generated in the atmospheric shower.

These differences lead these two classes of experiments to adopt different strategies in the search for DM. FERMI is very effective in rejecting hadronic events, and continuously monitors a large fraction of the sky, but has an effective area of only $\sim 1 \text{ m}^2$, far smaller than that of ground-based telescopes, $\sim 10^4 \text{ m}^2$. On the other hand, IACTs study small angular fields and have a lower rejection capability, but much greater overall exposure. As a consequence, diffuse gamma-ray signals are better probed by FERMI. Any unidentified sources detected by FERMI which lack a low-energy counterpart could be potentially attributed to DM substructure. IACTs would be very effective in providing detailed follow-up observations of such sources.

In addition, the accessible energy range is very different between these two classes of experiments: $\sim 100 \text{ MeV}$ to 300 GeV for FERMI, and above $\sim 100 \text{ GeV}$ for ACTs. This difference makes FERMI most sensitive to DM particles lighter than a few hundreds GeV, while IACTs are better suited for TeV-scale or heavier WIMPs. For both instrument classes, the search for indirect DM signatures is among the top physics priorities. The reach of FERMI and current and future IACTs has been recently assessed in Ref. [6] and Ref. [7], respectively.

3. The Dark Matter Signal

The energy and angular dependent flux of gamma-rays produced in DM annihilations is described by

$$\Phi_\gamma(E_\gamma, \Omega) = \left[\frac{dN_\gamma}{dE_\gamma}(E_\gamma) \frac{\langle \sigma v \rangle}{8\pi m_X^2} \right] \int_{\text{los}} \rho^2(\ell, \Omega) d\ell, \quad (1)$$

where $\langle \sigma v \rangle$ is the WIMP annihilation cross section multiplied by the relative velocity of the two WIMPs (averaged over the WIMP velocity distribution), m_X is the mass of the WIMP, ρ is the position-dependent DM density, and the integral is performed over the line-of-sight (los) in the direction of the sky, Ω . The gamma-ray spectrum generated per WIMP annihilation is dN_γ/dE_γ . If the DM is not its own antiparticle as assumed here, Eq. (1) should be multiplied further by a factor 1/2 (if X and \bar{X} are equally abundant). The factor in square brackets in Eq. (1) depends only on particle physics: in particular, cross section, mass, and the spectrum of gamma-rays produced through DM annihilations depends on the nature of the WIMP. The integral over the line-of-sight determines instead the angular dependence of the signal and is controlled by the astrophysical distribution of DM. We shall discuss each of these two terms in

the following, assuming that this factorization holds (see Appendix B for a discussion of possible violations of this hypothesis).

In convenient units, Eq. (1) can be recast as:

$$\frac{\Phi_\gamma(E_\gamma, \Omega)}{\text{cm}^{-2} \text{s}^{-1} \text{sr}^{-1}} \approx 2.8 \times 10^{-10} \frac{dN_\gamma}{dE_\gamma} \left(\frac{\langle \sigma v \rangle}{3 \times 10^{-26} \text{ cm}^3 \text{ s}^{-1}} \right) \left(\frac{100 \text{ GeV}}{m_X} \right)^2 J(\Omega). \quad (2)$$

The dimensionless function $J(\Omega)$ depends only on the DM distribution in the halo and is defined by convention as [8]

$$J(\Omega) = \frac{1}{8.5 \text{ kpc}} \left(\frac{1}{0.3 \text{ GeV/cm}^3} \right)^2 \int_{\text{los}} \rho^2(\ell, \Omega) d\ell. \quad (3)$$

The benchmark value for the cross-section, $\langle \sigma v \rangle \approx 3 \times 10^{-26} \text{ cm}^3/\text{s}$, is motivated by the fact that a WIMP annihilating with such a cross section during the freeze-out epoch will be generated as a thermal relic with a density similar to the measured DM abundance (for a review, see Ref. [9]). WIMPs constituting the *cold* DM annihilate in the non-relativistic limit. If annihilations take place largely through S -wave processes, then the annihilation cross section of WIMPs in the Galactic halo (i.e. in the low velocity limit) will also be approximately equal to this value, which justifies the benchmark value used in Eq. (2). Yet, it is worth stressing that much lower signals are possible (e.g. if P -wave annihilation dominates the freeze-out process), as well as significantly enhanced ones in models where the DM is non-thermally produced in the early universe (see e.g. Ref. [10]).

3.1. Energy Spectra

The gamma-ray spectrum from DM annihilation originates from several different contributions. Typically, the most abundant source of photons is the hadronization and/or decays of unstable particles. For example, neutralinos in the Minimal Supersymmetric Standard Model (MSSM), dominantly annihilate to final states consisting of heavy fermions ($b\bar{b}$, $t\bar{t}$, $\tau^+\tau^-$) or gauge and/or Higgs bosons (ZZ , W^+W^- , HA , hA , ZH , Zh , ZA , $W^\pm H^\pm$, where H , h , A and H^\pm are the Higgs bosons of the MSSM) [11]. With the exception of the $\tau^+\tau^-$ channel, each of these annihilation modes result in a very similar spectrum of gamma-rays, dominated eventually by the decay of mesons, especially π^0 , generated in the cascade. In Fig. 1 we show the predicted gamma-ray spectrum per annihilation, for several possible WIMP annihilation modes. The harder spectrum here shown from annihilation into the $\tau^+\tau^-$ channel is not typical of SUSY models, although it might be a distinctive feature of DM annihilating dominantly to charged leptons pairs; this arises, for example, in Kaluza-Klein DM models.

Although these secondary photons provide the dominant emission, other important channels can exist at the one-loop level. A particularly striking signature would be the mono-energetic photons resulting from final states such as $\gamma\gamma$, γZ or γh (see Ref. [13] for a discussion of these processes within the context of supersymmetry). Unfortunately, such processes are expected to produce far fewer events than continuum emission and in typical models can not easily be detected (for a counterexample, see Ref. [14]).




Figure 1. The gamma-ray spectrum per annihilation for a 100 GeV (left) and 500 GeV (right) WIMP. Each curve denotes the result for a different dominant annihilation mode. From Ref. [12].

Bremsstrahlung (with an additional photon appearing in the final state) is automatically present when annihilations produce charged final states, and can dominate the high energy region of the spectrum when those charged annihilation products are much lighter than the WIMPs. This is particularly important when the tree-level processes to light fermions are disfavored by “selections rules”, but without suppressing the three body final state [15, 16]. For more details, see the review by L. Bergstrom in this issue. Note that gamma-ray emission from DM annihilating into standard model particles is unavoidable, even in the most extreme case when only neutrino final states are allowed at tree level, due to W and Z -strahlung [17, 18].

3.2. Angular Shape Of The Signal

The gamma-ray spectrum and angular distribution predicted by Eq. (1) is rather general (see, however, Appendix B), and could be applied to the case of a smoothly distributed Galactic halo, or alternative targets such as dwarf spheroidal galaxies, microhalos/clumps, density spikes around intermediate mass black holes, or the integrated extragalactic diffuse background (see contributions from S. Koushiappas, G. Bertone, and L. Strigari). While some of these targets have interesting observational prospects, the intensity of these signals is strongly dependent on often unknown cosmological and astrophysical properties, such as the quantity of small-scale structures in DM halos or the population of intermediate mass black holes in the Milky Way. In this article, we will focus on the smooth Galactic halo.

Naively, by inputting the DM density profile inferred from kinematical observations of the Milky Way into Eq. (1), one could obtain an approximate lower bound on the gamma-ray flux from DM annihilation. In principle, this would enable us to translate the observations from ground or space-based gamma-ray telescopes into constraints on the

particle physics properties of the WIMP (mass, annihilation cross section and dominant modes). Unfortunately, even the average, smooth distribution of the DM particles in our Galaxy is not well known, especially in the volume within the solar circle. This is due to the fact that the baryonic material dominates the gravitational potential in the inner Galaxy, and lacking a detailed knowledge of its distribution, a reconstruction of the DM distribution “by subtraction” is unfeasible. It is not surprising, then, that very different profiles have been claimed to fit the observations (for example, compare the results of Ref. [19] and Ref. [20]).

On general grounds, a class of spherically symmetric, smooth halo distributions can be used to approximately fit both the observed rotation curves of galaxies and the results of numerical simulations of DM halos. A non-trivial angular dependence of the gamma-ray signal results from the off-center position of the Sun within the halo (see e.g. Ref. [21] and references therein for a discussion of sub-leading effects determining the angular distribution of the signal). The function J introduced in Eq. (2) and Eq. (3) then depends only on the angle θ between the observed direction of the sky and the Galactic Center or, in terms of galactic latitude b and longitude l , only on $\cos\theta = \cos b \cos l$. The radial variable, r , can be expressed in terms of the relevant quantities $\{\ell, \theta\}$ as

$$r(\ell, \theta) = \sqrt{r_\odot^2 + \ell^2 - 2r_\odot \ell \cos\theta}, \quad (4)$$

where $r_\odot \approx 8.33 \pm 0.35$ kpc [22] is the distance of the Solar System from the Galactic Center. Typically, one considers a distribution of the form

$$\rho(r) = \left(\frac{r_s}{r}\right)^\gamma \frac{\rho_0}{[1 + (r/r_s)^\alpha]^{(\beta-\gamma)/\alpha}}, \quad (5)$$

where ρ_0 is a normalization constant and r_s is a characteristic radius below which the profile scales as $r^{-\gamma}$. A very well known, universal profile of this class fit to DM-only (*ie.* neglecting baryons) N-body simulations has been proposed by Navarro, Frenk and White (NFW) [23], corresponding to the choice $\{\alpha, \beta, \gamma\} = \{1.0, 3, 1.0\}$. Steeper or softer profiles have also been extensively discussed, such as that proposed by Moore et al. [24] and Kravtsov et al. [25], respectively. While simulations (and data as well) typically agree on the shape of profile in the outskirts of the halos, a disagreement clearly exists concerning the inner slope, γ . More recent simulations [26, 27, 28, 29, 30] suggest that halo density profiles are better represented by a function with a continuously-varying slope, as the one proposed by Einasto [31]

$$\rho(r) = \rho_{-2} \exp\left[-\frac{2}{\alpha}(r/r_{-2} - 1)\right] \quad (6)$$

with clear hints for non-universality in the form of halo-to-halo variations in the quantity α . The quantity $r_{-2} \approx 25$ kpc denotes the radius at which the profile takes on a slope of -2. At $r \gtrsim 1$ kpc, these newly proposed fitting formulae provide only marginal improvement with respect to the more traditional ones. At smaller radii, however, these recent results lead us to expect the inner slope of DM halos to be shallower than that predicted by Moore *et al.*, and probably shallower than NFW, as well.

Model	α	β	γ	ρ_{\odot} [GeV cm ⁻³]	r_s [kpc]
Moore <i>et al.</i>	1.5	3	1.5	0.27	28
NFW	1.0	3	1.0	0.30	20
Kravtsov	2.0	3	0.4	0.37	10

Table 1. Parameters describing some common halo profiles of the form described by Eq. (5). See text for more details.

There are a number of astrophysical processes that may potentially modify the DM distribution, none of which are taken into account in the above-mentioned profiles. It is very difficult to reliably account for these effects in simulations, and only a few results are available (see e.g. Refs. [32, 33, 34]). Qualitatively, since baryons can cool and contract, one expects them to steepen the gravitational potential in the central regions galaxies and, as a result, enhance the DM density [35]. On the other hand, other feedback or frictional effects have been proposed that could reduce the DM density in the inner halos and bring the prediction closer to observations (for a recent review, see Ref. [36]). For some galaxies, it has actually been argued that flat-cored profiles better fit the observations better than cusped profiles.

Even greater uncertainties exist concerning the DM distribution at the very center of the Milky Way, in the region immediately surrounding the central supermassive black hole. Adiabatic accretion may lead to the formation of a spike in the DM distribution, resulting in a very high DM annihilation rate in the innermost parsecs of the galaxy [37]. Mergers as well as scattering on the dense stellar cusp around the central black hole may potentially destroy density enhancements, however. In general, these effects only affect regions too close to the Galactic Center to be resolved angularly by present detectors, leaving only the energy spectrum to be used for separating the DM signal from the background. Yet, such a spike might lead to a measurable gamma-ray flux from the innermost angular bin, even in presence of relatively large astrophysical backgrounds. For more details, we direct the reader to the references cited in Ref. [9] or Ref. [38].

Given these considerably uncertainties, we have chosen to use three illustrative choices for the DM halo profiles: the Moore *et al.*, NFW, and Kravtsov profiles (with the parameters given in Table I). In each case, the normalization has been chosen so that the mass contained within the solar circle provides the appropriate DM contribution to the local rotational curves. In all of the cases we have discussed, even the most conservative cored profiles, the DM signal peaks at the Galactic Center. In presence of an isotropic (or even vanishing) astrophysical background, the Galactic Center region thus becomes the natural location to look for a DM signal. Unfortunately, there is also a higher background density as we look toward the inner region of our Galaxy. Still, as a first step, one can consider the detection prospects for this region, a task we address in Sec. 4, before turning to more general arguments regarding the optimal regions of the




Figure 2. Spectral energy density of the Galactic Center source as measured by HESS in 2004 (full points) and 2003 [42] (open points). Upper limits are 95% CL. The shaded area shows the power-law fit $dN/dE \sim E^{-\Gamma}$. The dashed line illustrates typical spectra of phenomenological MSSM DM annihilation for best fit neutralino masses of 14 TeV. The dotted line shows the distribution predicted for Kaluza-Klein DM with a mass of 5 TeV. The solid line gives the spectrum of a 10 TeV DM particle annihilating into $\tau^+\tau^-$ (30%) and $b\bar{b}$ (70%). From Ref. [47].

sky for DM searches in Sec. 5.

4. The Galactic Center

The Galactic Center has long been considered to be among the most promising targets for the detection of DM annihilation, particularly if the halo profile of the Milky Way is cusped in its inner volume [8, 40]. This has been complicated, however, by the recent discovery of astrophysical gamma-ray sources from the Galactic Center. Following an earlier claim by the WHIPPLE IACT [41], very high-energy gamma-rays from the Galactic Center have been detected by HESS [42], MAGIC [43], and CANGAROO-II [44]. This source is consistent with point-like emission and is located at $l = 359^\circ 56' 41.1'' \pm 6.4''$ (stat), $b = -0^\circ 2' 39.2'' \pm 5.9''$ (stat) with a systematic pointing error of $28''$ [45], coincident with the position of Sgr A*, the black hole constituting the dynamical center of the Milky Way. The spectrum of this source is well described by a power-law with a spectral index of $\alpha = 2.25 \pm 0.04$ (stat) ± 0.10 (syst) over the range of approximately 160 GeV to 20 TeV. Although speculations were initially made that this source could be the product of annihilations of very heavy (~ 10 to 50 TeV) DM particles [46], this interpretation is disfavored by the power-law form of the observed spectrum and the wide energy range over which it extends (see Fig. 2). The source of these gamma-rays is, instead, likely an astrophysical accelerator associated with our Galaxy's central supermassive black hole [48]. In recent analyses, this source has been treated as a background for DM searches [49, 50]. Given the presence of this background, the prospects for detecting DM annihilation products from the Galactic Center appear considerably less promising than they had a few years ago.

The main challenge involved in DM searches with FERMI will be to distinguish the signal from this and other backgrounds. This is made particularly difficult by




Figure 3. The projected exclusion limits at 95% confidence level from FERMI/GLAST (after ten years) on the WIMP annihilation cross section, as a function of the WIMP mass. The region above the dotted line is already excluded by EGRET [52]. The dashed and solid lines show the projections for FERMI/GLAST for an assumed isotropic diffuse background and the limit case where the astrophysical background has exactly the same angular distribution of the DM signal, respectively. In the left and right frames, the NFW and Moore *et al.* halo profiles have been adopted, respectively. Also shown are points representing a random scan of supersymmetric models. From Ref. [12].

our ignorance regarding the nature of these backgrounds. The Galactic Center is indeed a complex region of the sky at all wavelengths, the gamma-ray window being no exception [51].

An attempt to use angular and spectral information to separate DM annihilation products from these backgrounds was performed in Ref. [12], studying a $2^\circ \times 2^\circ$ region around the Galactic Center (see also the analysis of the FERMI/DM team in Ref. [6]). This study considered two known point-source backgrounds: a yet unidentified source detected by EGRET approximately 0.2° away from the dynamical center of our galaxy [52, 53], and the HESS source discussed above. In addition, a diffuse spectrum with a free power-law index was also included.

A multi-parameter χ^2 analysis of the simulated sky against models including a contribution from DM annihilation radiation yields the projected exclusion limits at the 95% confidence level shown in Fig. 3 (for ten years of collection time by FERMI). The left and right panels refer to the cases of an NFW profile and Moore *et al.* profile (replaced by a flat core within 10^{-2} pc of the Galactic Center to avoid a divergency), respectively. The WIMP is assumed to annihilate dominantly to $b\bar{b}$. The solid line in each frame represents the limit found if the diffuse background is assumed to be distributed isotropically, while the dashed line represents the conservative limit obtained if the diffuse background has the same angular distribution as the DM signal (*i.e.* the case in which angular information is not useful in disentangling the signal from the diffuse




Figure 4. The ability of FERMI/GLAST to measure the annihilation cross section and mass of dark matter after ten years of observation. A benchmark scenario with $m_X = 100$ GeV, $\langle\sigma v\rangle = 3 \times 10^{-26}$ cm³/s and an NFW halo profile has been used. The inner and outer contours in each frame represent the 2 and 3σ regions, respectively. In the left frame, the halo profile shape was treated as if it is known in advance. In the right frame, a marginalization over the inner slope of the profile was performed. From [12].

background). For values of $\langle\sigma v\rangle$ below the corresponding lines, a pure background model is expected to be consistent with the data. The fact that the limits are significantly stronger in the uniform background case is the manifestation of the improved sensitivity which can be achieved by an analysis including both energy and angular information. For comparison, in Fig. 3 it is also shown the region already excluded by EGRET [52] (above the dotted line) and the mass and cross section of neutralino models found in a random scan over supersymmetric parameters, as calculated using DarkSUSY [54]. As expected, many of the models cluster around $\langle\sigma v\rangle \sim 3 \times 10^{-26}$ cm³/s, the value required of a thermal relic annihilating via an S -wave amplitude. Each point shown represents a model which respects all direct collider constraint and generates a thermal DM abundance consistent with the observed DM density. In the scan the SUSY parameters varied were M_2 , $|\mu|$ and $m_{\tilde{q}}$ up to 2 TeV, m_A and $m_{\tilde{\tau}}$ up to 1 TeV and $\tan\beta$ up to 60. Also, the gaugino masses were assumed to evolve to a single unified scale, such that $M_1 \approx 0.5M_2$, $M_3 \approx 2.7M_2$.

Should gamma-rays be identified as having been produced in DM annihilations, such observations could then be used to measure the characteristics of the DM particle, including its mass, annihilation cross section and spatial distribution. Such determinations are an important step toward identifying the particle nature of DM. A calculation similar to the one leading to the results of Fig. 3 can be performed, this time including in the template a contribution from a fiducial model of DM annihilation, and asking the accuracy by which a reconstruction of its input properties is possible. In Fig. 4, the ability of FERMI to determine the WIMP mass and annihilation cross




Figure 5. The ability of FERMI/GLAST to measure the inner slope of the halo profile and the mass of dark matter particle (marginalizing over the annihilation cross section) after ten years of observation. Here, a benchmark scenario with $m_X = 100$ GeV, $\langle\sigma v\rangle = 3 \times 10^{-26}$ cm³/s and an NFW halo profile has been used. The inner and outer contours represent the 2 and 3 σ regions, respectively. From Ref. [12].

section for WIMPs distributed with an NFW halo profile is illustrated, for the case of an annihilation cross section of 3×10^{-26} cm³/s and a mass of 100 GeV. In each frame the projected 2 and 3 σ constraints on the input parameters are reported, assuming an isotropic diffuse background (in addition to background point sources). In the left frame, we treat the shape of the halo profile (NFW) as if it is known in advance. Of course, this is not a realistic assumption, and a less accurate determination of the WIMP mass must be expected in a more realistic treatment. In the right panel of Fig. 4 we report the results obtained marginalizing over the inner slope of the halo profile, γ . Without a prior on the inner halo slope, the constraint on the DM mass worsens by a factor ~ 2 .

If the spectrum and angular distribution of gamma-rays from DM annihilations in the Galactic Center region are sufficiently well measured, it will also be possible to measure the underlying DM distribution. Fig. 5 shows the results from the right frame of Fig. 4 in the $\{m_X, \gamma\}$ plane, marginalized over the annihilation cross section. In the above benchmark model, the inner slope of the halo profile can be determined at approximately the $\sim 10\%$ level.

A caveat is in order: in the above estimates one is assuming that a limited number of discrete sources will “contaminate” the inner Galactic Center angular bins, and/or that they could be removed effectively. Also we are extrapolating the source properties from lower and higher energies. FERMI itself will be used to determine the actual background(s) properties. Finally, the bounds are slightly optimistic, in the sense that we are considering the longest plausible lifetime of FERMI (ten years) and that further effects degrading the angular resolution or accounting for dead-time may loosen the constraints by a factor $\sim \mathcal{O}(1)$.

5. The Inner Halo

The emission of radiation per unit solid angle from DM annihilation is expected to be maximized at the Galactic Center. Yet, geometric factors and the presence of point-like and diffuse backgrounds make the choice of the optimal window size a non-trivial problem [6, 7, 8, 55, 56, 57]. Unless the DM halo is very cuspy towards the Galactic Center (say, cuspier than NFW), the optimal strategy for DM searches is never to focus on the inner sub-degree around the Galactic Center. Rather, a window size from 1° to 50° or more is preferred, where the precise optimal size depends on the angular distribution of the signal and backgrounds. Although the optimal shape of the window depends on the details of the signal and backgrounds (and could be optimized from morphological studies of the low-energy emission measured by FERMI), a circular annulus around the Galactic Center or a “rectangular” window in Galactic coordinates (with an inner rectangular mask) generally work fairly well. The improvement in the sensitivity reaches a factor of a few (especially for cored distributions), thus affecting the prospects for the detectability of specific models. This is illustrated in Fig. 6, where we plot the relative signal-to-noise as a function of the maximum galactic latitude, b_{\max} , for a Galactic region $0.4^\circ < |b| < b_{\max}$, $0^\circ < |l| < l_{\max} = b_{\max}$. The dot-dashed line refers to the continuum spectrum of a 100 GeV neutralino annihilating into W^+W^- with an NFW distribution. The solid lines refer to an hypothetical gamma-line emission for the benchmark “model I” in the inert Higgs doublet scenario, see Ref. [14].




Figure 6. The relative signal-to-noise as a function of the window size, b_{\max} , for a region $0.4^\circ < |b| < b_{\max}$, $0^\circ < |l| < l_{\max} = b_{\max}$ for the continuum spectrum of a 100 GeV neutralino annihilating into W^+W^- (dot-dashed line, NFW profile) and for the gamma-line emission for the benchmark “model I” in the inert Higgs doublet scenario, see Ref. [14]. From left to right, the solid lines refer to the Moore, NFW and Kravtsov profiles, see Table I. From Ref. [57].

Besides improving the prospects for detection, other advantages in focusing on an extended region around the Galactic Center include:

- Enabling an empirical determination of the dark matter profile slope outside of the region dominated by the gravitational potential of the central supermassive black hole.
- By comparing the emission from many angular regions, independent evidence can be obtained that it is the product of DM annihilation. In contrast, if the spectrum is found to vary with location, it is most likely the product of astrophysical backgrounds.
- By observing the angular distribution of the emission, it may be possible to constrain the quantity of dark matter substructure in the halo. The shape of the smooth, unresolved inner profile [21] could be studied, in addition to an anisotropy/multipole analysis [58, 59, 60]. This, in turn, may have important implications for DM cosmology; see the review by T. Bringmann in this issue.

6. Conclusions

In contrast to other probes of particle dark matter, the most challenging task for indirect dark matter searches is not to detect a few events, but to confidently identify those events as the products of dark matter annihilations. In particular, any signal must be separated from astrophysical backgrounds if it is to be reliably claimed to be a detection of dark matter. This is certainly true in the case of gamma-ray telescopes hoping to observe dark matter annihilations in the region of the Galactic Center. The discovery of a bright TeV source of astrophysical origin at the Galactic Center by IACTs has changed the prospects for such searches considerably. One strategy in light of this is to focus on different targets with lesser astrophysical background (such as dwarf galaxies). Another is to search for a signal in the “noisy” Galactic Center region, by taking advantage of the peculiar spectral and angular properties expected from dark matter annihilation products.

In this paper, we have reviewed the latter strategy, reporting on recent studies of the Galactic Center and inner-Galaxy. Given the characteristics of this search, the FERMI satellite is better suited than existing IACTs. FERMI’s LAT instrument will detect a number of astrophysical sources in the region of the sky around the Galactic Center, including the point sources previously identified by HESS and EGRET, and perhaps others. A diffuse gamma-ray background will also likely be present. Although predictions of FERMI’s sensitivity are unavoidably limited by our incomplete knowledge of these backgrounds, we have shown that the spectral and angular differences between the signal and backgrounds should be distinctive enough to be distinguished over a significant region of the parameter space, at least for a sufficiently cusped dark matter profile (NFW-like or steeper).

If dark matter annihilation products are identified by FERMI, then it may also be possible to measure or constrain the properties of the dark matter, including its mass,

annihilation cross section, and spatial distribution. It is unlikely that FERMI will determine the WIMP’s mass with high precision, however. For example, for the case of a 100 GeV WIMP with an annihilation cross section of $3 \times 10^{-26} \text{ cm}^3/\text{s}$, distributed with an NFW halo profile, the mass could be determined to lie within approximately 50-300 GeV. In the same benchmark model, the inner slope of the dark matter halo profile could be determined to $\sim 10\%$ precision. The combination of several indirect detection channels will be crucial to both confirm such a detection, and to best constrain the WIMP’s properties. Even limiting our attention to gamma-rays, it would be important to detect the emission from an extended region of the sky with FERMI, or to obtain an independent detection from other dark matter substructures with both FERMI and IACTs.

Acknowledgments

PS would like to thank the Galileo Galilei Institute for Theoretical Physics for the hospitality and the INFN for partial support during the completion of this work. DH is supported in part by the Fermi Research Alliance, LLC under Contract No. DE-AC02-07CH11359 with the US Department of Energy and by NASA grant NNX08AH34G.

Appendix A. The Case of Decaying Dark Matter

In the case of decaying dark matter, Eq. (1) is modified to

$$\Phi_\gamma = \frac{dN_\gamma}{dE_\gamma} \frac{\Gamma}{4\pi m_X} \int_{\text{los}} \rho(\ell, \Omega) d\ell, \quad (\text{A.1})$$

where Γ is the decay width (inverse lifetime) and the spectrum now refers to the photons generated in the decay process. Unlike with the cross section in the case of annihilating dark matter, we do not have any strong theoretical motivation for considering any particular lifetime for an unstable DM particle. In any case, arguments have been put forward justifying the typical range of the lifetimes needed for significant signatures in astrophysics with $\sim \text{TeV}$ mass particles and GUT-scale physics mediating the process (in analogy with the expected proton decay in GUTs), see e.g. [61]. From the phenomenological point of view, there are a couple of points worth mentioning regarding decaying DM candidates:

- I. The DM distribution and the role of substructures in particular is of little importance in determining the level of the signal.
- II. The angular distribution of the gamma-ray signal is very distinctive, and much flatter than the corresponding annihilation signal, as illustrated for a NFW profile in Fig. (A1).

Should gamma-rays be detected from DM, a comparison between the emission in the inner Galaxy and the emission at high latitude would immediately reveal the nature of the particle physics process (annihilation or decay) responsible for the emission [62]. Notice that this information is very difficult to extract with other cosmic ray probes.




Figure A1. The angular profile of the gamma-ray signal as function of the angle, θ , to the center of the galaxy for a NFW halo distribution for decaying DM (solid red line), compared to the case of self-annihilating DM (dashed blue line). Both signals have been normalized to their values at the galactic poles, $\theta = \pm 90^\circ$. The central cusp is regularized by assuming in both cases an angular resolution of 0.1° . From Ref. [62].

Appendix B. A Comment on the Factorization Assumption in Eq. (1)

Although the factorization between the particle physics term and astrophysical term in Eq. (1) is a useful approximation and valid in most practical cases, there are exceptions. More correctly, one should write

$$\Phi_\gamma(E_\gamma, \Omega) = \frac{dN_\gamma}{dE_\gamma}(E_\gamma) \frac{1}{8\pi m_X^2} \int_{\text{los}} \left[\int \sigma(v_r) v_r u(\mathbf{v}_r) d^3\mathbf{v}_r \right] \rho^2(\ell, \Omega) d\ell, \quad (\text{B.1})$$

where \mathbf{v}_r is the relative velocity between the two particles (with $v_r \equiv |\mathbf{v}_r|$) and $u(\mathbf{v}_r)$ its distribution function (not necessarily isotropic), whose integral over $d^3\mathbf{v}_r$ is normalized to unity. The factorization assumed in Eq. (1) holds only if the integral in square brackets—which is nothing but $\langle \sigma v \rangle$ —is independent of position. A sufficient condition for this is that $\sigma(v_r) v_r$ is velocity-independent. In general, the integral depends on the kinematical structure of the halo via the position-dependent velocity dispersion, anisotropy, etc. Then, both the astrophysical distribution of the DM and the particle physics contribute in determining the angular shape of the signal.

One case in which the factorization is not valid can be found when the WIMP annihilations mainly through a P -wave process, such that $\sigma(v_r) v_r \propto v_r^2$ [63]. This is a largely academic case, however, since whenever P -wave annihilation is dominant the gamma-ray signal is expected to be suppressed. More interesting is the case in which non-perturbative processes lead to large “Sommerfeld enhancements” to the annihilation cross section at low velocities [64]. This effect can be thought of as the distortion of the wave-function due to a relatively long-range attraction between the WIMPs. In this

case, a further steepening of the signal towards the Galactic Center is expected, which in turn should ease the detection of gamma-rays from the inner Galaxy [65].

References

- [1] <http://www-glast.slac.stanford.edu/>
- [2] <http://www.mpi-hd.mpg.de/hfm/HESS/HESS.html>
- [3] <http://www.magic.iac.es/>
- [4] <http://veritas.sao.arizona.edu/>
- [5] <http://icrhp9.icrr.u-tokyo.ac.jp/>
- [6] E. A. Baltz *et al.*, JCAP **0807**, 013 (2008) [arXiv:0806.2911 [astro-ph]].
- [7] J. Buckley *et al.*, arXiv:0812.0795 [astro-ph].
- [8] L. Bergstrom, P. Ullio and J. H. Buckley, Astropart. Phys. **9**, 137 (1998) [arXiv:astro-ph/9712318].
- [9] G. Bertone, D. Hooper and J. Silk, Phys. Rept. **405**, 279 (2005) [arXiv:hep-ph/0404175].
- [10] T. Moroi and L. Randall, Nucl. Phys. B **570**, 455 (2000) [arXiv:hep-ph/9906527].
- [11] G. Jungman, M. Kamionkowski and K. Griest, Phys. Rept. **267**, 195 (1996) [hep-ph/9506380].
- [12] S. Dodelson, D. Hooper and P. D. Serpico, Phys. Rev. D **77**, 063512 (2008) [arXiv:0711.4621 [astro-ph]].
- [13] L. Bergstrom and P. Ullio, Nucl. Phys. B **504**, 27 (1997) [hep-ph/9706232]; P. Ullio and L. Bergstrom, Phys. Rev. D **57**, 1962 (1998) [hep-ph/9707333].
- [14] M. Gustafsson, E. Lundstrom, L. Bergstrom and J. Edsjo, Phys. Rev. Lett. **99**, 041301 (2007) [arXiv:astro-ph/0703512].
- [15] A. Birkedal, K. T. Matchev, M. Perelstein and A. Spray, arXiv:hep-ph/0507194.
- [16] T. Bringmann, L. Bergstrom and J. Edsjo, JHEP **0801**, 049 (2008) [arXiv:0710.3169 [hep-ph]].
- [17] M. Kachelriess and P. D. Serpico, Phys. Rev. D **76**, 063516 (2007) [arXiv:0707.0209 [hep-ph]].
- [18] N. F. Bell, J. B. Dent, T. D. Jacques and T. J. Weiler, Phys. Rev. D **78**, 083540 (2008) [arXiv:0805.3423 [hep-ph]].
- [19] J. J. Binney and N. W. Evans, Mon. Not. Roy. Astron. Soc. **327**, L27 (2001) [arXiv:astro-ph/0108505].
- [20] A. Klypin, H. Zhao and R. S. Somerville, Astrophys. J. **573**, 597 (2002) [arXiv:astro-ph/0110390].
- [21] D. Hooper and P. D. Serpico, JCAP **0706**, 013 (2007) [astro-ph/0702328].
- [22] S. Gillessen, F. Eisenhauer, S. Trippe, T. Alexander, R. Genzel, F. Martins and T. Ott, arXiv:0810.4674 [astro-ph].
- [23] J. F. Navarro *et al.*, Astrophys. J. **462** 563 (1996).
- [24] B. Moore *et al.*, Phys. Rev. D **64** 063508 (2001).
- [25] A. V. Kravtsov, A. A. Klypin, J. S. Bullock and J. R. Primack, Astrophys. J. **502**, 48 (1998) [astro-ph/9708176].
- [26] C. Power *et al.*, Mon. Not. Roy. Astron. Soc. **338**, 14 (2003) [astro-ph/0201544].
- [27] J. F. Navarro *et al.*, Mon. Not. Roy. Astron. Soc. **349**, 1039 (2004) [astro-ph/0311231].
- [28] D. Reed *et al.*, Mon. Not. Roy. Astron. Soc. **357**, 82 (2005) [astro-ph/0312544].
- [29] D. Merritt, J. F. Navarro, A. Ludlow and A. Jenkins, Astrophys. J. **624**, L85 (2005) [astro-ph/0502515].
- [30] J. F. Navarro *et al.*, arXiv:0810.1522 [astro-ph].
- [31] J. Einasto, Trudy Inst. Astroz. Alma-Ata, **51**, 87 (1965)
- [32] O. Y. Gnedin, A. V. Kravtsov, A. A. Klypin and D. Nagai, Astrophys. J. **616**, 16 (2004) [astro-ph/0406247];
- [33] M. Gustafsson, M. Fairbairn and J. Sommer-Larsen, Phys. Rev. D **74**, 123522 (2006) [arXiv:astro-ph/0608634].
- [34] J. I. Read, L. Mayer, A. M. Brooks, F. Governato and G. Lake, arXiv:0902.0009 [astro-ph.GA].
- [35] F. Prada, A. Klypin, J. Flix, M. Martinez and E. Simonneau, Phys. Rev. Lett. **93**, 241301 (2004) [arXiv:astro-ph/0401512]. G. Bertone and D. Merritt, Mod. Phys. Lett. A **20**, 1021

- (2005) [astro-ph/0504422]; G. Bertone and D. Merritt, *Phys. Rev. D* **72**, 103502 (2005) [astro-ph/0501555].
- [36] J. A. Sellwood, arXiv:0807.1973 [astro-ph].
- [37] P. Gondolo and J. Silk, *Phys. Rev. Lett.* **83**, 1719 (1999) [astro-ph/9906391]; P. Ullio, H. Zhao and M. Kamionkowski, *Phys. Rev. D* **64**, 043504 (2001) [astro-ph/0101481]; G. Bertone, G. Sigl and J. Silk, *Mon. Not. Roy. Astron. Soc.* **337**, 98 (2002) [astro-ph/0203488].
- [38] M. Fornasa and G. Bertone, *Int. J. Mod. Phys. D* **17**, 1125 (2008) [arXiv:0711.3148 [astro-ph]].
- [39] H. Yuksel, S. Horiuchi, J. F. Beacom and S. Ando, *Phys. Rev. D* **76**, 123506 (2007) [arXiv:0707.0196 [astro-ph]].
- [40] L. Bergstrom, J. Edsjo and P. Ullio, *Phys. Rev. Lett.* **87**, 251301 (2001) [astro-ph/0105048]; V. Berezhinsky, A. Bottino and G. Mignola, *Phys. Lett. B* **325**, 136 (1994) [hep-ph/9402215]; A. Cesarini, F. Fucito, A. Lionetto, A. Morselli and P. Ullio, *Astropart. Phys.* **21**, 267 (2004) [astro-ph/0305075]; P. Ullio, L. Bergstrom, J. Edsjo and C. G. Lacey, *Phys. Rev. D* **66**, 123502 (2002) [astro-ph/0207125].
- [41] K. Kosack *et al.* [The VERITAS Collaboration], *Astrophys. J.* **608**, L97 (2004) [astro-ph/0403422].
- [42] F. Aharonian *et al.* [The HESS Collaboration], *Astron. Astrophys.* **425**, L13 (2004) [astro-ph/0408145].
- [43] J. Albert *et al.* [MAGIC Collaboration], *Astrophys. J.* **638**, L101 (2006) [astro-ph/0512469].
- [44] K. Tsuchiya *et al.* [CANGAROO-II Collaboration], *Astrophys. J.* **606**, L115 (2004) [astro-ph/0403592].
- [45] C. van Eldik, O. Bolz, I. Braun, G. Hermann, J. Hinton and W. Hofmann [for the HESS Collaboration], arXiv:0709.3729 [astro-ph].
- [46] D. Hooper and J. March-Russell, *Phys. Lett. B* **608**, 17 (2005) [hep-ph/0412048]; D. Hooper, I. de la Calle Perez, J. Silk, F. Ferrer and S. Sarkar, *JCAP* **0409**, 002 (2004) [astro-ph/0404205]; S. Profumo, *Phys. Rev. D* **72**, 103521 (2005) [astro-ph/0508628]; L. Bergstrom, T. Bringmann, M. Eriksson and M. Gustafsson, *Phys. Rev. Lett.* **94**, 131301 (2005) [astro-ph/0410359].
- [47] F. Aharonian *et al.* [H.E.S.S. Collaboration], *Phys. Rev. Lett.* **97**, 221102 (2006) [Erratum-ibid. **97**, 249901 (2006)] [arXiv:astro-ph/0610509].
- [48] F. Aharonian and A. Neronov, *Astrophys. J.* **619**, 306 (2005) [astro-ph/0408303]; astro-ph/0503354; *AIP Conf. Proc.* **745**, 409 (2005); A. Atoyan and C. D. Dermer, *Astrophys. J.* **617**, L123 (2004) [astro-ph/0410243].
- [49] G. Zaharijas and D. Hooper, *Phys. Rev. D* **73**, 103501 (2006) [arXiv:astro-ph/0603540].
- [50] D. Hooper, G. Zaharijas, D. P. Finkbeiner and G. Dobler, *Phys. Rev. D* **77**, 043511 (2008) [arXiv:0709.3114 [astro-ph]].
- [51] C. van Eldik, arXiv:0811.0931 [astro-ph].
- [52] D. Hooper and B. L. Dingus, *Phys. Rev. D* **70**, 113007 (2004) [astro-ph/0210617]; D. Hooper and B. Dingus, *Proc. of the 34th COSPAR Scientific Assembly, Houston, Texas* (2002), astro-ph/0212509.
- [53] M. Pohl, *Astrophys. J.* **626**, 174 (2005).
- [54] P. Gondolo, J. Edsjo, P. Ullio, L. Bergstrom, M. Schelke and E. A. Baltz, *JCAP* **0407**, 008 (2004) [astro-ph/0406204].
- [55] F. Stoehr, S. D. M. White, V. Springel, G. Tormen and N. Yoshida, *Mon. Not. Roy. Astron. Soc.* **345**, 1313 (2003) [astro-ph/0307026].
- [56] N. W. Evans, F. Ferrer and S. Sarkar, *Phys. Rev. D* **69**, 123501 (2004) [astro-ph/0311145];
- [57] P. D. Serpico and G. Zaharijas, *Astropart. Phys.* **29**, 380 (2008) [arXiv:0802.3245 [astro-ph]].
- [58] J. M. Siegal-Gaskins, *JCAP* **0810**, 040 (2008) [arXiv:0807.1328 [astro-ph]].
- [59] S. K. Lee, S. Ando and M. Kamionkowski, arXiv:0810.1284 [astro-ph].
- [60] M. Fornasa, L. Pieri, G. Bertone and E. Branchini, arXiv:0901.2921 [astro-ph].
- [61] A. Arvanitaki, S. Dimopoulos, S. Dubovsky, P. W. Graham, R. Harnik and S. Rajendran, arXiv:0812.2075 [hep-ph].
- [62] G. Bertone, W. Buchmuller, L. Covi and A. Ibarra, *JCAP* **0711**, 003 (2007) [arXiv:0709.2299].

- [63] M. A. Amin and T. Wizansky, Phys. Rev. D **77**, 123510 (2008) [arXiv:0710.5517 [astro-ph]].
- [64] J. Hisano, S. Matsumoto, M. M. Nojiri and O. Saito, Phys. Rev. D **71**, 063528 (2005) [arXiv:hep-ph/0412403].
- [65] B. Robertson and A. Zentner, arXiv:0902.0362 [astro-ph.CO].