



Neotropical Ichthyology

OPEN ACCESS



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under
Creative Commons CC-BY 4.0

© 2024 The Authors.
Diversity and Distributions Published by SBI



Official Journal of the
Sociedade Brasileira de Ictiologia

FIGURE S4 | Phylogenetic relationships among Dasyatinae, Neotrygoninae, and Urogymninae representatives based on a Bayesian Inference on an alignment of 11,471 base pairs with 135 sequences. For each node, posterior probabilities are given. Genus *Hypanus* in red, subfamily Dasyatinae in blue (except *Hypanus*), subfamily Neotrygoninae in green, and subfamily Urogymninae in orange. Download the image here: <https://www.ni.bio.br/content/v22n1/1982-0224-2023-0046/supplementary/1982-0224-ni-22-01-e230046-gfs4.zip>

HOW TO CITE THIS ARTICLE

- Petean FF, Yang L, Corrigan S, Lima SMQ, Naylor GJP. How many lineages are there of the stingrays genus *Hypanus* (Myliobatiformes: Dasyatidae) and why does it matter? *Neotrop Ichthyol*. 2024; 22(1):e230046. <https://doi.org/10.1590/1982-0224-2023-0046>