

Norm-Based Capacity Control in Neural Networks

Behnam Neyshabur

BNEYSHABUR@TTIC.EDU

Ryota Tomioka

TOMIOKA@TTIC.EDU

Nathan Srebro

NATI@TTIC.EDU

Toyota Technological Institute at Chicago, Chicago, IL 60637, USA

Abstract

We investigate the capacity, convexity and characterization of a general family of norm-constrained feed-forward networks.

Keywords: Feed-forward neural networks, deep learning, scale-sensitive capacity control

1. Introduction

The statistical complexity, or capacity, of *unregularized* feed-forward neural networks, as a function of the network size and depth, is fairly well understood. With hard-threshold activations, the VC-dimension, and hence sample complexity, of the class of functions realizable with a feed-forward network is equal, up to logarithmic factors, to the number of edges in the network (Anthony and Bartlett, 2009; Shalev-Shwartz and Ben-David, 2014), corresponding to the number of parameters. With continuous activation functions the VC-dimension could be higher, but is fairly well understood and is still controlled by the size and depth of the network.¹

But feedforward networks are often trained with some kind of explicit or implicit regularization, such as weight decay, early stopping, “max regularization”, or more exotic regularization such as drop-outs. What is the effect of such regularization on the induced hypothesis class and its capacity?

For linear prediction (a one-layer feed-forward network) we know that using regularization the capacity of the class can be bounded only in terms of the norms, with no (or a very weak) dependence on the number of edges (i.e. the input dimensionality or number of linear coefficients). E.g., we understand very well how the capacity of ℓ_2 -regularized linear predictors can be bounded in terms of the norm alone (when the norm of the data is also bounded), even in infinite dimension.

A central question we ask is: can we bound the capacity of feed-forward network in terms of norm-based regularization alone, without relying on network size and even if the network size (number of nodes or edges) is unbounded or infinite? What type of regularizers admit such capacity control? And how does the capacity behave as a function of the norm, and perhaps other network parameters such as depth?

Beyond the central question of capacity control, we also analyze the convexity of the resulting hypothesis class—unlike unregularized size-controlled feed-forward networks, infinite magnitude-controlled networks have the potential of yielding convex hypothesis classes (this is the case, e.g.,

1. Using weights with very high precision and vastly different magnitudes it is possible to shatter a number of points quadratic in the number of edges when activations such as the sigmoid, ramp or hinge are used (Shalev-Shwartz and Ben-David, 2014, Chapter 20.4). But even with such activations, the VC dimension can still be bounded by the size and depth (Bartlett, 1998; Anthony and Bartlett, 2009; Shalev-Shwartz and Ben-David, 2014).

when we move from rank-based control on matrices, which limits the number of parameters to magnitude based control with the trace-norm or max-norm). A convex class might be easier to optimize over and might be convenient in other ways.

In this paper we focus on networks with rectified linear units and two natural types of norm regularization: bounding the norm of the incoming weights of each unit (per-unit regularization) and bounding the overall norm of all the weights in the system jointly (overall regularization, e.g. limiting the overall sum of the magnitudes, or square magnitudes, in the system). We generalize both of these with a single notion of group-norm regularization: we take the ℓ_p norm over the weights in each unit and then the ℓ_q norm over units. In Section 3 we present this regularizer and obtain a tight understanding of when it provides for size-independent capacity control and a characterization of when it induces convexity. We then apply these generic results to per-unit regularization (Section 4) and overall regularization (Section 5), noting also other forms of regularization that are equivalent to these two. In particular, we show how per-unit regularization is equivalent to a novel path-based regularizer and how overall ℓ_2 regularization for two-layer networks is equivalent to so-called “convex neural networks” (Bengio et al., 2005). In terms of capacity control, we show that per-unit regularization allows size-independent capacity-control only with a per-unit ℓ_1 -norm, and that overall ℓ_p regularization allows for size-independent capacity control only when $p \leq 2$, even if the depth is bounded. In any case, even if we bound the sum of all magnitudes in the system, we show that an exponential dependence on the depth is unavoidable.

As far as we are aware, prior work on size-independent capacity control for feed-forward networks considered only per-unit ℓ_1 regularization, and per-unit ℓ_2 regularization for two-layered networks (see discussion and references at the beginning of Section 4). Here, we extend the scope significantly, and provide a broad characterization of the types of regularization possible and their properties. In particular, we consider overall norm regularization, which is perhaps the most natural form of regularization used in practice (e.g. in the form of weight decay). We hope our study will be useful in thinking about, analyzing and designing learning methods using feed-forward networks. Another motivation for us is that complexity of large-scale optimization is often related to scale-based, not dimension-based complexity. Understanding when the scale-based complexity depends exponentially on the depth of a network might help shed light on understanding the difficulties in optimizing deep networks.

2. Preliminaries: Feedforward Neural Networks

A feedforward neural network that computes a function $f : \mathbb{R}^D \rightarrow \mathbb{R}$ is specified by a directed acyclic graph (DAG) $G(V, E)$ with D special “input nodes” $v_{\text{in}}[1], \dots, v_{\text{in}}[D] \in V$ with no incoming edges and a special “output node” $v_{\text{out}} \in V$ with no outgoing edges, weights $w : E \rightarrow \mathbb{R}$ on the edges, and an *activation function* $\sigma : \mathbb{R} \rightarrow \mathbb{R}$.

Given an input $x \in \mathbb{R}^D$, the output values of the input units are set to the coordinates of x , $o(v_{\text{in}}[i]) = x[i]$ (we might want to also add a special “bias” node with $o(v_{\text{in}}[0]) = 1$, or just rely on the inputs having a fixed “bias coordinate”), the output value of internal nodes (all nodes except the input and output nodes) are defined according to the forward propagation equation:

$$o(v) = \sigma \left(\sum_{(u \rightarrow v) \in E} w(u \rightarrow v) o(u) \right), \quad (1)$$

and the output value of the output unit is defined as $o(v_{\text{out}}) = \sum_{(u \rightarrow v_{\text{out}}) \in E} w(u \rightarrow v_{\text{out}})o(u)$. The network is then said to compute the function $f_{G,w,\sigma}(x) = o(v_{\text{out}})$. Given a graphs G and activation function σ , we can consider the hypothesis class of functions $\mathcal{N}^{G,\sigma} = \{f_{G,w,\sigma} : \mathbb{R}^D \rightarrow \mathbb{R} \mid w : E \rightarrow \mathbb{R}\}$ computable using some setting of the weights.

We will refer to the *size* of the network, which is the overall number of edges $|E|$, the *depth* d of the network, which is the length of the longest directed path in G , and the *in-degree* (or width) H of a network, which is the maximum in-degree of a vertex in G .

A special case of feedforward neural networks are layered fully connected networks where vertices are partitioned into layers and there is a directed edge from every vertex in layer i to every vertex in layer $i + 1$. We index the layers from the first layer, $i = 1$ whose inputs are the input nodes, up to the last layer $i = d$ which contains the single output node—the number of layers is thus equal to the depth and the in-degree is the maximal layer size. We denote by $\text{layer}(d, H)$ the layered fully connected network with d layers and H nodes per layer (except the output layer that has a single node), and also allow $H = \infty$. We will also use the shorthand $\mathcal{N}^{d,H,\sigma} = \mathcal{N}^{\text{layer}(d,H),\sigma}$ and $\mathcal{N}^{d,\sigma} = \mathcal{N}^{\text{layer}(d,\infty),\sigma}$.

Layered networks can be parametrized by a sequence of matrices $W_1 \in \mathbb{R}^{H \times D}, W_2, W_3, \dots, W_{d-1} \in \mathbb{R}^{H \times H}, W_d \in \mathbb{R}^{1 \times H}$ where the row $W_i[j, :]$ contains the input weights to unit j in layer i , and

$$f_W(x) = W_d \sigma(W_{d-1} \sigma(W_{d-2} (\dots \sigma(W_1 x))))), \quad (2)$$

where σ is applied element-wise.

We will focus mostly on the hinge, or RELU (REctified Linear Unit) activation, which is currently in popular use (Nair and Hinton, 2010; Bordes and Bengio, 2011; Zeiler et al., 2013), $\sigma_{\text{RELU}}(z) = [z]_+ = \max(z, 0)$. When the activation will not be specified, we will implicitly be referring to the RELU. The RELU has several convenient properties which we will exploit, some of them shared with other activation functions:

Lipshitz The hinge is Lipschitz continuous with Lipschitz constant one. This property is also shared by the sigmoid and the ramp activation $\sigma(z) = \min(\max(0, z), 1)$.

Idempotency The hinge is idempotent, i.e. $\sigma_{\text{RELU}}(\sigma_{\text{RELU}}(z)) = \sigma_{\text{RELU}}(z)$. This property is also shared by the ramp and hard threshold activations.

Non-Negative Homogeneity For a non-negative scalar $c \geq 0$ and any input $z \in \mathbb{R}$ we have $\sigma_{\text{RELU}}(c \cdot z) = c \cdot \sigma_{\text{RELU}}(z)$. This property is important as it allows us to scale the incoming weights to a unit by $c > 0$ and scale the outgoing edges by $1/c$ without changing the the function computed by the network. For layered graphs, this means we can scale W_i by c and compensate by scaling W_{i+1} by $1/c$.

We will consider various measures $\alpha(w)$ of the magnitude of the weights $w(\cdot)$. Such a measure induces a complexity measure on functions $f \in \mathcal{N}^{G,\sigma}$ defined by $\alpha^{G,\sigma}(f) = \inf_{f_{G,w,\sigma}=f} \alpha(w)$. The sublevel sets of the complexity measure $\alpha^{G,\sigma}$ form a family of hypothesis classes $\mathcal{N}_{\alpha \leq a}^{G,\sigma} = \{f \in \mathcal{N}^{G,\sigma} \mid \alpha^{G,\sigma}(f) \leq a\}$. Again we will use the shorthand $\alpha^{d,H,\sigma}$ and $\alpha^{d,\sigma}$ when referring to layered graphs $\text{layer}(d, H)$ and $\text{layer}(d, \infty)$ respectively, and frequently drop σ when RELU is implicitly meant.

For binary function $g : \{\pm 1\}^D \rightarrow \pm 1$ we say that g is realized by f with unit margin if $\forall_x f(x)g(x) \geq 1$. A set of points S is shattered with unit margin by a hypothesis class \mathcal{N} if all $g : S \rightarrow \pm 1$ can be realized with unit margin by some $f \in \mathcal{N}$.

3. Group Norm Regularization

Considering the grouping of weights going into each node of the network, we will consider the following generic group-norm type regularizer, parametrized by $1 \leq p, q \leq \infty$:

$$\mu_{p,q}(w) = \left(\sum_{v \in V} \left(\sum_{(u \rightarrow v) \in E} |w(u \rightarrow v)|^p \right)^{q/p} \right)^{1/q}. \quad (3)$$

Here and elsewhere we allow $q = \infty$ with the usual conventions that $(\sum z_i^q)^{1/q} = \sup z_i$ and $1/q = 0$ when it appears in other contexts. When $q = \infty$ the group regularizer (3) imposes a per-unit regularization, where we constrain the norm of the incoming weights of each unit separately, and when $q = p$ the regularizer (3) is an ‘‘overall’’ weight regularizer, constraining the overall norm of all weights in the system. E.g., when $q = p = 1$ we are paying for the sum of all magnitudes of weights in the network, and $q = p = 2$ corresponds to overall weight-decay where we pay for the sum of square magnitudes of all weights (i.e. the overall Euclidean norm of the weights).

For a layered graph, we have:

$$\begin{aligned} \mu_{p,q}(W) &= \left(\sum_{k=1}^d \sum_{i=1}^H \left(\sum_{j=1}^H |W_k[i, j]|^p \right)^{q/p} \right)^{1/q} = d^{1/q} \left(\frac{1}{d} \sum_{k=1}^d \|W_k\|_{p,q}^q \right)^{1/q} \\ &\geq d^{1/q} \left(\prod_{k=1}^d \|W_k\|_{p,q} \right)^{1/d} \triangleq d^{1/q} \sqrt[d]{\gamma_{p,q}(W)} \end{aligned} \quad (4)$$

where $\gamma_{p,q}(W) = \prod_{k=1}^d \|W_k\|_{p,q}$ aggregates the layers by multiplication instead of summation. The inequality (4) holds regardless of the activation function, and so for any σ we have:

$$\gamma_{p,q}^{d,H,\sigma}(f) \leq \left(\frac{\mu_{p,q}^{d,H,\sigma}(f)}{d^{1/q}} \right)^d. \quad (5)$$

But due to the homogeneity of the RELU activation, when this activation is used we can always balance the norm between the different layers without changing the computed function so as to achieve equality in (4):

Claim 1 For any $f \in \mathcal{N}^{d,H,\sigma_{\text{RELU}}}$, $\mu_{p,q}^{d,H,\sigma_{\text{RELU}}}(f) = d^{1/q} \sqrt[d]{\gamma_{p,q}^{d,H,\sigma_{\text{RELU}}}(f)}$.

Proof Let W be weights that realizes f and are optimal with respect to $\gamma_{p,q}$; i.e. $\gamma_{p,q}(W) = \gamma_{p,q}^{d,H}(f)$. Let $\widetilde{W}_k = \sqrt[d]{\gamma_{p,q}(W)} W_k / \|W_k\|_{p,q}$, and observe that they also realize f . We now have:

$$\mu_{p,q}^{d,H}(f) \leq \mu_{p,q}(\widetilde{W}) = \left(\sum_{k=1}^d \|\widetilde{W}_k\|_{p,q}^q \right)^{1/q} = \left(d \left(\gamma_{p,q}(W) \right)^{q/d} \right)^{1/q} = d^{1/q} \sqrt[d]{\gamma_{p,q}^{d,H,\sigma_{\text{RELU}}}(f)}$$

which together with (4) completes the proof. \blacksquare

The two measures are therefore equivalent when we use RELUs, and define the same level sets, or family of hypothesis classes, which we refer to simply as $\mathcal{N}_{p,q}^{d,H}$. In the remainder of this Section, we investigate convexity and generalization properties of these hypothesis classes.

3.1. Generalization and Capacity

In order to understand the effect of the norm on the sample complexity, we bound the Rademacher complexity of the classes $\mathcal{N}_{p,q}^{d,H}$. Recall that the Rademacher Complexity is a measure of the capacity of a hypothesis class on a specific sample, which can be used to bound the difference between empirical and expected error, and thus the excess generalization error of empirical risk minimization (see, e.g., [Bartlett and Mendelson \(2003\)](#) for a complete treatment, and [Appendix A](#) for the exact definitions we use). In particular, the Rademacher complexity typically scales as $\sqrt{C/m}$, which corresponds to a sample complexity of $O(C/\epsilon^2)$, where m is the sample size and C is the effective measure of capacity of the hypothesis class.

Theorem 1 *For any $d, q \geq 1$, any $1 \leq p < \infty$ and any set $S = \{x_1, \dots, x_m\} \subseteq \mathbb{R}^D$:*

$$\begin{aligned} \mathcal{R}_m(\mathcal{N}_{\gamma_{p,q} \leq \gamma}^{d,H,\sigma_{\text{RELU}}}) &\leq \gamma \left(2H^{\lceil \frac{1}{p^*} - \frac{1}{q} \rceil_+}\right)^{(d-1)} \mathcal{R}_{m,p,D}^{\text{linear}} \\ &\leq \sqrt{\frac{\gamma^2 \left(2H^{\lceil \frac{1}{p^*} - \frac{1}{q} \rceil_+}\right)^{2(d-1)} \min\{p^*, 4 \log(2D)\} \max_i \|x_i\|_{p^*}^2}{m}} \end{aligned}$$

and so:

$$\begin{aligned} \mathcal{R}_m(\mathcal{N}_{\mu_{p,q} \leq \mu}^{d,H,\sigma_{\text{RELU}}}) &\leq \mu^d \left(2H^{\lceil \frac{1}{p^*} - \frac{1}{q} \rceil_+} / \sqrt[q]{d}\right)^{(d-1)} \mathcal{R}_{m,p,D}^{\text{linear}} \\ &\leq \sqrt{\frac{\mu^{2d} \left(2H^{\lceil \frac{1}{p^*} - \frac{1}{q} \rceil_+} / \sqrt[q]{d}\right)^{2(d-1)} \min\{p^*, 4 \log(2D)\} \max_i \|x_i\|_{p^*}^2}{m}} \end{aligned}$$

where the second inequalities hold only if $1 \leq p \leq 2$, $\mathcal{R}_{m,p,D}^{\text{linear}}$ is the Rademacher complexity of D -dimensional linear predictors with unit ℓ_p norm with respect to a set of m samples and p^* is such that $\frac{1}{p^*} + \frac{1}{p} = 1$.

Proof sketch We prove the bound by induction, showing that for any $q, d > 1$ and $1 \leq p < \infty$,

$$\mathcal{R}_m(\mathcal{N}_{\gamma_{p,q} \leq \gamma}^{d,H,\sigma_{\text{RELU}}}) \leq 2H^{\lceil \frac{1}{p^*} - \frac{1}{q} \rceil_+} \mathcal{R}_m(\mathcal{N}_{\gamma_{p,q} \leq \gamma}^{d-1,H,\sigma_{\text{RELU}}}).$$

The intuition is that when $p^* < q$, the Rademacher complexity increases by simply distributing the weights among neurons and if $p^* \geq q$ then the supremum is attained when the output neuron is connected to a neuron with highest Rademacher complexity in the lower layer and all other weights in the top layer are set to zero. For a complete proof, see [Appendix A](#). \blacksquare

Note that for $2 \leq p < \infty$, the bound on the Rademacher complexity scales with $m^{\frac{1}{p}}$ (see [section A.1](#) in appendix) because:

$$\mathcal{R}_{m,p,D}^{\text{linear}} \leq \frac{\sqrt{2} \|X\|_{2,p^*}}{m} \leq \frac{\sqrt{2} \max_i \|x_i\|_{p^*}}{m^{\frac{1}{p}}} \quad (6)$$

The bound in [Theorem 1](#) depends on both the magnitude of the weights, as captured by $\mu_{p,q}(W)$ or $\gamma_{p,q}(W)$, and also on the width H of the network (the number of nodes in each layer). However, the dependence on the width H disappears, and the bound depends only on the magnitude, as long

as $q \leq p^*$ (i.e. $1/p + 1/q \geq 1$). This happens, e.g., for overall ℓ_1 and ℓ_2 regularization, for per-unit ℓ_1 regularization, and whenever $1/p + 1/q = 1$. In such cases, we can omit the size constraint and state the theorem for an infinite-width layered network (i.e. a network with an infinitely countable number of units, when the number of units is allowed to be as large as needed):

Corollary 2 *For any $d \geq 1$, $1 \leq p < \infty$ and $1 \leq q \leq p^* = p/(p-1)$, and any set $S = \{x_1, \dots, x_m\} \subseteq \mathbb{R}^D$,*

$$\mathcal{R}_m(\mathcal{N}_{\gamma_{p,q} \leq \gamma}^{d,H,\sigma_{\text{RELU}}}) \leq \gamma 2^{(d-1)} \mathcal{R}_{m,p,D}^{\text{linear}} \leq \sqrt{\frac{\gamma^2 \left(2H^{\lceil \frac{1}{p^*} - \frac{1}{q} \rceil +}\right)^{2(d-1)} \min\{p^*, 4 \log(2D)\} \max_i \|x_i\|_{p^*}^2}{m}}$$

and so:

$$\mathcal{R}_m(\mathcal{N}_{\mu_{p,q} \leq \mu}^{d,H,\sigma_{\text{RELU}}}) \leq \left(2\mu/\sqrt[q]{d}\right)^d \mathcal{R}_{m,p,D}^{\text{linear}} \leq \sqrt{\frac{\left(2\mu/\sqrt[q]{d}\right)^{2d} \min\{p^*, 4 \log(2D)\} \max_i \|x_i\|_{p^*}^2}{m}}$$

where the second inequalities hold only if $1 \leq p \leq 2$ and $\mathcal{R}_{m,p,D}^{\text{linear}}$ is the Rademacher complexity of D -dimensional linear predictors with unit ℓ_p norm with respect to a set of m samples.

3.2. Tightness

We next investigate the tightness of the complexity bound in Theorem 1, and show that when $1/p + 1/q < 1$ the dependence on the width H is indeed unavoidable. We show not only that the bound on the Rademacher complexity is tight, but that the implied bound on the sample complexity is tight, even for binary classification with a margin over binary inputs. To do this, we show how we can shatter the $m = 2^D$ points $\{\pm 1\}^D$ using a network with small group-norm:

Theorem 3 *For any $p, q \geq 1$ (and $1/p^* + 1/p = 1$) and any depth $d \geq 2$, the $m = 2^D$ points $\{\pm 1\}^D$ can be shattered with unit margin by $\mathcal{N}_{\gamma_{p,q} \leq \gamma}^{d,H}$ with:*

$$\gamma \leq D^{1/p} m^{1/p+1/q} H^{-(d-2)[1/p^*-1/q]_+}$$

The proof is given in Appendix B. To understand this lower bound, first consider the bound without the dependence on the width H . We have that for any depth $d \geq 2$, $\gamma \leq m^r D = m^r \log m$ (since $1/p \leq 1$ always) where $r = 1/p + 1/q \leq 2$. This means that for any depth $d \geq 2$ and any p, q the sample complexity of learning the class scales as $m = \Omega(\gamma^{1/r} / \log \gamma) \geq \tilde{\Omega}(\sqrt{\gamma})$. This shows a polynomial dependence on γ , though with a lower exponent than the γ^2 (or higher for $p > 2$) dependence in Theorem 1. Still, if we now consider the complexity control as a function of $\mu_{p,q}$ we get a sample complexity of at least $\Omega(\mu^{d/2} / \log \mu)$, establishing that if we control the group-norm as in (3), we cannot avoid a sample complexity which depends exponentially on the depth. Note that in our construction, all other factors in Theorem 1, namely $\max_i \|x_i\|$ and $\log D$, are logarithmic (or double-logarithmic) in m .

Next we consider the dependence on the width H when $1/p + 1/q < 1$. Here we have to use depth $d \geq 3$, and we see that indeed as the width H and depth d increase, the magnitude control γ can decrease as $H^{(1/p^*-1/q)(d-2)}$ without decreasing the capacity, matching Theorem 1 up to an offset of 2 on the depth. In particular, we see that in this regime we can shatter an arbitrarily large number of points with arbitrarily low γ by using enough hidden units, and so the capacity of $\mathcal{N}_{p,q}^d$ is indeed infinite and it cannot ensure any generalization.

3.3. Convexity

Finally we establish a sufficient condition for the hypothesis classes $\mathcal{N}_{p,q}^d$ to be convex. We are referring to convexity of the functions in the $\mathcal{N}_{p,q}^d$ independent of a specific representation. If we consider a, possibly regularized, empirical risk minimization problem on the weights, the objective (the empirical risk) would never be a convex function of the weights (for depth $d \geq 2$), even if the regularizer is convex in w (which it always is for $p, q \geq 1$). But if we do not bound the width of the network, and instead rely on magnitude-control alone, we will see that the resulting hypothesis class, and indeed the complexity measure, may be convex (with respect to taking convex combinations of functions, *not* of weights).

Theorem 4 For any $d, p, q \geq 1$ such that $\frac{1}{q} \leq \frac{1}{d-1}(1 - \frac{1}{p})$, $\gamma_{p,q}^d(f)$ is a semi-norm in \mathcal{N}^d .

In particular, under the condition of the Theorem, $\gamma_{p,q}^d$ is convex, and hence its sublevel sets $\mathcal{N}_{p,q}^d$ are convex, and so $\mu_{p,q}^d$ is quasi-convex (but not convex).

Proof sketch To show convexity, consider two functions $f, g \in \mathcal{N}_{\gamma_{p,q} \leq \gamma}^d$ and $0 < \alpha < 1$, and let U and V be the weights realizing f and g respectively with $\gamma_{p,q}(U) \leq \gamma$ and $\gamma_{p,q}(V) \leq \gamma$. We will construct weights W realizing $\alpha f + (1 - \alpha)g$ with $\gamma_{p,q}(W) \leq \gamma$. This is done by first balancing U and V s.t. at each layer $\|U_i\|_{p,q} = \sqrt[d]{\gamma_{p,q}(U)}$ and $\|V_i\|_{p,q} = \sqrt[d]{\gamma_{p,q}(V)}$ and then placing U and V side by side, with no interaction between the units calculating f and g until the output layer. The output unit has weights αU_d coming in from the f -side and weights $(1 - \alpha)V_d$ coming in from the g -side. In Appendix C we show that under the condition in the theorem, $\gamma_{p,q}(W) \leq \gamma$. To complete the proof, we also show $\gamma_{p,q}^d$ is homogeneous and that this is sufficient for convexity. ■

4. Per-Unit and Path Regularization

In this Section we will focus on the special case of $q = \infty$, i.e. when we constrain the norm of the incoming weights of each unit separately.

Per-unit ℓ_1 -regularization was studied by [Bartlett \(1998\)](#); [Koltchinskii and Panchenko \(2002\)](#); [Bartlett and Mendelson \(2003\)](#) who showed generalization guarantees. A two-layer network of this form with RELU activation was also considered by [Bach \(2014\)](#), who studied its approximation ability and suggested heuristics for learning it. Per-unit ℓ_2 regularization in a two-layer network was considered by [Cho and Saul \(2009\)](#), who showed it is equivalent to using a specific kernel. We now introduce *Path regularization* and discuss its equivalence to Per-Unit regularization.

Path Regularization Consider a regularizer which looks at the sum over all paths from input nodes to the output node, of the product of the weights along the path:

$$\phi_p(w) = \left(\sum_{v_{\text{in}}[i] \xrightarrow{e_1} v_1 \xrightarrow{e_2} v_2 \dots \xrightarrow{e_k} v_{\text{out}}} \prod_{i=1}^k |w(e_i)|^p \right)^{1/p} \quad (7)$$

where $p \geq 1$ controls the norm used to aggregate the paths. We can motivate this regularizer as follows: if a node does not have any high-weight paths going out of it, we really don't care much about what comes into it, as it won't have much effect on the output. The path-regularizer thus looks at the aggregated influence of all the weights.

Referring to the induced regularizer $\phi_p^G(f) = \min_{f_G, w=f} \phi_p(w)$ (with the usual shorthands for layered graphs), we now observe that for layered graphs, path regularization and per-unit regularization are equivalent:

Theorem 5 For $p \geq 1$, any d and (finite or infinite) H , for any $f \in \mathcal{N}^{d,H}$: $\phi_p^{d,H}(f) = \gamma_{p,\infty}^{d,H}(f)$

It is important to emphasize that even for layered graphs, it is not the case that for all weights $\phi_p(w) = \gamma_{p,\infty}(w)$. E.g., a high-magnitude edge going into a unit with no non-zero outgoing edges will affect $\gamma_{p,\infty}(w)$ but not $\phi_p(w)$, as will having high-magnitude edges on different layers in different paths. In a sense path regularization is as more careful regularizer less fooled by imbalance. Nevertheless, in the proof of Theorem 5 in Appendix D.1, we show we can always balance the weights such that the two measures are equal.

The equivalence does not extend to non-layered graphs, since the lengths of different paths might be different. Again, we can think of path regularizer as more refined regularizer taking into account the local structure. However, if we consider all DAGs of depth at most d (i.e. with paths of length at most d), the notions are again equivalent (see proof in Appendix D.2):

Theorem 6 For any $p \geq 1$ and any d : $\gamma_{p,\infty}^d(f) = \min_{G \in \text{DAG}(d)} \phi_p^G(f)$.

In particular, for any graph G of depth d , we have that $\phi_p^G(f) \geq \gamma_{p,\infty}^d(f)$. Combining this observation with Corollary 2 allows us to immediately obtain a generalization bound for path regularization on any, even non-layered, graph:

Corollary 7 For any graph G of depth d and any set $S = \{x_1, \dots, x_m\} \subseteq \mathbb{R}^D$:

$$\mathcal{R}_m(\mathcal{N}_{\phi_1 \leq \phi}^G) \leq \sqrt{\frac{4^{d-1} \phi^2 \cdot 4 \log(2D) \sup \|x_i\|_\infty^2}{m}}$$

Note that in order to apply Corollary 2 and obtain a width-independent bound, we had to limit ourselves to $p = 1$. We further explore this issue next.

Capacity As was previously noted, size-independent generalization bounds for bounded depth networks with bounded per-unit ℓ_1 norm have long been known (and make for a popular homework problem). These correspond to a specialization of Corollary 2 for the case $p = 1, q = \infty$. Furthermore, the kernel view of Cho and Saul (2009) allows obtaining size-independent generalization bound for *two-layer* networks with bounded per-unit ℓ_2 norm (i.e. a single infinite hidden layer of all possible unit-norm units, and a bounded ℓ_2 -norm output unit). However, the lower bound of Theorem 3 establishes that for any $p > 1$, once we go beyond two layers, we cannot ensure generalization without also controlling the size (or width) of the network.

Convexity An immediate consequence of Theorem 4 is that per-unit regularization, if we do not constrain the network width, is convex for any $p \geq 1$. In fact, $\gamma_{p,\infty}^d$ is a (semi)norm. However, as discussed above, for depth $d > 2$ this is meaningful only for $p = 1$, as $\gamma_{p,\infty}^d$ collapses for $p > 1$.

Hardness Since the classes $\mathcal{N}_{1,\infty}^d$ are convex, we might hope that this might make learning computationally easier. Indeed, one can consider functional-gradient or boosting-type strategies for learning a predictor in the class (Lee et al., 1996). However, as Bach (2014) points out, this is not so easy as it requires finding the best fit for a target with a RELU unit, which is not easy. Indeed,

applying results on hardness of learning intersections of halfspaces, which can be represented with small per-unit norm using two-layer networks, we can conclude that, subject to certain complexity assumptions, it is not possible to efficiently PAC learn $\mathcal{N}_{1,\infty}^d$, even for depth $d = 2$ when $\gamma_{1,\infty}$ increases superlinearly:

Corollary 8 *Subject to the the strong random CSP assumptions in Daniely et al. (2014), it is not possible to efficiently PAC learn (even improperly) functions $\{\pm 1\}^D \rightarrow \{\pm 1\}$ realizable with unit margin by $\mathcal{N}_{1,\infty}^2$ when $\gamma_{1,\infty} = \omega(D)$ (e.g. when $\gamma_{1,\infty} = D \log D$). Moreover, subject to intractability of $\tilde{Q}(D^{1.5})$ -unique shortest vector problem, for any $\epsilon > 0$, it is not possible to efficiently PAC learn (even improperly) functions $\{\pm 1\}^D \rightarrow \{\pm 1\}$ realizable with unit margin by $\mathcal{N}_{1,\infty}^2$ when $\gamma_{1,\infty} = D^{1+\epsilon}$.*

This is a corollary of Theorem 22 in the Appendix E. Either versions of corollary 8 precludes the possibility of learning in time polynomial in $\gamma_{1,\infty}$, though it still might be possible to learn in $\text{poly}(D)$ time when $\gamma_{1,\infty}$ is sublinear.

Sharing We conclude this Section with an observation on the type of networks obtained by per-unit, or equivalently path, regularization.

Theorem 9 *For any $p \geq 1$ and $d > 1$ and any $f \in \mathcal{N}^d$, there exists a layered graph $G(V, E)$ of depth d , such that $f \in \mathcal{N}^G$ and $\gamma_{p,\infty}^G(f) = \phi_p^G(f) = \gamma_{p,\infty}^d(f)$, and the out-degree of every internal (non-input) node in G is one. That is, the subgraph of G induced by the non-input vertices is a tree directed toward the output vertex.*

The proof is given in Appendix D.3. What the Theorem tells us is that we can realize every function as a tree with optimal per-unit norm. If we think of learning with an infinite fully-connected layered network, we can always restrict ourselves to models in which the non-zero-weight edges form a tree. This means that when using per-unit regularization we have no incentive to “share” lower-level units—each unit will only have a single outgoing edge and will only be used by a single down-stream unit. This seems to defy much of the intuition and power of using deep networks, where we expect lower layers to represent generic feature useful in many higher-level features. In effect, we are not encouraging any transfer between learning different aspects of the function (or between different tasks or classes, if we do have multiple output units). Per-unit regularization therefore misses out on much of the inductive bias that we might like to impose when using deep learning (namely, promoting sharing).

5. Overall Regularization

In this Section, we will focus on “overall” ℓ_p regularization, corresponding to the choice $q = p$, i.e. when we bound the overall (vectorized) norm of all weights in the system:

$$\mu_{p,p}(w) = \left(\sum_{e \in E} |w(e)|^p \right)^{1/p}.$$

Capacity For $p \leq 2$, Corollary 2 provides a generalization guarantee that is independence of the width—we can conclude that if we use weight decay (overall ℓ_2 regularization), or any tighter ℓ_p regularization, there is no need to limit ourselves to networks of finite size (as long as the corresponding dual-norm of the inputs are bounded). However, in Section 3.2 we saw that with $d \geq 3$

layers, the regularizer degenerates and leads to infinite capacity classes if $p > 2$. In any case, even if we bound the overall ℓ_1 -norm, the complexity increases exponentially with the depth.

Convexity The conditions of Theorem 4 for convexity of $\mathcal{N}_{2,2}^d$ are ensured when $p \geq d$. For depth $d = 1$, i.e. a single unit, this just confirms that ℓ_p -regularized linear prediction is convex for $p \geq 1$. For depth $d = 2$, we get convexity with ℓ_2 regularization, but not ℓ_1 . For depth $d > 2$ we would need $p > d \geq 3$, however for such values of p we know from Theorem 3 that $\mathcal{N}_{p,p}^d$ degenerates to an infinite capacity class if we do not control the width (if we do control the width, we do not get convexity). This leaves us with $\mathcal{N}_{2,2}^2$ as the interesting convex class. Below we show an explicit convex characterization of $\mathcal{N}_{2,2}^2$ by showing it is equivalent to so-called ‘‘convex neural nets’’.

Convex Neural Nets (Bengio et al., 2005) over inputs in \mathbb{R}^D are two-layer networks with a fixed infinite hidden layer consisting of all units with weights $w \in \mathcal{G}$ for some base class $\mathcal{G} \in \mathbb{R}^D$, and a second ℓ_1 -regularized layer. Since over finite data the weights in the second layer can always be taken to have finite support (i.e. be non-zero for only a finite number of first-layer units), and we can approach any function with countable support, we can instead think of a network in \mathcal{N}^2 where the bottom layer is constraint to \mathcal{G} and the top layer is ℓ_1 regularized. Focusing on $\mathcal{G} = \{w \mid \|w\|_p \leq 1\}$, this corresponds to imposing an ℓ_p constraint on the bottom layer, and ℓ_1 regularization on the top layer and yields the following complexity measure over \mathcal{N}^2 :

$$\nu_p(f) = \inf_{f_{\text{layer}(d)}, W=f, \text{s.t. } \forall_j \|W_1[j, :]\|_p \leq 1} \|W_2\|_1. \quad (8)$$

This is similar to per-unit regularization, except we impose different norms at different layers (if $p \neq 1$). We can see that $\mathcal{N}_{\nu_p \leq \nu}^2 = \nu \cdot \overline{\text{conv}}(\sigma(\mathcal{G}))$, and is thus convex for any p . Focusing on RELU activation we have the equivalence:

Theorem 10 $\mu_{2,2}^2(f) = 2\nu_2(f)$.

That is, overall ℓ_2 regularization with two layers is equivalent to a convex neural net with ℓ_2 -constrained units on the bottom layer and ℓ_1 (not ℓ_2 !) regularization on the output.

Proof We can calculate:

$$\begin{aligned} \min_{f_W=f} \mu_{2,2}^2(W) &= \min_{f_W=f} \sum_{j=1}^H \left(\sum_{i=1}^D |W_1[j, i]|^2 + |W_2[j]|^2 \right) \\ &= \min_{f_W=f} \sum_{j=1}^H 2 \sqrt{\sum_{i=1}^D |W_1[j, i]|^2 \cdot |W_2[j]|^2} \end{aligned} \quad (9)$$

$$= 2 \min_{f_W=f} \sum_{j=1}^H |W_2[j]| \quad \text{s.t.} \quad \sqrt{\sum_{i=1}^D |W_1[j, i]|^2} \leq 1. \quad (10)$$

Here (9) is the arithmetic-geometric mean inequality for which we can achieve equality by balancing the weights (as in Claim 1) and (10) again follows from the homogeneity of the RELU which allows us to rebalance the weights. \blacksquare

Hardness As with $\mathcal{N}_{1,\infty}^d$, we might hope that the convexity of $\mathcal{N}_{2,2}^2$ might make it computationally easy to learn. However, by the same reduction from learning intersection of halfspaces (Theorem 22 in Appendix E) we can again conclude that we cannot learn in time polynomial in $\mu_{2,2}^2$:

Corollary 11 *Subject to the the strong random CSP assumptions in Daniely et al. (2014), it is not possible to efficiently PAC learn (even improperly) functions $\{\pm 1\}^D \rightarrow \{\pm 1\}$ realizable with unit margin by $\mathcal{N}_{p,p}^2$ when $\mu_{p,p}^2 = \omega(D^{\frac{1}{p}})$. (e.g. when $\gamma_{1,\infty} = D \log D$). Moreover, subject to intractability of $\tilde{Q}(D^{1.5})$ -unique shortest vector problem, for any $\epsilon > 0$, it is not possible to efficiently PAC learn (even improperly) functions $\{\pm 1\}^D \rightarrow \{\pm 1\}$ realizable with unit margin by $\mathcal{N}_{1,\infty}^2$ when $\gamma_{1,\infty} = D^{\frac{1}{p}+\epsilon}$.*

6. Depth Independent Regularization

Up until now we discussed relying on magnitude-based regularization instead of directly controlling network size, thus allowing unbounded and even infinite width. But we still relied on a finite bound on the depth in all our derivations. Can the explicit dependence on the depth be avoided, and replaced with only a measure of scale of the weights?

We already know we cannot rely only on a bound on the group-norm $\mu_{p,q}$ when the depth is unbounded, as we know from Theorem 3 that in terms of $\mu_{p,q}$ the sample complexity necessarily increases exponentially with the depth: if we allow arbitrarily deep graphs we can shrink $\mu_{p,q}$ toward zero without changing the scale of the computed function. However, controlling the γ -measure, or equivalently the path-regularizer ϕ , in arbitrarily-deep graphs is sensible, and we can define:

$$\gamma_{p,q} = \inf_{d \geq 1} \gamma_{p,q}^d(f) = \lim_{d \rightarrow \infty} \gamma_{p,q}^d(f) \quad \text{or:} \quad \phi_p = \inf_G \phi_p^G(f) \quad (11)$$

where the minimization is over *any* DAG. From Theorem 6 we can conclude that $\phi_p(f) = \gamma_{p,\infty}(f)$. In any case, $\gamma_{p,q}(f)$ is a sensible complexity measure, that does not collapse despite the unbounded depth. Can we obtain generalization guarantees for the class $\mathcal{N}_{\gamma_{p,q} \leq \gamma}$?

Unfortunately, even when $1/p + 1/q \geq 1$ and we can obtain width-independent bounds, the bound in Corollary 2 still has a dependence on 4^d , even if $\gamma_{p,q}$ is bounded. Can such a dependence be avoided?

For *anti-symmetric* Lipschitz-continuous activation functions (i.e. such that $\sigma(-z) = -\sigma(z)$), such as the ramp, and for per-unit ℓ_p -regularization $\mu_{1,\infty}^d$ we can avoid the factor of 4^d

Theorem 12 *For any anti-symmetric 1-Lipschitz function σ and any set $S = \{x_1, \dots, x_m\} \subseteq \mathbb{R}^D$:*

$$\mathcal{R}_m(\mathcal{N}_{\mu_{1,\infty} \leq \mu}^{d,\sigma}) \leq \sqrt{\frac{4\mu^{2d} \log(2D) \sup \|x_i\|_\infty^2}{m}}$$

The proof is again based on an inductive argument similar to Theorem 1 and you can find it in appendix A.4.

However, the ramp is not homogeneous and so the equivalent between μ , γ and ϕ breaks down. Can we obtain such a bound also for the RELU? At the very least, what we can say is that an inductive argument such that used in the proofs of Theorems 1 and 12 cannot be used to avoid an

exponential dependence on the depth. To see this, consider $\gamma_{1,\infty} \leq 1$ (this choice is arbitrary if we are considering the Rademacher complexity), for which we have

$$\mathcal{N}_{\gamma_{1,\infty} < 1}^{d+1} = \left[\overline{\text{conv}}(\mathcal{N}_{\gamma_{1,\infty} < 1}^d) \right]_+, \quad (12)$$

where $\overline{\text{conv}}(\cdot)$ is the symmetric convex hull, and $[\cdot]_+ = \max(z, 0)$ is applied to each function in the class. In order to apply the inductive argument without increasing the complexity exponentially with the depth, we would need the operation $[\overline{\text{conv}}(\mathcal{H})]_+$ to preserve the Rademacher complexity, at least for non-negative convex cones \mathcal{H} . However we show a simple example of a non-negative convex cone \mathcal{H} for which $\mathcal{R}_m([\overline{\text{conv}}(\mathcal{H})]_+) > \mathcal{R}_m(\mathcal{H})$.

We will specify \mathcal{H} as a set of vectors in \mathbb{R}^m , corresponding to the evaluation of $h(x_i)$ of different functions in the class on the m points x_i in the sample. In our construction, we will have only $m = 3$ points. Consider $\mathcal{H} = \text{conv}(\{(1, 0, 1), (0, 1, 1)\})$, in which case $\mathcal{H}' \triangleq [\overline{\text{conv}}(\mathcal{H})]_+ = \text{conv}(\{(1, 0, 1), (0, 1, 1), (0.5, 0, 0)\})$. It is not hard to verify that $\mathcal{R}_m(\mathcal{H}') = \frac{13}{16} > \frac{12}{16} = \mathcal{R}_m(\mathcal{H})$.

7. Summary and Open Issues

We presented a general framework for norm-based capacity control for feed-forward networks, and analyzed when the norm-based control is sufficient and to what extent capacity still depends on other parameters. In particular, we showed that in depth $d > 2$ networks, per-unit control with $p > 1$ and overall regularization with $p > 2$ is not sufficient for capacity control without also controlling the network size. This is in contrast with linear models, where with any $p < \infty$ we have only a weak dependence on dimensionality, and two-layer networks where per-unit $p = 2$ is also sufficient for capacity control. We also obtained generalization guarantees for perhaps the most natural form of regularization, namely ℓ_2 regularization, and showed that even with such control we still necessarily have an exponential dependence on the depth.

Although the additive μ -measure and multiplication γ -measure are equivalent at the optimum, they behave rather differently in terms of optimization dynamics (based on anecdotal empirical experience) and understanding the relationship between them, as well as the novel path-based regularizer can be helpful in practical regularization of neural networks.

Although we obtained a tight characterization of when size-independent capacity control is possible, the precise polynomial dependence of margin-based classification (and other tasks) on the norm in might not be tight and can likely be improved, though this would require going beyond bounding the Rademacher complexity of the real-valued class. In particular, Theorem 1 gives the same bound for per-unit ℓ_1 regularization and overall ℓ_1 regularization, although we would expect the later to have lower capacity.

Beyond the open issue regarding depth-independent γ -based capacity control, another interesting open question is understanding the expressive power of $\mathcal{N}_{\gamma_{p,q} \leq \gamma}^d$, particularly as a function of the depth d . Clearly going from depth $d = 1$ to depth $d = 2$ provides additional expressive power, but it is not clear how much additional depth helps. The class \mathcal{N}^2 already includes all binary functions over $\{\pm 1\}^D$ and is dense among continuous real-valued functions. But can the γ -measure be reduced by increasing the depth? Viewed differently: $\gamma_{p,q}^d(f)$ is monotonically non-increasing in d , but are there functions for it continues decreasing? Although it seems obvious there are functions that require high depth for efficient representation, these questions are related to decade-old problems in circuit complexity and might not be easy to resolve.

Acknowledgments

This research was partially supported by NSF grant IIS-1302662 and an Intel ICRI-CI award. We thank the COLT anonymous reviewers for pointing out an error in the statement of Lemma 15 and suggesting other corrections.

References

- Martin Anthony and Peter L. Bartlett. *Neural network learning: Theoretical foundations*. Cambridge University Press, 2009.
- Francis Bach. Breaking the curse of dimensionality with convex neural networks. *Technical report, HAL-01098505*, 2014.
- Maria-Florina Balcan and Christopher Berlind. A new perspective on learning linear separators with large lq1p margins. *Proceedings of the Seventeenth International Conference on Artificial Intelligence and Statistics*, pages 68–76, 2014.
- Peter L. Bartlett. The sample complexity of pattern classification with neural networks: the size of the weights is more important than the size of the network. *IEEE transactions on information theory*, 44(2):525–536, 1998.
- Peter L. Bartlett and Shahar Mendelson. Rademacher and gaussian complexities: Risk bounds and structural results. *The Journal of Machine Learning Research*, pages 463–482, 2003.
- Yoshua Bengio, Nicolas L. Roux, Pascal Vincent, Olivier Delalleau, and Patrice Marcotte. Convex neural networks. *Advances in neural information processing systems*, pages 123–130, 2005.
- Xavier Glorot Antoine Bordes and Yoshua Bengio. Deep sparse rectifier networks. *AISTATS*, 2011.
- Youngmin Cho and Lawrence K. Saul. Kernel methods for deep learning. *Advances in neural information processing systems*, pages 342–350, 2009.
- Amit Daniely, Nati Linial, and Shai Shalev-Shwartz. From average case complexity to improper learning complexity. *STOC*, 2014.
- Uffe Haagerup. The best constants in the khintchine inequality. *Studia Mathematica*, 70(3):231–283, 1981.
- Sham M Kakade, Karthik Sridharan, and AmbujTewari. On the complexity of linear prediction: Risk bounds, margin bounds, and regularization. *Advances in neural information processing systems*, pages 793–800, 2009.
- Adam R Klivans and Alexander A Sherstov. Cryptographic hardness for learning intersections of halfspaces. *FOCS*, pages 553–562, 2006.
- Vladimir Koltchinskii and Dmitry Panchenko. Empirical margin distributions and bounding the generalization error of combined classifiers. *Annals of Statistics*, pages 1–50, 2002.

Wee Sun Lee, Peter L Bartlett, and Robert C Williamson. Efficient agnostic learning of neural networks with bounded fan-in. *Information Theory, IEEE Transactions on*, 42(6):2118–2132, 1996.

Roi Livni, Shai Shalev-Shwartz, and Ohad Shamir. On the computational efficiency of training neural networks. *Advances in Neural Information Processing Systems*, pages 855–863, 2014.

Vinod Nair and Geoffrey E. Hinton. Rectified linear units improve restricted boltzmann machines. *ICML*, 2010.

Shai Shalev-Shwartz and Shai Ben-David. *Understanding Machine Learning: From Theory to Algorithms*. Cambridge University Press, 2014.

M.D. Zeiler, M. Ranzato, R. Monga, M. Mao, K. Yang, Q.V. Le, P. Nguyen, A. Senior, V. Vanhoucke, J. Dean, and G.E. Hinton. On rectified linear units for speech processing. *ICASSP*, 2013.

Appendix A. Rademacher Complexities

The sample based Rademacher complexity of a class \mathcal{F} of function mapping from \mathcal{X} to \mathbb{R} with respect to a set $S = \{x_1, \dots, x_m\}$ is defined as:

$$\mathcal{R}_m(\mathcal{F}) = \mathbb{E}_{\xi \in \{\pm 1\}^m} \left[\frac{1}{m} \sup_{f \in \mathcal{F}} \left| \sum_{i=1}^m \xi_i f(x_i) \right| \right]$$

In this section, we prove an upper bound for the Rademacher complexity of the class $\mathcal{N}_{\gamma_{p,q} \leq \gamma}^{d,H,\sigma_{\text{RELU}}}$, i.e., the class of functions that can be represented as depth d , width H network with rectified linear activations, and the layer-wise group norm complexity $\gamma_{p,q}$ bounded by γ . As mentioned in the main text, our proof is an induction with respect to the depth d . We start with $d = 1$ layer neural networks, which is essentially the class of linear separators.

A.1. ℓ_p -regularized Linear Predictors

For completeness, we prove the upper bounds on the Rademacher complexity of class of linear separators with bounded ℓ_p norm. The upper bounds presented here are particularly similar to generalization bounds in [Kakade et al. \(2009\)](#) and [Balcan and Berlind \(2014\)](#). We first mention two already established lemmas that we use in the proofs.

Theorem 13 (Khintchine-Kahane Inequality) For any $0 < p < \infty$ and $S = \{z_1, \dots, z_m\}$, if the random variable ξ is uniform over $\{\pm 1\}^m$, then

$$\left(\mathbb{E}_{\xi} \left[\left| \sum_{i=1}^m \xi_i z_i \right|^p \right] \right)^{\frac{1}{p}} \leq C_p \left(\sum_{i=1}^m |z_i|^2 \right)^{\frac{1}{2}}$$

where C_p is a constant depending only on p .

The sharp value of the constant C_p was found by [Haagerup \(1981\)](#) but for our analysis, it is enough to note that if $p \geq 1$ we have $C_p \leq \sqrt{p}$.

Lemma 14 (*Massart Lemma*) *Let A be a finite set of m dimensional vectors. Then*

$$\mathbb{E}_\xi \left[\max_{a \in A} \frac{1}{m} \sum_{i=1}^m \xi_i a_i \right] \leq \max_{a \in A} \|a\|_2 \frac{\sqrt{2 \log |A|}}{m},$$

where $|A|$ is the cardinality of A .

We are now ready to show upper bounds on Rademacher complexity of linear separators with bounded ℓ_p norm.

Lemma 15 (*Rademacher complexity of linear separators with bounded ℓ_p norm*) *For any $d, q \geq 1$, For any $1 \leq p \leq 2$,*

$$\mathcal{R}_m(\mathcal{N}_{\gamma_{p,q} \leq \gamma}^1) \leq \sqrt{\frac{\gamma^2 \min\{p^*, 4 \log(2D)\} \max_i \|x_i\|_{p^*}^2}{m}}$$

and for any $2 < p < \infty$

$$\mathcal{R}_m(\mathcal{N}_{\gamma_{p,q} \leq \gamma}^1) \leq \frac{\sqrt{2}\gamma \|X\|_{2,p^*}}{m} \leq \frac{\sqrt{2}\gamma \max_i \|x_i\|_{p^*}}{m^{\frac{1}{p}}}$$

where p^* is such that $\frac{1}{p^*} + \frac{1}{p} = 1$.

Proof First, note that \mathcal{N}^1 is the class of linear functions and hence for any function $f_w \in \mathcal{N}^1$, we have that $\gamma_{p,q}(w) = \|w\|_p$. Therefore, we can write the Rademacher complexity for a set $S = \{x_1, \dots, x_m\}$ as:

$$\begin{aligned} \mathcal{R}_m(\mathcal{N}_{\gamma_{p,q} \leq \gamma}^1) &= \mathbb{E}_{\xi \in \{\pm 1\}^m} \left[\frac{1}{m} \sup_{\|w\|_p \leq \gamma} \left| \sum_{i=1}^m \xi_i w^\top x_i \right| \right] \\ &= \mathbb{E}_{\xi \in \{\pm 1\}^m} \left[\frac{1}{m} \sup_{\|w\|_p \leq \gamma} \left| w^\top \sum_{i=1}^m \xi_i x_i \right| \right] \\ &= \gamma \mathbb{E}_{\xi \in \{\pm 1\}^m} \left[\frac{1}{m} \left\| \sum_{i=1}^m \xi_i x_i \right\|_{p^*} \right] \end{aligned}$$

For $1 \leq p \leq \min \left\{ 2, \frac{2 \log(2D)}{2 \log(2D) - 1} \right\}$ (and therefore $2 \log(2D) \leq p^*$), we have

$$\begin{aligned}
 \mathcal{R}_m(\mathcal{N}_{\gamma p, q \leq \gamma}^1) &= \gamma \mathbb{E}_{\xi \in \{\pm 1\}^m} \left[\frac{1}{m} \left\| \sum_{i=1}^m \xi_i x_i \right\|_{p^*} \right] \\
 &\leq D^{\frac{1}{p^*}} \gamma \mathbb{E}_{\xi \in \{\pm 1\}^m} \left[\frac{1}{m} \left\| \sum_{i=1}^m \xi_i x_i \right\|_{\infty} \right] \\
 &\leq D^{\frac{1}{2 \log(2D)}} \gamma \mathbb{E}_{\xi \in \{\pm 1\}^m} \left[\frac{1}{m} \left\| \sum_{i=1}^m \xi_i x_i \right\|_{\infty} \right] \\
 &\leq \sqrt{2} \gamma \mathbb{E}_{\xi \in \{\pm 1\}^m} \left[\frac{1}{m} \left\| \sum_{i=1}^m \xi_i x_i \right\|_{\infty} \right]
 \end{aligned}$$

We now use the Massart Lemma viewing each feature $(x_i[j])_{i=1}^m$ for $j = 1, \dots, D$ as a member of a finite hypothesis class and obtain

$$\begin{aligned}
 \mathcal{R}_m(\mathcal{N}_{\gamma p, q \leq \gamma}^1) &\leq \sqrt{2} \gamma \mathbb{E}_{\xi \in \{\pm 1\}^m} \left[\frac{1}{m} \left\| \sum_{i=1}^m \xi_i x_i \right\|_{\infty} \right] \\
 &\leq 2\gamma \frac{\sqrt{\log(2D)}}{m} \max_{j=1, \dots, D} \|(x_i[j])_{i=1}^m\|_2 \\
 &\leq 2\gamma \sqrt{\frac{\log(2D)}{m}} \max_{i=1, \dots, m} \|x_i\|_{\infty} \\
 &\leq 2\gamma \sqrt{\frac{\log(2D)}{m}} \max_{i=1, \dots, m} \|x_i\|_{p^*}
 \end{aligned}$$

If $\min \left\{ 2, \frac{2 \log(2D)}{2 \log(2D) - 1} \right\} < p < \infty$, by Khintchine-Kahane inequality we have

$$\begin{aligned}
 \mathcal{R}_m(\mathcal{N}_{\gamma p, q \leq \gamma}^1) &= \gamma \mathbb{E}_{\xi \in \{\pm 1\}^m} \left[\frac{1}{m} \left\| \sum_{i=1}^m \xi_i x_i \right\|_{p^*} \right] \\
 &\leq \gamma \frac{1}{m} \left(\sum_{j=1}^D \mathbb{E}_{\xi \in \{\pm 1\}^m} \left[\left\| \sum_{i=1}^m \xi_i x_i[j] \right\|^{p^*} \right] \right)^{1/p^*} \\
 &\leq \gamma \frac{\sqrt{p^*}}{m} \left(\sum_{j=1}^D \|(x_i[j])_{i=1}^m\|_2^{p^*} \right)^{1/p^*} = \gamma \frac{\sqrt{p^*}}{m} \|X\|_{2, p^*}
 \end{aligned}$$

If $p^* \geq 2$, by Minskowski inequality we have that $\|X\|_{2, p^*} \leq m^{1/2} \max_i \|x_i\|_{p^*}$. Otherwise, by subadditivity of the function $f(z) = z^{\frac{p^*}{2}}$, we get $\|X\|_{2, p^*} \leq m^{1/p^*} \max_i \|x_i\|_{p^*}$. ■

A.2. Theorem 1

We define the hypothesis class $\mathcal{N}^{d,H,H}$ to be the class of functions from \mathcal{X} to \mathbb{R}^H computed by a layered network of depth d , layer size H and H outputs.

For the proof of theorem 1, we need the following two technical lemmas. The first is the well-known contraction lemma:

Lemma 16 (Contraction Lemma) *Let function $\phi : \mathbb{R} \rightarrow \mathbb{R}$ be Lipschitz with constant \mathcal{L}_ϕ such that ϕ satisfies $\phi(0) = 0$. Then for any class \mathcal{F} of functions mapping from \mathcal{X} to \mathbb{R} and any set $S = \{x_1, \dots, x_m\}$:*

$$\mathbb{E}_{\xi \in \{\pm 1\}^m} \left[\frac{1}{m} \sup_{f \in \mathcal{F}} \left| \sum_{i=1}^m \xi_i \phi(f(x_i)) \right| \right] \leq 2\mathcal{L}_\phi \mathbb{E}_{\xi \in \{\pm 1\}^m} \left[\frac{1}{m} \sup_{f \in \mathcal{F}} \left| \sum_{i=1}^m \xi_i f(x_i) \right| \right]$$

Next, the following lemma reduces the maximization over a matrix $W \in \mathbb{R}^{H \times H}$ that appears in the computation of Rademacher complexity to H independent maximizations over a vector $w \in \mathbb{R}^H$ (the proof is deferred to Subsection A.3):

Lemma 17 *For any $p, q \geq 1$, $d \geq 2$, $\xi \in \{\pm 1\}^m$ and $f \in \mathcal{N}^{d,H,H}$ we have*

$$\sup_W \frac{1}{\|W\|_{p,q}} \left\| \sum_{i=1}^m \xi_i [W[f(x_i)]_+] \right\|_{p^*} = H^{\lceil \frac{1}{p^*} - \frac{1}{q} \rceil_+} \sup_w \frac{1}{\|w\|_p} \left| \sum_{i=1}^m \xi_i [w^\top [f(x_i)]_+] \right|$$

where p^* is such that $\frac{1}{p^*} + \frac{1}{p} = 1$.

Theorem 1 *For any $d, p, q \geq 1$ and any set $S = \{x_1, \dots, x_m\} \subseteq \mathbb{R}^D$:*

$$\mathcal{R}_m(\mathcal{N}_{\gamma_{p,q} \leq \gamma}^{d,H,\sigma_{\text{RELU}}}) \leq \sqrt{\frac{\gamma^2 \left(2H^{\lceil \frac{1}{p^*} - \frac{1}{q} \rceil_+} \right)^{2(d-1)} \min\{p^*, 2 \log(2D)\} \sup \|x_i\|_{p^*}^2}{m}}$$

and so:

$$\mathcal{R}_m(\mathcal{N}_{\mu_{p,q} \leq \mu}^{d,H,\sigma_{\text{RELU}}}) \leq \sqrt{\frac{\mu^{2d} \left(2H^{\lceil \frac{1}{p^*} - \frac{1}{q} \rceil_+} / \sqrt[d]{d} \right)^{2(d-1)} \min\{p^*, 2 \log(2D)\} \sup \|x_i\|_{p^*}^2}{m}}$$

where p^* is such that $\frac{1}{p^*} + \frac{1}{p} = 1$.

Proof By the definition of Rademacher complexity if ξ is uniform over $\{\pm 1\}^m$, we have:

$$\begin{aligned}
 \mathcal{R}_m(\mathcal{N}_{\gamma_{p,q} \leq \gamma}^{d,H}) &= \mathbb{E}_\xi \left[\frac{1}{m} \sup_{f \in \mathcal{N}_{\gamma_{p,q} \leq \gamma}^{d,H}} \left| \sum_{i=1}^m \xi_i f(x_i) \right| \right] \\
 &= \mathbb{E}_\xi \left[\frac{1}{m} \sup_{f \in \mathcal{N}^{d,H}} \frac{\gamma}{\gamma_{p,q}(f)} \left| \sum_{i=1}^m \xi_i f(x_i) \right| \right] \\
 &= \mathbb{E}_\xi \left[\frac{1}{m} \sup_{g \in \mathcal{N}^{d-1,H,H}} \sup_w \frac{\gamma}{\gamma_{p,q}(g) \|w\|_p} \left| \sum_{i=1}^m \xi_i w^\top [g(x_i)]_+ \right| \right] \\
 &= \mathbb{E}_\xi \left[\frac{1}{m} \sup_{g \in \mathcal{N}^{d-1,H,H}} \frac{\gamma}{\gamma_{p,q}(g)} \left\| \sum_{i=1}^m \xi_i [g(x_i)]_+ \right\|_{p^*} \right] \\
 &= \mathbb{E}_\xi \left[\frac{1}{m} \sup_{h \in \mathcal{N}^{d-2,H,H}} \frac{\gamma}{\gamma_{p,q}(h)} \sup_W \frac{1}{\|W\|_{p,q}} \left\| \sum_{i=1}^m \xi_i [W[h(x_i)]_+]_+ \right\|_{p^*} \right] \\
 &= H^{[\frac{1}{p^*} - \frac{1}{q}]_+} \mathbb{E}_\xi \left[\frac{1}{m} \sup_{h \in \mathcal{N}^{d-2,H,H}} \frac{\gamma}{\gamma_{p,q}(h)} \sup_w \frac{1}{\|w\|_p} \left| \sum_{i=1}^m \xi_i [w^\top [h(x_i)]_+]_+ \right| \right] \quad (13)
 \end{aligned}$$

$$\begin{aligned}
 &= H^{[\frac{1}{p^*} - \frac{1}{q}]_+} \mathbb{E}_\xi \left[\frac{1}{m} \sup_{g \in \mathcal{N}_{\gamma_{p,q} \leq \gamma}^{d-1,H}} \left| \sum_{i=1}^m \xi_i [g(x_i)]_+ \right| \right] \\
 &\leq 2H^{[\frac{1}{p^*} - \frac{1}{q}]_+} \mathbb{E}_\xi \left[\frac{1}{m} \sup_{g \in \mathcal{N}_{\gamma_{p,q} \leq \gamma}^{d-1,H}} \left| \sum_{i=1}^m \xi_i g(x_i) \right| \right] \quad (14) \\
 &= 2H^{[\frac{1}{p^*} - \frac{1}{q}]_+} \mathcal{R}_m(\mathcal{N}_{\gamma_{p,q} \leq \gamma}^{d-1,H})
 \end{aligned}$$

where the equality (13) is obtained by lemma 17 and inequality (14) is by Contraction Lemma. This will give us the bound on Rademacher complexity of $\mathcal{N}_{\gamma_{p,q} \leq \gamma}^{d,H}$ based on the Rademacher complexity of $\mathcal{N}_{\gamma_{p,q} \leq \gamma}^{d-1,H}$. Applying the same argument on all layers and using lemma 15 to bound the complexity of the first layer completes the proof. \blacksquare

A.3. Proof of Lemma 17

Proof It is immediate that the right hand side of the equality in the statement is always less than or equal to the left hand side because given any vector w in the right hand side, by setting each row of matrix W in the left hand side we get the equality. Therefore, it is enough to prove that the left hand side is less than or equal to the right hand side. For the convenience of notations, let $g(w) \triangleq \left| \sum_{i=1}^m \xi_i w^\top [f(x_i)]_+ \right|$. Define \tilde{w} to be:

$$\tilde{w} \triangleq \arg \max_w \frac{g(w)}{\|w\|_p}$$

If $q \leq p^*$, then the right hand side of equality in the lemma statement will reduce to $g(\tilde{w})/\|\tilde{w}\|_p$ and therefore we need to show that for any matrix V ,

$$\frac{g(\tilde{w})}{\|\tilde{w}\|_p} \geq \frac{\|g(V)\|_{p^*}}{\|V\|_{p,q}}.$$

Since $q \leq p^*$, we have $\|V\|_{p,p^*} \leq \|V\|_{p,q}$ and hence it is enough to prove the following inequality:

$$\frac{g(\tilde{w})}{\|\tilde{w}\|_p} \geq \frac{\|g(V)\|_{p^*}}{\|V\|_{p,p^*}}.$$

On the other hand, if $q > p^*$, then we need to prove the following inequality holds:

$$H^{\frac{1}{p^*} - \frac{1}{q}} \frac{g(\tilde{w})}{\|\tilde{w}\|_p} \geq \frac{\|g(V)\|_{p^*}}{\|V\|_{p,q}}$$

Since $q > p^*$, we have that $\|V\|_{p,p^*} \leq H^{\frac{1}{p^*} - \frac{1}{q}} \|V\|_{p,q}$. Therefore, it is again enough to show that:

$$\frac{g(\tilde{w})}{\|\tilde{w}\|_p} \geq \frac{\|g(V)\|_{p^*}}{\|V\|_{p,p^*}}.$$

We can rewrite the above inequality in the following form:

$$\sum_{i=1}^H \left(\frac{g(\tilde{w}) \|V_i\|_p}{\|\tilde{w}\|_p} \right)^{p^*} \geq \sum_{i=1}^H g(V_i)^{p^*}$$

By the definition of \tilde{w} , we know that the above inequality holds for each term in the sum and hence the inequality is true. \blacksquare

A.4. Theorem 12

The proof is similar to the proof of theorem 1 but here bounding $\mu_{1,\infty}$ by μ means the ℓ_1 norm of input weights to each neuron is bounded by μ . We use a different version of Contraction Lemma in the proof that is without the absolute value:

Lemma 18 (*Contraction Lemma (without the absolute value)*) *Let function $\phi : \mathbb{R} \rightarrow \mathbb{R}$ be Lipschitz with constant \mathcal{L}_ϕ . Then for any class \mathcal{F} of functions mapping from \mathcal{X} to \mathbb{R} and any set $S = \{x_1, \dots, x_m\}$:*

$$\mathbb{E}_{\xi \in \{\pm 1\}^m} \left[\frac{1}{m} \sup_{f \in \mathcal{F}} \sum_{i=1}^m \xi_i \phi(f(x_i)) \right] \leq \mathcal{L}_\phi \mathbb{E}_{\xi \in \{\pm 1\}^m} \left[\frac{1}{m} \sup_{f \in \mathcal{F}} \sum_{i=1}^m \xi_i f(x_i) \right]$$

Theorem 12 *For any anti-symmetric 1-Lipschitz function σ and any set $S = \{x_1, \dots, x_m\} \subseteq \mathbb{R}^D$:*

$$\mathcal{R}_m(\mathcal{N}_{\mu_{1,\infty} \leq \mu}^{d,\sigma}) \leq \sqrt{\frac{2\mu^{2d} \log(2D) \sup \|x_i\|_\infty^2}{m}}$$

Proof Assuming ξ is uniform over $\{\pm 1\}^m$, we have:

$$\begin{aligned}
 \mathcal{R}_m(\mathcal{N}_{\mu_1, \infty \leq \mu}^{d, H}) &= \mathbb{E}_\xi \left[\frac{1}{m} \sup_{f \in \mathcal{N}_{\mu_1, \infty \leq \mu}^{d, H}} \left| \sum_{i=1}^m \xi_i f(x_i) \right| \right] \\
 &= \mathbb{E}_\xi \left[\frac{1}{m} \sup_{f \in \mathcal{N}_{\mu_1, \infty \leq \mu}^{d, H}} \sum_{i=1}^m \xi_i f(x_i) \right] \\
 &= \mathbb{E}_\xi \left[\frac{1}{m} \sup_{g \in \mathcal{N}_{\mu_1, \infty \leq \mu}^{d-1, H, H}} \sup_{\|w\|_1 \leq \mu} w^\top \sum_{i=1}^m \xi_i \sigma(g(x_i)) \right] \\
 &= \mathbb{E}_\xi \left[\frac{1}{m} \sup_{g \in \mathcal{N}_{\mu_1, \infty \leq \mu}^{d-1, H, H}} \left\| \sum_{i=1}^m \xi_i \sigma(g(x_i)) \right\|_\infty \right] \\
 &= \mathbb{E}_\xi \left[\frac{1}{m} \sup_{g \in \mathcal{N}_{\mu_1, \infty \leq \mu}^{d-1, H}} \left| \sum_{i=1}^m \xi_i \sigma(g(x_i)) \right| \right] \tag{15}
 \end{aligned}$$

$$\begin{aligned}
 &= \mathbb{E}_\xi \left[\frac{1}{m} \sup_{g \in \mathcal{N}_{\mu_1, \infty \leq \mu}^{d-1, H}} \sum_{i=1}^m \xi_i \sigma(g(x_i)) \right] \\
 &\leq \mathbb{E}_\xi \left[\frac{1}{m} \sup_{g \in \mathcal{N}_{\mu_1, \infty \leq \mu}^{d-1, H}} \sum_{i=1}^m \xi_i g(x_i) \right] \tag{16} \\
 &= \mathbb{E}_\xi \left[\frac{1}{m} \sup_{g \in \mathcal{N}_{\mu_1, \infty \leq \mu}^{d-1, H}} \left| \sum_{i=1}^m \xi_i g(x_i) \right| \right] \\
 &= \mathcal{R}_m(\mathcal{N}_{\mu_1, \infty \leq \mu}^{d-1, H})
 \end{aligned}$$

where the equality (15) is by anti-symmetric property of σ and inequality (16) is by the version of Contraction Lemma without the absolute value. This will give us the bound on Rademacher complexity of $\mathcal{N}_{\mu_1, \infty \leq \mu}^{d, H}$ based on the Rademacher complexity of $\mathcal{N}_{\mu_1, \infty \leq \mu}^{d-1, H}$. Applying the same argument on all layers and using lemma 15 to bound the complexity of the first layer completes the proof. \blacksquare

Appendix B. Tightness

We repeat the theorem statement here for convenience.

Theorem 3 For any $p, q \geq 1$ (and $1/p^* + 1/p = 1$) and any depth $d \geq 2$, the $m = 2^D$ points $\{\pm 1\}^D$ can be shattered with unit margin by $\mathcal{N}_{\gamma_{p, q} \leq \gamma}^{d, H}$ with:

$$\gamma \leq D^{1/p} m^{1/p+1/q} H^{-(d-2)[1/p^*-1/q]_+}.$$

Proof Consider a size m subset S_m of 2^D vertices of the D dimensional hypercube $\{-1, +1\}^D$. We construct the first layer using m units. Each unit has a unique weight vector consisting of $+1$ and -1 's and will output a positive value if and only if the sign pattern of the input $x \in S_m$ matches that of the weight vector. The second layer has a single unit and connects to all m units in the first layer. For any m dimensional sign pattern $b \in \{-1, +1\}^m$, we can choose the weights of the second layer to be b , and the network will output the desired sign for each $x \in S_m$ with unit margin. The norm of the network is at most $(m \cdot D^{q/p})^{1/q} \cdot m^{1/p} = D^{1/p} \cdot m^{(1/p+1/q)}$. This establishes the claim for $d = 2$. For $d > 2$ and $1/p + 1/q \geq 1$, we obtain the same norm and unit margin by adding $d - 2$ layers with one unit in each layer connected to the previous layer by a unit weight. For $d > 2$ and $1/p + 1/q < 1$, we show the dependence on H by recursively replacing the top unit with H copies of it and adding an averaging unit on top of that. More specifically, given the above $d = 2$ layer network, we make H copies of the output unit with rectified linear activation and add a 3rd layer with one output unit with uniform weight $1/H$ to all the copies in the 2nd layer. Since this operation does not change the output of the network, we have the same margin and now the norm of the network is $(m \cdot D^{q/p})^{1/q} \cdot (Hm^{q/p})^{1/q} \cdot (H(1/H^p))^{1/p} = D^{1/p} \cdot m^{(1/p+1/q)} \cdot H^{1/q-1/p^*}$. That is, we have reduced the norm by factor $H^{1/q-1/p^*}$. By repeating this process, we get the geometric reduction in the norm $H^{(d-2)(1/q-1/p^*)}$, which concludes the proof. \blacksquare

Appendix C. Proof that $\gamma_{p,q}^d(f)$ is a semi-norm in \mathcal{N}^d

Theorem 4 For any $d, p, q \geq 1$ such that $\frac{1}{q} \leq \frac{1}{d-1}(1 - \frac{1}{p})$, $\gamma_{p,q}^d(f)$ is a semi-norm in \mathcal{N}^d .

Proof The proof consists of three parts. First we show that the level set $\mathcal{N}_{\gamma_{p,q}^d \leq \gamma}^d = \{f \in \mathcal{N}^d : \gamma_{p,q}^d(f) \leq \gamma\}$ is a convex set if the condition on d, p, q is satisfied. Next, we establish the non-negative homogeneity of $\gamma_{p,q}^d(f)$. Finally, we show that if a function $\alpha : \mathcal{N}^d \rightarrow \mathbb{R}$ is non-negative homogeneous and every sublevel set $\{f \in \mathcal{N}^d : \alpha(f) \leq \gamma\}$ is convex, then α satisfies the triangular inequality.

Convexity of the level sets First we show that for any two functions $f_1, f_2 \in \mathcal{N}_{\gamma_{p,q}^d \leq \gamma}^d$ and $0 \leq \alpha \leq 1$, the function $g = \alpha f_1 + (1 - \alpha)f_2$ is in the hypothesis class $\mathcal{N}_{\gamma_{p,q}^d \leq \gamma}^d$. We prove this by constructing weights W that realizes g . Let U and V be the weights of two neural networks such that $\gamma_{p,q}(U) = \gamma_{p,q}^d(f_1) \leq \gamma$ and $\gamma_{p,q}(V) = \gamma_{p,q}^d(f_2) \leq \gamma$. For every layer $i = 1, \dots, d$ let

$$\tilde{U}_i = \sqrt[d]{\gamma_{p,q}(U)U_i/\|U_i\|_{p,q}}, \quad \tilde{V}_i = \sqrt[d]{\gamma_{p,q}(V)V_i/\|V_i\|_{p,q}}$$

and set $W_1 = \begin{bmatrix} \tilde{U}_1 \\ \tilde{V}_1 \end{bmatrix}$ for the first layer, $W_i = \begin{bmatrix} \tilde{U}_i & 0 \\ 0 & \tilde{V}_i \end{bmatrix}$ for the intermediate layers and $W_d = \begin{bmatrix} \alpha\tilde{U}_d & (1-\alpha)\tilde{V}_d \end{bmatrix}$ for the output layer.

Then for the defined W , we have $f_W = \alpha f_1 + (1 - \alpha)f_2$ for rectified linear and any other non-negative homogeneous activation function. Moreover, for any $i < d$, the norm of each layer is

$$\|W_i\|_{p,q} = \left(\gamma_{p,q}(U)^{\frac{q}{d}} + \gamma_{p,q}(V)^{\frac{q}{d}} \right)^{\frac{1}{q}} \leq 2^{\frac{1}{q}} \gamma^{\frac{1}{d}} \quad (17)$$

and in layer d we have:

$$\|W_d\|_p = \left(\alpha^p \gamma_{p,q}(U)^{\frac{p}{d}} + (1 - \alpha)^p \gamma_{p,q}(V)^{\frac{p}{d}} \right)^{\frac{1}{p}} \leq 2^{1/p-1} \gamma^{1/d} \quad (18)$$

Combining inequalities (17) and (18), we get $\gamma_{p,q}^d(f_W) \leq 2^{\frac{d-1}{q} + \frac{1}{p}} \gamma \leq \gamma$, where the last inequality holds because we assume that $\frac{1}{q} \leq \frac{1}{d-1} (1 - \frac{1}{p})$. Thus for every $\gamma \geq 0$, $\mathcal{N}_{\gamma_{p,q}^d \leq \gamma}^d$ is a convex set.

Non-negative homogeneity For any function $f \in \mathcal{N}^d$ and any $\alpha \geq 0$, let U be the weights realizing f with $\gamma_{p,q}^d(f) = \gamma_{p,q}(U)$. Then $\sqrt[d]{\alpha}U$ realizes αf establishing $\gamma_{p,q}^d(\alpha f) \leq \gamma_{p,q}(\sqrt[d]{\alpha}U) = \alpha \gamma_{p,q}(U) = \alpha \gamma_{p,q}^d(f)$. This establishes the non-negative homogeneity of $\gamma_{p,q}^d$.

Convex sublevel sets and homogeneity imply triangular inequality Let $\alpha(f)$ be non-negative homogeneous and assume that every sublevel set $\{f \in \mathcal{N}^d : \alpha(f) \leq \gamma\}$ is convex. Then for $f_1, f_2 \in \mathcal{N}^d$, defining $\gamma_1 \triangleq \alpha(f_1)$, $\gamma_2 \triangleq \alpha(f_2)$, $\tilde{f}_1 \triangleq (\gamma_1 + \gamma_2)f_1/\gamma_1$, and $\tilde{f}_2 \triangleq (\gamma_1 + \gamma_2)f_2/\gamma_2$, we have

$$\alpha(f_1 + f_2) = \alpha \left(\frac{\gamma_1}{\gamma_1 + \gamma_2} \tilde{f}_1 + \frac{\gamma_2}{\gamma_1 + \gamma_2} \tilde{f}_2 \right) \leq \gamma_1 + \gamma_2 = \alpha(f_1) + \alpha(f_2).$$

Here the inequality is due to the convexity of the level set and the fact that $\alpha(\tilde{f}_1) = \alpha(\tilde{f}_2) = \gamma_1 + \gamma_2$, because of the homogeneity. Therefore α satisfies the triangular inequality and thus it is a seminorm. \blacksquare

Appendix D. Path Regularization

D.1. Theorem 5

Lemma 19 For any function $f \in \mathcal{N}_{\gamma_{p,\infty}^d \leq \gamma}^{d,H}$ there is a layered network with weights w such that $\gamma_{p,\infty}(w) = \gamma_{p,\infty}^d(f)$ and for any internal unit v , $\sum_{(u \rightarrow v) \in E} |w(u \rightarrow v)|^p = 1$.

Proof Let w be the weights of a network such that $\gamma_{p,\infty}(w) = \gamma_{p,\infty}^d(f)$. We now construct a network with weights \tilde{w} such that $\gamma_{p,\infty}(w) = \gamma_{p,\infty}^d(f)$ and for any internal unit v , $\sum_{(u \rightarrow v) \in E} |\tilde{w}(u \rightarrow v)|^p = 1$. We do this by an incremental algorithm. Let $w_0 = w$. At each step i , we do the following.

Consider the first layer, Set V_k to be the set of neurons in the layer k . Let x be the maximum of ℓ_p norms of input weights to each neuron in set V_1 and let $U_x \subseteq V_1$ be the set of neurons whose ℓ_p norms of their input weight is exactly x . Now let y be the maximum of ℓ_p norms of input weights to each neuron in the set $V_1 \setminus U_x$ and let U_y be the set of the neurons such that the ℓ_p norms of their input weights is exactly y . Clearly $y < x$. We now scale down the input weights of neurons in set U_x by y/x and scale up all the outgoing edges of vertices in U_x by x/y (y cannot be zero for internal neurons based on the definition). It is straightforward that the new network realizes the same function and the $\ell_{p,\infty}$ norm of the first layer has changed by a factor y/x . Now for every neuron $v \in V_2$, let $r(v)$ be the ℓ_p norm of the new incoming weights divided by ℓ_p norm of the original incoming weights. We know that $r(v) \leq x/y$. We again scale down the input weights of every $v \in V_2$ by $1/r(v)$ and scale up all the outgoing edges of v by $r(v)$. Continuing this operation to on each layer, each time we propagate the ratio to the next layer while the network always realizes

the same function and for each layer k , we know that for every $v \in V_k$, $r(v) \leq x/y$. After this operation, in the network, the $\ell_{p,\infty}$ norm of the first layer is scaled down by y/x while the $\ell_{p,\infty}$ norm of the last layer is scaled up by at most x/y and the $\ell_{p,\infty}$ norm of the rest of the layers has remained the same. Therefore, if w_i is the new weight setting, we have $\gamma_{p,\infty}(w_i) \leq \gamma_{p,\infty}(w_{i-1})$.

After continuing the above step at most $|V_1| - 1$ times, the ℓ_p norm of input weights is the same for all neurons in V_1 . We can then run the same algorithm on other layers and at the end we have a network with weight setting \tilde{w} such that the for each $k < d$, ℓ_p norm of input weight to each of the neurons in layer k is equal to each other and $\gamma_{p,\infty}(\tilde{w}) \leq \gamma_{p,\infty}(w)$. This is in fact an equality because weight setting w' realizes function f and we know that $\gamma_{p,\infty}(w) = \gamma_{p,\infty}^{d,H}(f)$. A simple scaling of weights in layers gives completes the proof. \blacksquare

Theorem 5 For $p \geq 1$, any d and (finite or infinite) H , for any $f \in \mathcal{N}^{d,H}$: $\phi_p^{d,H}(f) = \gamma_{p,\infty}^{d,H}$.

Proof By the Lemma 19, there is a layered network with weights \tilde{w} such that $\gamma_{p,\infty}(\tilde{w}) = \gamma_{p,\infty}^{d,H}(f)$ and for any internal unit v , $\sum_{(u \rightarrow v) \in E} |\tilde{w}(u \rightarrow v)|^p = 1$. Let W be the weights of the layered network that corresponds to the function \tilde{w} . Then we have:

$$v_p(\tilde{w}) = \left(\sum_{v_{\text{in}}[i] \xrightarrow{e_1} v_1 \xrightarrow{e_2} v_2 \cdots \xrightarrow{e_k} v_{\text{out}}} \prod_{i=1}^k |\tilde{w}(e_i)|^p \right)^{\frac{1}{p}} \quad (19)$$

$$= \left(\sum_{i_{d-1}=1}^H \cdots \sum_{i_1=1}^H \sum_{i_0=1}^D |W_d[i_{d-1}]|^p \prod_{k=1}^{d-1} |W_k[i_k, i_{k-1}]|^p \right)^{\frac{1}{p}} \quad (20)$$

$$= \left(\sum_{i_{d-1}=1}^H |W_d[i_{d-1}]|^p \cdots \sum_{i_1=1}^H |W_k[i_2, i_1]|^p \sum_{i_0=1}^D |W_k[i_1, i_0]|^p \right)^{\frac{1}{p}} \quad (21)$$

$$= \left(\sum_{i_{d-1}=1}^H |W_d[i_{d-1}]|^p \cdots \sum_{i_1=1}^H |W_k[i_2, i_1]|^p \right)^{\frac{1}{p}} \quad (22)$$

$$= \left(\sum_{i_{d-1}=1}^H |W_d[i_{d-1}]|^p \cdots \sum_{i_2=1}^H |W_k[i_3, i_2]|^p \right)^{\frac{1}{p}} \quad (23)$$

$$= \left(\sum_{i_{d-1}=1}^H |W_d[i_{d-1}]|^p \right)^{\frac{1}{p}} = \ell_p(W_d) = \gamma_{p,\infty}(W) \quad (24)$$

$$(25)$$

where inequalities 20 to 24 are due to the fact that the ℓ_p norm of input weights to each internal neuron is exactly 1 and the last equality is again because $\ell_{p,\infty}$ of all layers is exactly 1 except the layer d . \blacksquare

D.2. Proof of Theorem 6

In this section, without loss of generality, we assume that all the internal nodes in a DAG have incoming edges and outgoing edges because otherwise we can just discard them. Let $d_{\text{out}}(v)$ be the longest directed path from vertex v to v_{out} and $d_{\text{in}}(v)$ be the longest directed path from any input vertex $v_{\text{in}}[i]$ to v . We say graph G is a sublayered graph if G is a subgraph of a layered graph.

We first show the necessary and sufficient conditions under which a DAG is a sublayered graph.

Lemma 20 *The graph $G(E, V)$ is a sublayered graph if and only if any path from input nodes to the output nodes has length d where d is the length of the longest path in G*

Proof Since the internal nodes have incoming edges and outgoing edges; hence if G is a sublayered graph it is straightforward by induction on the layers that for every vertex v in layer i , there is a vertex u in layer $i + 1$ such that $(v \rightarrow u) \in E$ and this proves the necessary condition for being sublayered graph.

To show the sufficient condition, for any internal node u , u has $d_{\text{in}}(v)$ distance from the input node in every path that includes u (otherwise we can build a path that is longer than d). Therefore, for each vertex $v \in V$, we can place vertex v in layer $d_{\text{in}}(v)$ and all the outgoing edges from v will be to layer $d_{\text{in}}(v) + 1$. ■

Lemma 21 *If the graph $G(E, V)$ is not a sublayered graph then there exists a directed edge $(u \rightarrow v)$ such that $d_{\text{in}}(u) + d_{\text{out}}(v) < d - 1$ where d the length of the longest path in G .*

Proof We prove the lemma by an inductive argument. If G is not sublayered, by lemma 20, we know that there exists a path $v_0 \rightarrow \dots v_i \dots \rightarrow v_{d'}$ where v_0 is an input node ($d_{\text{in}}(v_0) = 0$), $v_{d'} = v_{\text{out}}$ ($d_{\text{out}}(v_{d'}) = 0$) and $d' < d$. Now consider the vertex v_1 . We need to have $d_{\text{out}}(v_1) = d - 1$ otherwise if $d_{\text{out}}(v_1) < d - 1$ we get $d_{\text{in}}(u) + d_{\text{out}}(v) < d - 1$ and if $d_{\text{out}}(v_1) > d - 1$ there will be path in G that is longer than d . Also, since $d_{\text{out}}(v_1) = d - 1$ and the longest path in G has length d , we have $d_{\text{in}}(v_1) = 1$.

By applying the same inductive argument on each vertex v_i in the path we get $d_{\text{in}}(v_i) = i$ and $d_{\text{out}}(v_i) = d - i$. Note that if the condition $d_{\text{in}}(u) + d_{\text{out}}(v) < d - 1$ is not satisfied in one of the steps of the inductive argument, the lemma is proved. Otherwise, we have $d_{\text{in}}(v_{d'-1}) = d' - 1$ and $d_{\text{out}}(v_{d'-1}) = d - d' + 1$ and therefore $d_{\text{in}}(v_{d'-1}) + d_{\text{out}}(v_{\text{out}}) = d' - 1 < d - 1$ that proves the lemma. ■

Theorem 6 *For any $p \geq 1$ and any d : $\gamma_{p,\infty}^d(f) = \min_{G \in \text{DAG}(d)} \phi_p^G(f)$.*

Proof Consider any $f_{G,w} \in \mathcal{N}^{\text{DAG}(d)}$ where the graph $G(E, V)$ is not sublayered. Let ρ be the total number of paths from input nodes to the output nodes. Let T be sum over paths of the length of the path. We indicate an algorithm to change G into a sublayered graph \tilde{G} of depth d with weights \tilde{w} such that $f_{G,w} = f_{\tilde{G},\tilde{w}}$ and $\phi(w) = \phi(\tilde{w})$. Let $G_0 = G$ and $w_0 = w$.

At each step i , we consider the graph G_{i-1} . If G_{i-1} is sublayered, we are done otherwise by lemma 21, there exists an edge $(u \rightarrow v)$ such that $d_{\text{in}}(u) + d_{\text{out}}(v) < d - 1$. Now we add a new vertex \tilde{v}_i to graph G_{i-1} , remove the edge $(u \rightarrow v)$, add two edges $(u \rightarrow \tilde{v}_i)$ and $(\tilde{v}_i \rightarrow v)$ and return the

graph as G_i and since we had $d_{\text{in}}(u) + d_{\text{out}}(v) < d - 1$ in G_{i-1} , the longest path in G_i still has length d . We also set $w(u \rightarrow \tilde{v}_i) = \sqrt{|w(u \rightarrow v)|}$ and $w(\tilde{v}_i \rightarrow v) = \text{sign}(w(u \rightarrow v))\sqrt{|w(u \rightarrow v)|}$. Since we are using rectified linear units activations, for any $x > 0$, we have $[x]_+ = x$ and therefore:

$$\begin{aligned} w(\tilde{v}_i \rightarrow v) [w(u \rightarrow \tilde{v}_i) o(u)]_+ &= \text{sign}(w(u \rightarrow v)) \sqrt{|w(u \rightarrow v)|} \left[\sqrt{|w(u \rightarrow v)|} o(u) \right]_+ \\ &= \text{sign}(w(u \rightarrow v)) \sqrt{|w(u \rightarrow v)|} \sqrt{|w(u \rightarrow v)|} o(u) \\ &= w(u \rightarrow v) o(u) \end{aligned}$$

So we conclude that $f_{G_i, w_i} = f_{G_{i-1}, w_{i-1}}$. Clearly, since we didn't change the length of any path from input vertices to the output vertex, we have $\phi(w) = \phi(\tilde{w})$. Let T_i be sum over paths of the length of the path in G_i . It is clear that $T_{i-1} \leq T_i$ because we add a new edge into a path at each step. We also know by lemma 20 that if $T_i = \rho d$, then G_i is a sublayered graph. Therefore, after at most $\rho d - T_0$ steps, we return a sublayered graph \tilde{G} and weights \tilde{w} such that $f_{G, w} = f_{\tilde{G}, \tilde{w}}$. We can easily turn the sublayered graph \tilde{G} a layered graph by adding edges with zero weights and this together with Theorem 5 completes the proof. ■

D.3. Proof of Theorem 9

Theorem 9 *For any $p \geq 1$ and $d > 1$ and any $f \in \mathcal{N}^d$, there exists a layered graph $G(V, E)$ of depth d , such that $f \in \mathcal{N}^G$ and $\gamma_{p, \infty}^G(f) = \phi_p^G(f) = \gamma_{p, \infty}^d(f)$, and the out-degree of every internal (non-input) node in G is one. That is, the subgraph of G induced by the non-input vertices is a tree directed toward the output vertex.*

Proof For any $f_{G, w} \in \mathcal{N}^{\text{DAG}(d)}$, we show how to construct such \tilde{G} and \tilde{w} . We first sort the vertices of G based on topological ordering such that the out-degree of the first vertex is zero. Let $G_0 = G$ and $w_0 = w$. At each step i , we first set $G_i = G_{i-1}$ and $w_i = w_{i-1}$ and then pick the vertex u that is the i th vector in the topological ordering. If the out-degree of u is at most 1. Otherwise, for any edge $(u \rightarrow v)$ we create a copy of vertex u that we call it u_v , add the edge $(u_v \rightarrow v)$ to G_i and connect all incoming edges of u with the same weights to every such u_v and finally we delete the vertex u from G_i together with all incoming and outgoing edges of u . It is easy to indicate that $f_{G_i, w_i} = f_{G_{i-1}, w_{i-1}}$. After at most $|V|$ such steps, all internal nodes have out-degree one and hence the subgraph induced by non-input vertices will be a tree. ■

Appendix E. Hardness of Learning Neural Networks

Daniely et al. (2014) show in Theorem 5.4 and in Section 7.2 that subject to the strong random CSP assumption, for any $k = \omega(1)$ the hypothesis class of intersection of homogeneous halfspaces over $\{\pm 1\}^n$ with normals in $\{\pm 1\}$ is not efficiently PAC learnable (even improperly)². Furthermore, for any $\epsilon > 0$, Klivans and Sherstov (2006) prove this hardness result subject to intractability of $\tilde{Q}(D^{1.5})$ -unique shortest vector problem for $k = D^\epsilon$.

2. Their Theorem 5.4 talks about unrestricted halfspaces, but the construction in Section 7.2 uses only data in $\{\pm 1\}^D$ and halfspaces specified by $\langle w, x \rangle > 0$ with $w \in \{\pm 1\}^D$

If it is not possible to efficiently PAC learn intersection of halfspaces (even improperly), we can conclude it is also not possible to efficiently PAC learn any hypothesis class which can represent such intersection. In Theorem 22 we show that intersection of homogeneous half spaces can be realized with unit margin by neural networks with bound norm.

Theorem 22 *For any $k > 0$, the intersection of k homogeneous half spaces is realizable with unit margin by $\mathcal{N}_{\gamma_{p,q}}^2$ where $\gamma = 4D^{\frac{1}{p}}k^2$.*

Proof The proof is by a construction that is similar to the one in Livni et al. (2014). For each hyperplane $\langle w_i, x \rangle > 0$, where $w_i \in \{\pm 1\}^D$, we include two units in the first layer: $g_i^+(x) = [\langle w_i, x \rangle]_+$ and $g_i^-(x) = [\langle w_i, x \rangle - 1]_+$. We set all incoming weights of the output node to be 1. Therefore, this network is realizing the following function:

$$f(x) = \sum_{i=1}^k ([\langle w_i, x \rangle]_+ - [\langle w_i, x \rangle - 1]_+)$$

Since all inputs and all weights are integer, the outputs of the first layer will be integer, $([\langle w_i, x \rangle]_+ - [\langle w_i, x \rangle - 1]_+)$ will be zero or one, and f realizes the intersection of the k halfspaces with unit margin. Now, we just need to make sure that $\gamma_{p,q}^2(f)$ is bounded by $\gamma = 4D^{\frac{1}{p}}k^2$:

$$\begin{aligned} \gamma_{p,q}^2(f) &= D^{\frac{1}{p}}(2k)^{\frac{1}{q}}(2k)^{\frac{1}{p}} \\ &\leq D^{\frac{1}{p}}(2k)^2 = \gamma. \end{aligned}$$

■