

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/339761758>

Measuring evidence: a probabilistic approach to an extension of Belnap–Dunn logic

Article in *Synthese* · March 2020

DOI: 10.1007/s11229-020-02571-w

CITATIONS

3

READS

33

3 authors:



Abilio Rodrigues

Federal University of Minas Gerais

22 PUBLICATIONS 64 CITATIONS

SEE PROFILE



Juliana Bueno-Soler

Universidade Federal do ABC (UFABC)

16 PUBLICATIONS 72 CITATIONS

SEE PROFILE



Walter Carnielli

Centre for Logic, Epistemology and the History of Science

70 PUBLICATIONS 165 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Brouwerian Relevance Logic [View project](#)



Paraconsistent probabilities and possibilistic measures [View project](#)

Measuring evidence: a probabilistic approach to an extension of Belnap-Dunn Logic *

Abilio Rodrigues¹, Juliana Bueno-Soler², Walter Carnielli²

¹Federal University of Minas Gerais
abilio.rodrigues@gmail.com

²University of Campinas
juliana@ft.unicamp.br
walter.carnielli@gmail.com

February 27, 2020

Abstract

This paper introduces the logic of evidence and truth LET_F as an extension of the Belnap-Dunn four-valued logic FDE . LET_F is a slightly modified version of the logic LET_J , presented in Carnielli and Rodrigues (2017). While LET_J is equipped only with a classicality operator \circ , LET_F is equipped with a non-classicality operator \bullet as well, dual to \circ . Both LET_F and LET_J are logics of formal inconsistency and undeterminedness in which the operator \circ recovers classical logic for propositions in its scope. Evidence is a notion weaker than truth in the sense that there may be evidence for a proposition α even if α is not true. As well as LET_J , LET_F is able to express preservation of evidence and preservation of truth. The primary aim of this paper is to propose a probabilistic semantics for LET_F where statements $P(\alpha)$ and $P(\circ\alpha)$ express, respectively, the amount of evidence available for α and the degree to which the evidence for α is expected to behave classically – or non-classically for $P(\bullet\alpha)$. A probabilistic scenario is paracomplete when $P(\alpha) + P(\neg\alpha) < 1$, and paraconsistent when $P(\alpha) + P(\neg\alpha) > 1$, and in both cases, $P(\circ\alpha) < 1$. If $P(\circ\alpha) = 1$, or $P(\bullet\alpha) = 0$, classical probability is recovered for α . The proposition $\circ\alpha \vee \bullet\alpha$, a theorem of LET_F , partitions what we call the information space, and thus allows us to obtain some new versions of known results of standard probability theory.

*The authors acknowledge support from the National Council for Scientific and Technological Development (CNPq, Brazil) under research grants 311911/2018-8, 308077/2018-0, and 307376/2018-4. The first author has also been supported by the CNPq fellowship grant 204479/2017-71 for a research stay at the University of Oxford, 2018. The authors owe a great debt to the referees for the careful reading of an earlier version of this text, which lead to relevant suggestions for improving it.

1 Introduction

In Carnielli and Rodrigues (2017) two paraconsistent and paracomplete formal systems were presented, the Basic Logic of Evidence (*BLE*) and the Logic of Evidence and Truth (*LET_J*). *BLE* ends up being equivalent to Nelson’s well-known logic *N4* but has been conceived to express preservation of evidence instead of truth. Scenarios with conflicting evidence – that is, non-conclusive evidence for the truth and the falsity of α – as well as scenarios with no evidence at all about α are possible, so neither explosion nor excluded middle hold in *BLE*. *LET_J* is an extension of *BLE* equipped with a classicality operator \circ . When $\circ\alpha$ holds, classical negation – and so full classical logic – for α is recovered. According to the intended interpretation, $\circ\alpha$ in *LET_J* means that there is conclusive evidence for the truth or falsity of α , so the truth-value of α has been established as true or false.

Let us call \vdash_C and \vdash_{BLE} , respectively, the relation of logical consequence in classical logic and in *BLE*. Classical consequence is defined in terms of preservation of truth: $\Gamma \vdash_C \alpha$ just in case there is no model M such that all propositions of Γ are true in M , but α is not true in M . The intended interpretation of *BLE*, on the other hand, is not based on preservation of truth, but rather on preservation of evidence: $\Gamma \vdash_{BLE} \alpha$ means that the availability of evidence for the premises in Γ implies that there is also evidence available for α . Classical logic and *BLE*, therefore, express different properties of propositions: truth and availability of evidence. The logic *LET_J*, in its turn, is able to express preservation of evidence and preservation of truth – it ‘combines’, in one and the same formal system, the relations \vdash_C and \vdash_{BLE} . The operator \circ works like a *context switch* that divides propositions into those that have a classical and those that have a non-classical behavior, and *BLE* is the underlying logic of the latter.

Adequate valuation semantics and decision procedures for *BLE* and *LET_J* have been proposed. These semantics, however, are only able to express the fact that a given proposition α has or does not have evidence available by attributing, respectively, the semantic value 1 or 0 to α . Evidence, thus, is treated from a purely qualitative point of view. A question that presents itself is whether the amount of evidence available for a given proposition α could be quantified. Here we give a positive answer to this question.

The aim of this paper is to propose a probabilistic semantics for a modified version of *LET_J* obtained by dropping the implication symbol \rightarrow and adding a non-classicality operator \bullet dual to \circ . While $\circ\alpha$ implies that α behaves classically, a non-classical behavior of α implies $\bullet\alpha$. The logic so obtained is an extension of the well-known logic of First-Degree Entailment (*FDE*), and we call it *LET_F*, the Logic of Evidence and Truth based on *FDE*. As well as *LET_J*, *LET_F* is suitable to an intuitive reading in terms of evidence and truth.

In order to capture this idea of preservation of degrees of evidence a non-classical notion of probability will be employed. The probabilistic semantics proposed here follows the ideas presented in Bueno-Soler and Carnielli (2016, 2017). Let $P(\alpha) = \epsilon$ mean that ϵ is the measure of evidence available for α . We call a probabilistic scenario *paracomplete* when $P(\alpha) + P(-\alpha) < 1$, and

46 *paraconsistent* when $P(\alpha) + P(\neg\alpha) > 1$. These two cases can be explained,
 47 respectively, as ‘too little information’ and ‘too much information’ about α .¹ In
 48 both cases, $P(\circ\alpha) < 1$, which means that the probability measures of α and $\neg\alpha$
 49 are not behaving classically. So, $P(\circ\alpha) < 1$ means that the information available
 50 about α is not reliable, and something must be wrong. If $P(\circ\alpha) = 1$, standard
 51 probability is recovered for α .

52 With the purpose of understanding the probabilistic semantics proposed here
 53 better, we adopt a notion of *information space* instead of the standard notion
 54 of sample space. The intuitive idea is to collect all the relevant information
 55 about a proposition α (or about a set of propositions Γ) and the corresponding
 56 measures of evidence. So, roughly speaking, an information space is constituted
 57 by propositions that represent evidence that can be non-conclusive, contradic-
 58 tory or incomplete, more reliable or less reliable, and sometimes conclusive (we
 59 return to this point in Section 4.3 below). Such a notion of information space
 60 requires a generalization of the notion of a partition, and consequently allows
 61 us to obtain generalized versions of standard results of probability theory such
 62 as total probability theorem and Bayes’ rule.²

63 The remainder of this paper is organized in four sections. Section 2 is dedi-
 64 cated to the logic *FDE*. It is shown that *FDE* is suited to an interpretation in
 65 terms of preservation of evidence. We also present adequate valuation semantics
 66 and a decision procedure for *FDE*. In Section 3, *FDE* is extended to *LET_F*,
 67 and an adequate semantics, a decision procedure, and some relevant results are
 68 presented and discussed. In Section 4, a probabilistic semantics for *LET_F*
 69 is defined, and paraconsistent and paracomplete versions of total probability the-
 70 orems and Bayes’ rule are also presented and discussed. Finally, in Section 5, we
 71 discuss some points related to the topics of this paper that could be developed
 72 further.

73 2 *FDE* as a logic of preservation of evidence

74 The inference rules of *BLE* were obtained by asking whether an inference rule
 75 preserves evidence. Since evidence can be incomplete (no evidence at all) and
 76 contradictory (conflicting evidence), explosion and excluded middle do not hold.
 77 In *BLE*, when α (resp. $\neg\alpha$) holds, the intended meaning is that there is evidence
 78 for the truth (resp. falsity) of α . Evidence that α is true and evidence that α

¹The connections between the notions of evidence and information will be explained in Section 2.2.1.

²Our approach differs from the so-called Dempster-Shafer (DS) theory of evidence, developed by Glenn Shafer in Shafer (1976) and based on earlier work of Arthur Dempster. DS is focused on degrees of belief and degrees of plausibility. As Lofti Zadeh points out in his review (Zadeh, 1984), the DS theory falls short as a useful tool for the management of uncertainty (even for expert systems, for which it was designed). Our approach, as we try to make clear throughout this paper, uses probabilistic semantics intended to quantify the evidence attributed to a proposition and introduces a new logic with an intuitive reading in terms of preservation of evidence and truth. That is the reason we cannot rely on the DS ‘mathematical’ theory of evidence: it is not so attractive as it seems to be at first glance, and lacks the features we are interested in.

79 is false are independent of each other, and are treated as such by the formal
80 system. *BLE* can express the following four scenarios:

- 81 1. Only evidence that α is true: α holds, $\neg\alpha$ does not hold.
- 82 2. Only evidence that α is false: $\neg\alpha$ holds, α does not hold.
- 83 3. No evidence at all: neither α nor $\neg\alpha$ hold.
- 84 4. Conflicting evidence: both α and $\neg\alpha$ hold.³

85 Evidence for a proposition α is explained in Carnielli and Rodrigues (2017,
86 Section 2) as reasons for believing in α , but these reasons may be non-conclusive
87 or even wrong, and do not imply the truth of α , nor the belief in α . Thus,
88 evidence is a notion weaker than truth in the sense that there may be evidence
89 for a proposition α even if α is not true. Below, in Section 2.2.1, starting from
90 the notion of information proposed by Dunn (2008), we explain evidence in
91 terms of a (perhaps) non-conclusive justification added to a proposition α or, as
92 Fitting (2016b) puts it, “justifications that might be wrong”.⁴ Notice that the
93 notion of evidence encompasses non-conclusive as well as conclusive evidence,
94 and the latter is evidence that establishes the truth-value of a proposition α .

95 The logic of First-Degree Entailment (*FDE*) is a paraconsistent and para-
96 complete propositional logic in a language with conjunction, disjunction, and
97 negation, with no theorems nor bottom particles (cf. Anderson and Belnap,
98 1963, 1975; Anderson et al., 1992; Belnap, 1977a,b; Dunn, 1976). *FDE* is a
99 fragment of *BLE/N4*, obtained by dropping the implication symbol and the
100 corresponding rules, and it can be interpreted in terms of preservation of evi-
101 dence, as well as *BLE* – the four scenarios above clearly correspond to the four
102 truth-values proposed by Belnap (1977a,b) (we return to this point in Section 2.2
103 below).⁵

104 **Definition 1.** *The Logic of First-Degree Entailment (FDE)*

105 *Let L_1 be a language with a denumerable set of sentential letters $\{p_1, p_2, p_3, \dots\}$,*
106 *the set of connectives $\{\neg, \wedge, \vee\}$, and parentheses. The set of formulas of L_1*

³The expression ‘ α holds/does not hold’ here means that α holds/does not hold in *BLE*.
So, here, it does not mean that α is true/false.

⁴Fitting (2016a) presents an embedding of *BLE* into the modal logic *KX4*, and an em-
bedding of the later into the justification logic *JX4*. The latter is equipped with justification
terms that stand for “justification, or evidence, which may be non-factual, uncertain, or con-
tradictory” (Fitting, 2016a, p. 1159). In *JX4*, ‘ $t : \alpha$ ’ means that α is justified by reason t .
The notion of evidence expressed by *KX4* (implicit evidence) and *JX4* (explicit evidence) is
a “formal alternative” of the “informal” notion of evidence expressed by *BLE*.

⁵The move from *BLE* and *LET_J* to (respectively) *FDE* and *LET_F* has been motivated by
some difficulties in interpreting the implication of *BLE* in probabilistic terms. The implication
of *BLE* is located somewhere in between classical and intuitionistic implication: it is not
classical because Peirce’s Law does not hold, and it is not intuitionistic because the equivalence
between $\neg(\alpha \rightarrow \beta)$ and $\alpha \wedge \neg\beta$ holds. It is not clear what would be the intuitive meaning of
the attribution of a probabilistic measure to a formula $\alpha \rightarrow \beta$ of *BLE*, and how this measure
would relate to the probabilistic values of α and β . So we decided, at least in this paper, to
work with *FDE*, the implication-free fragment of *BLE*.

107 is obtained recursively in the usual way. The logic FDE is defined over the
 108 language L_1 by the following natural deduction rules:

$$\begin{array}{c}
 109 \quad \frac{\alpha \quad \beta}{\alpha \wedge \beta} \wedge I \quad \frac{\alpha \wedge \beta}{\alpha} \wedge E \quad \frac{\alpha \wedge \beta}{\beta} \\
 110 \\
 111 \quad \frac{\alpha}{\alpha \vee \beta} \vee I \quad \frac{\beta}{\alpha \vee \beta} \quad \frac{\begin{array}{c} [\alpha] \\ \vdots \\ \gamma \end{array} \quad \begin{array}{c} [\beta] \\ \vdots \\ \gamma \end{array}}{\gamma} \vee E \\
 112 \\
 113 \quad \frac{\neg \alpha}{\neg(\alpha \wedge \beta)} \neg \wedge I \quad \frac{\neg \beta}{\neg(\alpha \wedge \beta)} \quad \frac{\begin{array}{c} [\neg \alpha] \\ \vdots \\ \gamma \end{array} \quad \begin{array}{c} [\neg \beta] \\ \vdots \\ \gamma \end{array}}{\gamma} \neg \wedge E \\
 114 \\
 115 \quad \frac{\neg \alpha \quad \neg \beta}{\neg(\alpha \vee \beta)} \neg \vee I \quad \frac{\neg(\alpha \vee \beta)}{\neg \alpha} \neg \vee E \quad \frac{\neg(\alpha \vee \beta)}{\neg \beta} \\
 116 \\
 117 \quad \frac{\alpha}{\neg \neg \alpha} DN \quad \frac{\neg \neg \alpha}{\alpha}
 \end{array}$$

118 A deduction of α from a set of premises Γ , $\Gamma \vdash_{FDE} \alpha$, is defined as follows:
 119 there is a derivation with conclusion α and all uncanceled hypotheses in Γ , and
 120 the definition of a derivation is the usual one for natural deduction systems (see
 121 e.g. van Dalen (2008, pp. 35-36)).

122 Other deductive systems have already been presented for FDE (see Omori and
 123 Wansing, 2017, Section 2.2), but the natural deduction system proposed here
 124 makes the symmetry between positive and negative rules explicit: $\wedge I$ and $\neg \vee I$
 125 are symmetrical, $\vee E$ and $\neg \wedge E$ are symmetrical, and so on. This mirrors the
 126 fact that positive and negative evidence are primitive and non-complementary
 127 notions, but have symmetric deductive behavior: the rule $\wedge I$ expresses the idea
 128 that when there is positive evidence available for both α and β , there is positive
 129 evidence for $\alpha \wedge \beta$, while the rule $\neg \vee I$ means that when there is negative evidence
 130 available for both α and β , there is negative evidence for $\alpha \vee \beta$.

131 **Theorem 2.**

132 *Reflexivity, monotonicity, transitivity, and compactness hold for FDE .*

133 *Proof.* These well-known properties of FDE can be easily proved by means of
 134 the natural deduction system above. \square

135 **2.1 Valuation semantics for FDE**

136 We now propose a non-deterministic valuation semantic for FDE .

137 **Definition 3.** *Valuation semantics for FDE*

138 *A valuation semantics for FDE is a collection of FDE -valuations defined as fol-*
 139 *lows: A function $v : L_1 \rightarrow \{0, 1\}$ is a FDE -valuation if it satisfies the following*
 140 *clauses:*

- 141 $v1. v(\alpha \wedge \beta) = 1$ iff $v(\alpha) = 1$ and $v(\beta) = 1$,
- 142 $v2. v(\alpha \vee \beta) = 1$ iff $v(\alpha) = 1$ or $v(\beta) = 1$,
- 143 $v3. v(\neg(\alpha \wedge \beta)) = 1$ iff $v(\neg\alpha) = 1$ or $v(\neg\beta) = 1$,
- 144 $v4. v(\neg(\alpha \vee \beta)) = 1$ iff $v(\neg\alpha) = 1$ and $v(\neg\beta) = 1$,
- 145 $v5. v(\alpha) = 1$ iff $v(\neg\neg\alpha) = 1$.

146 **Definition 4.** We say that a formula α is a *semantical consequence* of Γ ,
 147 $\Gamma \models_{FDE} \alpha$, iff for every valuation v , if $v(\beta) = 1$ for all $\beta \in \Gamma$, then $v(\alpha) = 1$.

148 This semantics is sound and complete, and provides a decision procedure for
 149 *FDE*. From now on, in this section, when there is no risk of ambiguity, we will
 150 just write \vdash and \models in the place of \vdash_{FDE} and \models_{FDE} .

151 **Theorem 5.** *Soundness*

152 Let Γ be a set of formulas, and α a formula of *FDE*. So, $\Gamma \vdash \alpha$ implies $\Gamma \models \alpha$.

153 *Proof.* The proof is routine. It shows that assuming there are sound derivations
 154 for the premise(s), the derivation obtained by the application of a rule is sound.
 155 □

156 **Theorem 6.** *Completeness*

157 Let Γ be a set of formulas, and α a formula of *FDE*. Then $\Gamma \models \alpha$ implies $\Gamma \vdash \alpha$.

158 *Proof.* Completeness can be proved by a Henkin-style proof. Given Γ and α
 159 such that $\Gamma \not\models \alpha$, a set Δ maximal w.r.t α can be obtained in the usual way. So,
 160 the proof of the following propositions is straightforward:

- 161 $v1'. \alpha \wedge \beta \in \Delta$ iff $\alpha \in \Delta$ and $\beta \in \Delta$;
- 162 $v2'. \alpha \vee \beta \in \Delta$ iff $\alpha \in \Delta$ or $\beta \in \Delta$;
- 163 $v3'. \neg(\alpha \wedge \beta) \in \Delta$ iff $\neg\alpha \in \Delta$ or $\neg\beta \in \Delta$;
- 164 $v4'. \neg(\alpha \vee \beta) \in \Delta$ iff $\neg\alpha \in \Delta$ and $\neg\beta \in \Delta$;
- 165 $v5'. \alpha \in \Delta$ iff $\neg\neg\alpha \in \Delta$.

166 Let v be the mapping from the language L_1 to $\{0, 1\}$ defined as follows: for
 167 every $\gamma \in L_1$, $v(\gamma) = 1$ iff $\gamma \in \Delta$. v is a valuation for *FDE* such that: for every
 168 $\beta \in \Gamma$, $v(\beta) = 1$, since $\Gamma \subseteq \Delta$; but $v(\alpha) = 0$, since $\alpha \notin \Delta$ (Δ is maximal w.r.t. α).
 169 Therefore, $\Gamma \not\models \alpha$. □

170
 171 The valuation semantics proposed in Definition 3 is non-deterministic in the
 172 sense that the semantic value of a formula $\neg\alpha$ is not a function of the value of

173 α . The possible values a formula can receive are given by quasi-matrices.⁶ In
 174 Example 7 below, we illustrate how a quasi-matrix works.

175 **Example 7.** *In FDE:*

- 176 1. $p, \neg p \vee q \neq q$;
- 177 2. $p, \neg(p \wedge q) \neq \neg q$;
- 178 3. $\neg p \wedge \neg q \neq \neg(p \vee q)$;
- 179 4. $\neg p \vee \neg q \neq \neg(p \wedge q)$.

180 *Proof.* Consider the following quasi-matrix:

	0								1							
p																
$\neg p$	0				1				0				1			
q	0		1		0		1		0		1		0		1	
$\neg q$	0	1	0	1	0	1	0	1	0	1	0	1	0	1	0	1
$\neg p \vee q$	0	0	1	1	1	1	1	1	0	0	1	1	1	1	1	1
$\neg(p \wedge q)$	0	1	0	1	1	1	1	1	0	1	0	1	1	1	1	1
<i>valuation</i>	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16

183 The valuations 13 and 14 above show that 1 is invalid, and the valuations 13
 184 and 15 show that 2 is invalid. The remaining cases (De Morgan laws) are left
 185 to the reader. \square

186 **Remark 8.** *The first four rows of the quasi-matrix above display the semantic*
 187 *values of the propositional variables and the negations of propositional variables*
 188 *that occur in the formulas at stake. The 5th and 6th rows are given by clauses*
 189 *v_2 and v_3 of Definition 3. Note that the semantic value of $\neg p$ is not determined*
 190 *by the value of p : the value of $\neg p$ bifurcates into 0 and 1 below $v(p) = 1$ and*
 191 *also below $v(p) = 0$. So, being n the number of propositional variables of a given*
 192 *formula, the number of valuations is finite and bounded by 2^{2n} . It is intuitively*
 193 *clear that the valuation semantics provides a decision procedure for FDE. A*
 194 *detailed algorithm, however, will be presented elsewhere.*

195 2.1.1 Some facts about FDE

196 **Fact 9.** *Modus ponens and the deduction theorem do not hold in FDE for an*
 197 *implication $\alpha \rightarrow \beta$ defined as $\neg\alpha \vee \beta$.*

198 *Proof.* That disjunctive syllogism does not hold in FDE is shown by the fol-
 199 lowing valuation: $v(\alpha) = 1$, $v(\neg\alpha) = 1$, $v(\beta) = 0$. In order to show that the
 200 deduction theorem does not hold, suppose $\Gamma, \alpha \vDash \beta$ implies $\Gamma \vDash \neg\alpha \vee \beta$. So, from
 201 $\alpha \vDash \alpha$ we would get $\vDash \neg\alpha \vee \alpha$, but the latter is invalid in FDE. \square

⁶A quasi-matrix is a non-deterministic matrix that represents non-deterministic valuation semantics. The notion of quasi-matrix was introduced by da Costa and Alves in da Costa and Alves (1977), where a valuation semantics was proposed for da Costa's logic C_1 (in da Costa and Alves (1977, p. 624, Def. 11) a detailed explanation of how to construct a quasi-matrix for C_1 can be found). See also Loparic (1986, 2010); Loparic and Alves (1979), where decision procedures based on quasi-matrices are provided for da Costa's C_ω and for intuitionistic logic.

202 **Fact 10.** *Grounding of contradictoriness*
 203 *A compound formula α is contradictory in a valuation v , i.e. $v(\alpha) = 1$ and*
 204 *$v(\neg\alpha) = 1$, only if at least one propositional letter p that occurs in α is contra-*
 205 *dictory in v .*

206 *Proof.* Suppose there is a valuation v such that $v(\alpha) = v(\neg\alpha) = 1$. We prove
 207 that there is at least one propositional letter p in α such that $v(p) = v(\neg p) = 1$.
 208 If $\alpha = p$, clearly, $v(\alpha) = v(\neg\alpha) = v(p) = v(\neg p) = 1$. The remaining cases are
 209 proved by induction on the complexity of α .

210 Case 1. $\alpha = \neg\neg\beta$. I.H.: if $v(\beta) = v(\neg\beta) = 1$, there is a p in β such that
 211 $v(p) = v(\neg p) = 1$. Suppose $v(\neg\neg\beta) = v(\neg\neg\neg\beta) = 1$. So, by Definition 3, $v(\beta) =$
 212 $v(\neg\beta) = 1$. The result follows by the inductive hypothesis.

213 Case 2. $\alpha = \beta \wedge \gamma$. I.H.: if $v(\beta) = v(\neg\beta) = 1$, there is a p in β such that
 214 $v(p) = v(\neg p) = 1$; *mutatis mutandis* for γ . Suppose $v(\beta \wedge \gamma) = v(\neg(\beta \wedge \gamma)) = 1$.
 215 So, by Definition 3, $v(\beta) = v(\gamma) = 1$, and either $v(\neg\beta) = 1$ or $v(\neg\gamma) = 1$. By the
 216 inductive hypothesis, there is a p either in β or in γ such that $v(p) = v(\neg p) = 1$.
 217 The remaining cases are left to the reader. \square

218 **Fact 11.** *Grounding of incompleteness*
 219 *A compound formula α is incomplete in a valuation v , i.e. $v(\alpha) = 0$ and*
 220 *$v(\neg\alpha) = 0$, only if at least one propositional letter p that occurs in α is incom-*
 221 *plete in v .*

222 *Proof.* Similar to the proof of Fact 10 above. \square

223 It is to be noted that the converse of Facts 10 and 11 do not hold: there may
 224 be a contradictory (resp. incomplete) atom p in a formula α without α being
 225 contradictory (resp. incomplete). Let α be the formula $p \vee q$ and consider the
 226 valuation v such that $v(p) = v(\neg p) = 1$, $v(q) = 1$ and $v(\neg q) = 0$. In this case,
 227 p is a contradictory propositional letter, but $p \vee q$ is not contradictory. On the
 228 other hand, in the valuation $v(p) = v(\neg p) = 0$, $v(q) = 1$ and $v(\neg q) = 0$, p is a
 229 incomplete propositional letter, but $p \vee q$ is not incomplete. Both valuations
 230 make $v(p \vee q) = 1$ and $v(\neg(p \vee q)) = 0$.

231 2.2 Equivalence with Belnap's four-valued and Dunn's re- 232 lational semantics

233 The valuation semantics proposed above, as expected, is equivalent both to the
 234 two-valued relational semantics proposed by Dunn (1976) and to the four-valued
 235 semantics presented by Belnap (1977b).⁷

236 **Definition 12.** *Dunn's relational semantics for FDE*
 237 *A Dunn-interpretation for FDE is a relation ρ between the set of formulas of*
 238 *FDE and the values T and F , $\rho \subseteq L \times \{T, F\}$, satisfying the following clauses:*

⁷The literature has a variety of algorithmic procedures that provide translations between finite-valued semantics and valuation semantics. One of them is given in Caleiro, Carnielli, Coniglio, and Marcos (2005). For the ease of the reader, however, we give below a direct proof of the equivalence between FDE-valuations, Dunn's and Belnap's semantics for FDE.

- 239 1. $\neg\alpha\rho T$ iff $\alpha\rho F$,
- 240 2. $\neg\alpha\rho F$ iff $\alpha\rho T$,
- 241 3. $(\alpha \wedge \beta)\rho T$ iff $\alpha\rho T$ and $\beta\rho T$,
- 242 4. $(\alpha \vee \beta)\rho T$ iff $\alpha\rho T$ or $\beta\rho T$,
- 243 5. $(\alpha \wedge \beta)\rho F$ iff $\alpha\rho F$ or $\beta\rho F$,
- 244 6. $(\alpha \vee \beta)\rho F$ iff $\alpha\rho F$ and $\beta\rho F$.

245 **Definition 13.** A formula α is a Dunn semantic consequence of Γ , $\Gamma \vDash_D \alpha$, iff
 246 for all Dunn-interpretations ρ , if $\beta\rho T$ for all $\beta \in \Gamma$, then $\alpha\rho T$.

247 **Definition 14.** Belnap's four-valued semantics for FDE
 248 A four-valued interpretation for FDE is a function v_B from the set of formulas
 249 of FDE to the semantic values $\{T, F, B, N\}$ satisfying the following matrices:

α	$\neg\alpha$	$\alpha \wedge \beta$	T	F	B	N	$\alpha \vee \beta$	T	F	B	N
T	F	T	T	F	B	N	T	T	T	T	T
F	T	F	F	F	F	F	F	T	F	B	N
B	B	B	B	F	B	F	B	T	B	B	T
N	N	N	N	F	F	N	N	T	N	T	N

251

252 **Definition 15.** Let $\mathbf{D} = \{T, B\}$ be the set of designated values of Belnap's
 253 four-valued semantics. A formula α is a four-valued semantic consequence of
 254 Γ , $\Gamma \vDash_B \alpha$, iff for all four-valued interpretations v_B , if $v_B(\beta) \in \mathbf{D}$ for all $\beta \in \Gamma$,
 255 then $v_B(\alpha) \in \mathbf{D}$.

256 The valuation semantics of Definition 3, Dunn's relational semantics of Def-
 257 inition 12, and Belnap's four-valued semantics of Definition 14 intend to repre-
 258 sent four scenarios. Belnap (1977b, p. 11) explains the semantic values T , F ,
 259 N , and B with the notion of a computer 'being told', so, these values mean,
 260 respectively, 'just told true', 'just told false', 'told neither true nor false', and
 261 'told both true and false' (we return to this point in Section 2.2.1 below). Dunn
 262 (1976, p. 156) explains them in terms of subsets of $\{T, F\}$, so a proposition can
 263 be related to $\{T\}$, $\{F\}$, \emptyset , and $\{T, F\}$. In Section 2 above we explained these
 264 four scenarios in terms of availability of evidence.

265 Although both the valuation semantics proposed here and Dunn's relational
 266 semantics are bi-valued, and end up being equivalent, they have an essential
 267 difference: a valuation is a function from the set of formulas to $\{0, 1\}$, while a
 268 Dunn interpretation is a relation between the set of formulas and $\{T, F\}$. In the
 269 latter, a formula can be related simultaneously to both T and F , when it is, in
 270 the Dunn-Belnap reading, both true and false, or not related to T nor F , when
 271 it is neither true nor false. But these three semantics, as expected, validate the
 272 same inferences, i.e. $\Gamma \vDash_{FDE} \alpha$ iff $\Gamma \vDash_D \alpha$ iff $\Gamma \vDash_B \alpha$.

273 **Definition 16.** (*Dunn interpretation induced by an FDE-valuation*)
 274 *Given a FDE-valuation v , we define a Dunn-interpretation ρ_v , based on v , as*
 275 *follows:*

276 $\alpha\rho_v T$ iff $v(\alpha) = 1$

277 $\alpha\neg\rho_v T$ iff $v(\alpha) = 0$

278 $\alpha\rho_v F$ iff $v(\neg\alpha) = 1$

279 $\alpha\neg\rho_v F$ iff $v(\neg\alpha) = 0$

280 **Definition 17.** (*FDE-valuation induced by a Dunn-interpretation*)
 281 *Given a Dunn-interpretation ρ , we define a FDE-valuation v_ρ , based on ρ , as*
 282 *follows:*

283 $v_\rho(\alpha) = 1$ iff $\alpha\rho T$

284 $v_\rho(\alpha) = 0$ iff $\alpha\neg\rho T$

285 $v_\rho(\neg\alpha) = 1$ iff $\alpha\rho F$

286 $v_\rho(\neg\alpha) = 0$ iff $\alpha\neg\rho F$

287 **Lemma 18.** *Given an FDE-valuation v , then ρ_v is a Dunn-interpretation.*

288 *Proof.* We have to prove that ρ_v is a Dunn's relational semantics as in Defini-
 289 tion 12.

290 1. $\neg\alpha\rho_v T$ iff $v(\neg\alpha) = 1$ iff $\alpha\rho_v F$

291 2. $\neg\alpha\rho_v F$ iff $v(\neg\neg\alpha) = 1$ iff $v(\alpha) = 1$ iff $\alpha\rho_v T$

292 3. $(\alpha \wedge \beta)\rho_v T$ iff $v(\alpha \wedge \beta) = 1$ iff $v(\alpha) = 1$ and $v(\beta) = 1$ iff $\alpha\rho_v T$ and $\beta\rho_v T$

293 4. $(\alpha \vee \beta)\rho_v T$ iff $v(\alpha \vee \beta) = 1$ iff $v(\alpha) = 1$ or $v(\beta) = 1$ iff $\alpha\rho_v T$ or $\beta\rho_v T$

294 5. $(\alpha \wedge \beta)\rho_v F$ iff $v(\neg(\alpha \wedge \beta)) = 1$ iff $v(\neg\alpha) = 1$ or $v(\neg\beta) = 1$ iff $\alpha\rho_v F$ or $\beta\rho_v F$

295 6. $(\alpha \vee \beta)\rho_v F$ iff $v(\neg(\alpha \vee \beta)) = 1$ iff $v(\neg\alpha) = 1$ and $v(\neg\beta) = 1$ iff $\alpha\rho_v F$ and
 296 $\beta\rho_v F$

297 □

298 **Lemma 19.** *Given a Dunn-interpretation ρ , then v_ρ is a FDE-valuation.*

299 *Proof.* We have to prove that v_ρ is a FDE-valuation as in Definition 3.

300 1. $v_\rho(\alpha \wedge \beta) = 1$ iff $(\alpha \wedge \beta)\rho T$ iff $\alpha\rho T$ and $\beta\rho T$ iff $v_\rho(\alpha) = 1$ and $v_\rho(\beta) = 1$

301 2. $v_\rho(\alpha \vee \beta) = 1$ iff $(\alpha \vee \beta)\rho T$ iff $\alpha\rho T$ or $\beta\rho T$ iff $v_\rho(\alpha) = 1$ or $v_\rho(\beta) = 1$

302 3. $v_\rho(\neg(\alpha \wedge \beta)) = 1$ iff $(\alpha \wedge \beta)\rho F$ iff $\alpha\rho F$ or $\beta\rho F$ iff $v_\rho(\neg\alpha) = 1$ or $v_\rho(\neg\beta) = 1$

303 4. $v_\rho(\neg(\alpha \vee \beta)) = 1$ iff $(\alpha \vee \beta)\rho F$ iff $\alpha\rho F$ and $\beta\rho F$ iff $v_\rho(\neg\alpha) = 1$ and $v_\rho(\neg\beta) = 1$

304 5. $v_\rho(\alpha) = 1$ iff $(\alpha)\rho T$ iff $\neg\alpha\rho F$ iff $\neg\neg\alpha\rho T$ iff $v_\rho(\neg\neg\alpha) = 1$

305

□

306 **Lemma 20.**

307 *The valuation semantics (Definition 3) and Dunn-interpretation (Definition 12)*
308 *are equivalent, that is, given a valuation semantics v there exists a Dunn-*
309 *interpretation ρ_v such that*

310 $v_\rho(\alpha) = 1$ iff $\alpha\rho T$

311 $v_\rho(\alpha) = 0$ iff $\alpha\neg\rho T$

312 $v_\rho(\neg\alpha) = 1$ iff $\alpha\rho F$

313 $v_\rho(\neg\alpha) = 0$ iff $\alpha\neg\rho F$

314 *for any proposition α ; and vice-versa, given a Dunn-interpretation ρ , there exists*
315 *a valuation v_ρ such that:*

316 $\alpha\rho_v T$ iff $v(\alpha) = 1$

317 $\alpha\neg\rho_v T$ iff $v(\alpha) = 0$

318 $\alpha\rho_v F$ iff $v(\neg\alpha) = 1$

319 $\alpha\neg\rho_v F$ iff $v(\neg\alpha) = 0$

320 *for any proposition α .*

321 *Proof.* Immediate from Lemma 18 and Lemma 19 above. □

322 **Lemma 21.** *The valuation semantics (Definition 3) and Belnap's four-valued*
323 *semantics for FDE (Definition 14) are equivalent.*

324 *Proof.* It follows from Lemma 20 and the well-known fact that Dunn's and
325 Belnap's semantics are equivalent. □

326 **Theorem 22.** *The valuation semantics, the Dunn interpretation and the Bel-*
327 *nap interpretation define equivalent notions of logical consequence: $\Gamma \models_{FDE} \alpha$*
328 *iff $\Gamma \models_B \alpha$ iff $\Gamma \models_D \alpha$.*

329 *Proof.* It follows from Lemma 20 and Lemma 21. □

330 **2.2.1 On paraconsistency, evidence, and information**

331 *FDE* is the well-known and widely studied ‘useful four-valued logic’ proposed by
332 Belnap and Dunn as the underlying logic of an artificial information processor,
333 i.e. a computer, capable of dealing with information received from different
334 sources that are not entirely reliable (cf. Belnap, 1977a,b; Dunn, 1976). The
335 semantic value *Both* is intended to represent the circumstance in which there is
336 conflicting information about α , i.e. both α and $\neg\alpha$ hold, and *None* is intended
337 to represent the circumstance in which there is no information at all about α ,
338 i.e. neither α nor $\neg\alpha$ holds.

339 When Belnap explains these four values, he talks about a computer ‘being
340 told’ that a proposition α is true, or false. The computer should be able to com-
341 pute the values of complex propositions and draw inferences from the received
342 information, but it “can only accept and report information without divesting
343 itself of it” (Belnap, 1977b, p. 9). Of course, contradictory information stored
344 in a database should not be taken as true, as Belnap (1977a, p. 47) remarks
345 that

346 these sentences *have* truth-values independently of what the com-
347 puter has been told; but who can gainsay that the computer cannot
348 *use* the actual truth-value of the sentences in which it is interested?
349 All it can possibly *use* as a basis for inference is what it knows or
350 believes, i.e., what it has been told.

351 The computer, when asked, must provide information based only on what it
352 has been told, otherwise “we would have no way of knowing that its data-base
353 harbored contradictory information” (Belnap, 1977b, p. 9).⁸

354 This notion of ‘a computer being told’ is clearly weaker than truth, since a
355 computer may be told that α is true even if it is not the case. So, Belnap is not
356 really talking about truth *simpliciter*. On the other hand, Dunn (1976, p. 157)
357 seems not to be totally comfortable with the interpretation of *FDE* in terms of
358 the simultaneous truth of α and $\neg\alpha$:

359 Do not get me wrong – I am not claiming that there are sentences
360 which are in fact both true and false. I am merely pointing out
361 that there are plenty of situations where we suppose, assert, believe,
362 etc., contradictory sentences to be true, and we therefore need a
363 semantics which expresses the truth conditions of contradictions in
364 terms of the truth values that the ingredient sentences would have
365 to take for the contradictions to be true.

366 Indeed, we should consider Dunn’s relational semantics as a *façon de parler*,
367 rather than a claim that true contradictions are possible. Obviously, the si-
368 multaneous attribution of the semantic value *True* to a pair of propositions α

⁸Belnap’s approach to the problem is akin to the idea, defended by us in a number of places, that a contradiction α and $\neg\alpha$ can be ‘more informative’ than a single assertion of α , or of $\neg\alpha$, when neither α nor $\neg\alpha$ has been conclusively established. Indeed, in such cases, the contradiction makes it explicit that something is wrong and must be further investigated.

369 and $\neg\alpha$ is not to be understood as an acceptance of dialetheism. It is worth
370 noting that at the time Belnap’s and Dunn’s papers were published, although
371 there were already several paraconsistent formal systems available, the concep-
372 tual discussion about the nature of contradictions accepted by paraconsistent
373 logics was still in its beginnings. It was a ‘lateral issue’ that had not yet been
374 brought to the center of debate.

375 That the four values represented by Belnap-Dunn’s semantics correspond to
376 the four scenarios of availability of evidence the logic *BLE* expresses has been
377 shown in Section 2.2 above. The notions of evidence and information, indeed,
378 are akin to each other, and both are well-suited to a non-dialetheist reading of
379 paraconsistency. Let us take a closer look at these two notions.

380 In Carnielli and Rodrigues (2017, Section 2) the notion of evidence for a
381 proposition α was explained as ‘reasons for believing and/or accepting α ’. Ev-
382 idence, when conclusive, gives support to the truth (or falsity) of α , and thus
383 it has to do with the justification of α (or $\neg\alpha$). The idea behind the recovery
384 operator \circ , introduced in Section 3 below, is that if there is conclusive evidence
385 for the truth, or falsity, of a proposition α , then α is subjected to classical logic.
386 But evidence can be non-conclusive, and so there may be conflicting evidence
387 for a proposition α . Besides being weaker than truth, evidence does not imply
388 belief: there may be evidence for α , an agent may be aware of such evidence but
389 still does not believe in α . If there is non-conclusive evidence for α , it means
390 that there is some degree of justification for α that, however, is not conclusive
391 and might be wrong.⁹

392 Dunn (2008, p. 589) explains a ‘bare-boned’ notion of information as:

393 what is left from knowledge when you subtract, justification, truth,
394 belief, and any other ingredients such as reliability that relate to
395 justification. Information is, as it were, a mere “idle thought.” Oh,
396 one other thing, I want to subtract the thinker. (...) Anyone who
397 has searched for information on the Web does not have to have this
398 concept drummed home. So much of what we find on the Web has
399 no truth or justification, and one would have to be a fool to believe
400 it (...) [Information] is something like a Fregean “thought,” i.e., the
401 “content” of a belief that is equally shared by a doubt, a concern, a
402 wish, etc.

403 Information, so understood, is what is expressed by a proposition, indeed similar
404 to a Fregean thought but without its platonic ingredient. It is objective, does not
405 imply belief, does not need to be true. The difference between this bare-boned
406 notion of information and the notion of non-conclusive evidence is that the latter
407 has an epistemic ingredient that is lacking by the former. So, we can characterize
408 non-conclusive evidence as *bare-boned information plus a justification that might*
409 *be wrong*. Indeed, situations in which we have something that may be or may be
410 not a justification for some proposition α are quite common, and there is nothing

⁹This notion of evidence is in line with the discussion carried out in Achinstein (2010a,b); Kelly (2014).

411 wrong in saying that evidence, conclusive or non-conclusive, is still information:
 412 a proposition α is information, as well as the claim that α has been established
 413 as true. The notion of information is thus more general than evidence. It is not
 414 surprising, therefore, that both *BLE* and *FDE* are suitable to a non-dialetheist
 415 interpretation in terms of evidence and information.

416 3 Extending *FDE* to a logic of evidence and truth

417 *FDE* will now be extended to the logic LET_F , in a similar way to what was
 418 done with *BLE* obtaining LET_J in Carnielli and Rodrigues (2017). Both LET_J
 419 and LET_F are Logics of Formal Inconsistency and Undeterminedness (*LFIs*)
 420 (cf. Carnielli and Rodrigues, 2017; Carnielli, Coniglio, and Rodrigues, 2019;
 421 Marcos, 2005). In *LFIs*,

422 $\alpha, \neg\alpha \not\vdash \beta$, while $\circ\alpha, \alpha, \neg\alpha \vdash \beta$,

423 and in *LFUs*,

424 $\not\vdash \alpha \vee \neg\alpha$, while $\circ\alpha \vdash \alpha \vee \neg\alpha$.¹⁰

425 When $\circ\alpha$ holds, and so excluded middle and explosion are valid, we say that α
 426 is classical. For this reason, in *LFIs*, like the logics LET_J and LET_F , we say
 427 that \circ is a *classicality operator*.

428 Like *BLE*, the logic *FDE*, interpreted from the viewpoint of preservation
 429 of evidence, is not able to express preservation of truth. Indeed, none of the
 430 semantics presented for *FDE* in Section 2.1 can distinguish a context (i) where
 431 there there is non-conclusive evidence for α , so α has not been established as
 432 true, but no evidence for $\neg\alpha$. from another context (ii) where there is conclusive
 433 evidence for α and so α has been established as true, and $\neg\alpha$ does not hold.
 434 In both (i) and (ii), α and $\neg\alpha$ receive respectively the values 1 and 0 by the
 435 valuation semantics (Definition 3), or the values *T* and *F* by the Belnap's four
 436 valued semantics (Definition 14), and so we cannot distinguish between (i) and
 437 (ii). The logic LET_F , on the other hand, is able to distinguish these contexts.

438 **Definition 23.** *The Logic of Evidence and Truth based on FDE (LET_F)*
 439 *Let L_2 be a language with a denumerable set of sentential letters $\{p_1, p_2, p_3, \dots\}$,*
 440 *the set of connectives $\{\circ, \bullet, \neg, \wedge, \vee, \}$ and parentheses. The set of formulas of L_2*
 441 *is obtained recursively in the usual way. The logic LET_F is defined over the*
 442 *language L_2 by adding the following rules to the natural deduction system of*
 443 *FDE (Definition 1):*

¹⁰Definitions of Logics of Formal Inconsistency and Undeterminedness can be found in Carnielli, Coniglio, and Rodrigues (2019) (Defs. 9 and 11). Note that the notion of incompleteness in the interpretation of *FDE* in terms of evidence/information (e.g. Fact 11) is analogous to the notion of undeterminedness in *LFUs*. Actually, in our view, except for the same acronym of *LFIs*, *LFUs* could well be called Logics of Formal Incompleteness. The name *LFU* was established in Marcos (2005) and adopted in Carnielli and Rodrigues (2017) and Carnielli, Coniglio, and Rodrigues (2019).

$$\begin{array}{l}
444 \quad \frac{\circ\alpha \quad \bullet\alpha}{\beta} \textit{Cons} \quad \frac{}{\circ\alpha \vee \bullet\alpha} \textit{Comp} \\
445 \\
446 \quad \frac{\circ\alpha \quad \alpha \quad \neg\alpha}{\beta} \textit{EXP}^\circ \quad \frac{\circ\alpha}{\alpha \vee \neg\alpha} \textit{PEM}^\circ
\end{array}$$

447 A deduction of α from a set of premises Γ in LET_F , $\Gamma \vdash_{LET_F} \alpha$, is defined as
448 follows: there is a derivation with conclusion α and all uncanceled hypotheses in
449 Γ . The definition of a derivation is the usual one for natural deduction systems
450 (see e.g. van Dalen (2008, pp. 35-36)).

451 **Theorem 24.** *The following properties hold for LET_F :*

- 452 1. *Reflexivity: if $\alpha \in \Gamma$, then $\Gamma \vdash_{LET_F} \alpha$;*
- 453 2. *Monotonicity: if $\Gamma \vdash_{LET_F} \beta$, then $\Gamma, \alpha \vdash_{LET_F} \beta$, for any α ;*
- 454 3. *Transitivity (cut): if $\Delta \vdash_{LET_F} \alpha$ and $\Gamma, \alpha \vdash_{LET_F} \beta$, then $\Delta, \Gamma \vdash_{LET_F} \beta$;*
- 455 4. *Compactness: if $\Gamma \vdash_{LET_F} \alpha$, then there is $\Delta \subseteq \Gamma$, Δ finite such that*
456 $\Delta \vdash_{LET_F} \alpha$.

457 *Proof.* Straightforward, from the definition of a deduction of α from premises
458 in Γ in LET_F . \square

Fact 25. *The following rules hold in LET_F :*

$$\frac{\alpha \quad \neg\alpha}{\bullet\alpha} \bullet R1 \quad \frac{}{\alpha \vee \neg\alpha \vee \bullet\alpha} \bullet R2$$

Proof. We prove $\bullet R1$. The proof of $\bullet R2$ is left to the reader.

$$\frac{\frac{}{\circ\alpha \vee \bullet\alpha} \textit{Comp} \quad \frac{[\circ\alpha]^1 \quad \alpha \quad \neg\alpha}{\bullet\alpha} \textit{EXP}^\circ}{\bullet\alpha} \quad \frac{[\bullet\alpha]^1}{1, \vee E}$$

459 \square

460 3.1 On the connectives \circ and \bullet

461 The rules PEM° and EXP° recover classical logic for propositions in the scope
462 of \circ (this claim will be made precise by Fact 31 below). As well as LET_J , LET_F
463 is suitable to an intuitive reading in terms of different contexts concerned with
464 preservation of evidence and preservation of truth. But unlike LET_J , LET_F has
465 a non-classicality operator \bullet , dual to the classicality operator \circ . This duality is
466 made clear by the rules above (Fact 25): $R1$ is the dual of EXP° , and $R2$ is the
467 dual of PEM° ¹¹. While $\circ\alpha$ implies a classical behavior for α , a non-classical

¹¹Actually, different versions of LET_F can be obtained by adding to FDE , besides $Cons$ and $Comp$, the following pair of rules: PEM° and EXP° ; $\bullet R1$ and $\bullet R2$; PEM° and $\bullet R1$; EXP° and $\bullet R2$. Notice that the rules EXP° and $\bullet R2$ are dual, as well as PEM° and $\bullet R1$ (cf. Carnielli et al., 2019).

468 behavior of α implies $\bullet\alpha$. Notice that: (i) $\circ\alpha$ does not imply α , rather, it implies
 469 that one and at most one between α and $\neg\alpha$ holds; (ii) $\bullet\alpha$ does not imply that α
 470 and $\neg\alpha$ hold; indeed, according to *R2*, if both α and $\neg\alpha$ do not hold, $\bullet\alpha$ holds.

471 Strictly speaking, $\circ\alpha$ recovers classical logic for α . The intended interpreta-
 472 tion of $\circ\alpha$ is that there is conclusive evidence for α or $\neg\alpha$, and so the truth-value
 473 of α is conclusively established as true or false. On the other hand, if evidence
 474 for α is non-conclusive, or it is contradictory, or there is no evidence at all, then
 475 $\bullet\alpha$ holds. The rule *Cons* prohibits the circumstance such that *there is* and *there*
 476 *is not* conclusive evidence for α , while *Comp* expresses the fact that either *there*
 477 *is* or *there is not* conclusive evidence for α .

478 Since LET_F distinguishes conclusive from non-conclusive evidence, it is able
 479 to express the following six scenarios:¹²
 480

481 When $\bullet\alpha$ holds, four scenarios of non-conclusive evidence can be expressed:

- 482 1. Only evidence that α is true: α holds, $\neg\alpha$ does not hold.
- 483 2. Only evidence that α is false: $\neg\alpha$ holds, α does not hold.
- 484 3. No evidence at all: both α and $\neg\alpha$ do not hold.
- 485 4. Conflicting evidence: both α and $\neg\alpha$ hold.

486 When $\circ\alpha$ holds, two scenarios can be expressed, tantamount to classical
 487 truth and falsity:

- 488 5. Conclusive evidence for α : α is true ($\circ\alpha \wedge \alpha$ holds).
- 489 6. Conclusive evidence for $\neg\alpha$: $\neg\alpha$ is true ($\circ\alpha \wedge \neg\alpha$ holds).

490 Of course, a scenario with conclusive evidence for both α and $\neg\alpha$ is not allowed,
 491 since it would imply that α is true and false simultaneously. Indeed, if classical
 492 logic holds for α , it cannot be that there is any residual conflicting evidence for
 493 α and $\neg\alpha$.

494 3.2 Valuation semantics for LET_F

495 **Definition 26.** A valuation semantics for LET_F is obtained by adding the
 496 following clauses to the valuation semantics of *FDE* (Definition 3):

- 497 v6. $v(\bullet\alpha) = 1$ iff $v(\circ\alpha) = 0$,
- 498 v7. If $v(\circ\alpha) = 1$, then $v(\alpha) = 1$ if and only if $v(\neg\alpha) = 0$.

¹² In classical logic, ‘ α holds’ means that α is true, while in *FDE*, according to the intended interpretation in terms of evidence, ‘ α holds’ means that there is evidence available for α . In LET_F , the meaning of ‘ α holds’ depends on the context: if the context is classical, it means that α is true. This is precisely the point of the classicality operator \circ . So, two additional scenarios can be expressed, besides the four scenarios of *FDE*.

499 **Definition 27.** We say that a formula α is a *semantical consequence* of Γ ,
500 $\Gamma \vDash_{LET_F} \alpha$ iff for every valuation v , if $v(\beta) = 1$ for all $\beta \in \Gamma$, then $v(\alpha) = 1$.

501 The valuation semantics given above in Definition 26 is sound, complete, and
502 provides a decision procedure for LET_F . From now on, when there is no risk of
503 ambiguity, we will just write \vdash and \vDash in the place of \vdash_{LET_F} and \vDash_{LET_F} .

504 **Theorem 28.** *Soundness and completeness of LET_F w.r.t. the valuation se-*
505 *mantics:* $\Gamma \vDash \alpha$ iff $\Gamma \vdash \alpha$.

506 *Proof.* In order to prove completeness, the proof of Theorem 6 has to be ex-
507 tended to include clauses 6' and 7' below:

508 $v6'$. $\circ\alpha \in \Delta$ iff $\bullet\alpha \notin \Delta$,

509 $v7'$. $\circ\alpha \in \Delta$ implies $\neg\alpha \in \Delta$ iff $\alpha \notin \Delta$.

510 For soundness, it can be shown that rules *Cons*, *Comp*, *EXP*^o, and *PEM*^o are
511 sound with respect to clauses 6 and 7 of Definition 26 above. Details are left to
512 the reader. □

513
514 The quasi-matrix below displays the behavior of the connectives \circ and \bullet in
515 LET_F .

p	0		1			
$\neg p$	0	1	0	1	0	1
$\circ p$	0	1	0	1	0	0
$\bullet p$	1	0	1	0	1	1
<i>valuation</i>	1	2	3	4	5	6

517 The first two lines display the possible values of p and $\neg p$. The connectives \circ
518 and \bullet are primitive and unary, but the semantic values of $\circ p$ and $\bullet p$ depend
519 (non-deterministically) on the semantic values of p and $\neg p$. When $v(p) = 1$ and
520 $v(\neg p) = 0$, $v(p) = 0$ and $v(\neg p) = 1$, the value of $\circ p$ and $\bullet p$ bifurcates into 0 and
521 1. This expresses the fact that $\circ p$ is undetermined in LET_F when $v(p) \neq v(\neg p)$,
522 as explained in page 14 above. In terms of evidence, valuations v_1 and v_6 show,
523 respectively, that no evidence at all, as well as conflicting evidence, implies
524 $v(\bullet p) = 1$ and $v(\circ p) = 0$. But if only one holds among p and $\neg p$ (valuations v_2 to
525 v_5), then $v(\bullet p)$ and $v(\circ p)$ are left undetermined. The rationale of this is that in
526 order to say that p is true, or false, only the information that there is evidence
527 for the truth, or for the falsity, of p is not enough. Something else is needed,
528 namely, the information that such evidence is conclusive.¹³

529 In Example 29 below we illustrate how quasi-matrices work in LET_F .

¹³Note that valuations express evidence available from a purely qualitative point of view. An analogy with analytical chemistry at this point may be illuminating. Qualitative analysis is concerned with whether or not some sample contains a given substance, while quantitative analysis asks how much of a substance is contained in a sample. Analogously, the valuation semantics represents only that there is or there is not positive and negative evidence available for α , while the probabilistic semantics, presented in Section 4 below, intends to express the amount of such evidence.

530 **Example 29.** In LET_F :

531 1. $p \vee \neg p \neq \circ p$

532 2. $\bullet p \neq p \wedge \neg p$

533 3. $\circ p, p, \neg p \vee q \models q$;

534 4. $\circ p, p, \neg(p \wedge q) \models \neg q$;

535 *Proof.* Consider the following quasi-matrix (divided into two parts):

	p	0															
	$\neg p$	0						1									
	q	0				1				0				1			
	$\neg q$	0	1	0	1	0	1	0	1	0	1	0	1	0	1	0	1
536	$\neg p \vee q$	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1
	$\neg(p \wedge q)$	0	1	0	1	1	1	1	1	1	1	1	1	1	1	1	1
	$\circ p$	0	0	0	0	0	1	0	1	0	1	0	1	0	1	0	1
	$\bullet p$	1	1	1	1	1	0	1	0	1	0	1	0	1	0	1	0
	<i>valuation</i>	1	2	3	4	5	6	7	8	9	10	11	12				

	p	1															
	$\neg p$	0						1									
	q	0				1				0				1			
	$\neg q$	0	1	0	1	0	1	0	1	0	1	0	1	0	1	0	1
537	$\neg p \vee q$	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1
	$\neg(p \wedge q)$	0	1	0	1	0	1	0	1	1	1	1	1	1	1	1	1
	$\circ p$	0	1	0	1	0	1	0	1	0	0	0	0	0	0	0	0
	$\bullet p$	1	0	1	0	1	0	1	0	1	1	1	1	1	1	1	1
	<i>valuation</i>	13	14	15	16	17	18	19	20	21	22	23	24				

538 Item 1: since $v_{24}(p) = v_{24}(\neg p) = 1$, $v_{24}(p \vee \neg p) = 1$, but $v_{24}(\circ p) = 0$. Item 2:
539 $v_1(\bullet p) = 1$, but $v_1(p \wedge \neg p) = 0$, since $v_1(p) = v_1(\neg p) = 0$. For items 3 and 4, it
540 is easy to check that there is no valuation v such that the premises receive the
541 value 1 but the conclusion receives 0 in v (compare with items 1 and 2 of Fact
542 7). \square

543 **Remark 30.** The 7th row of the quasi-matrix above is given by clause $v7$ and
544 the 8th by clause $v6$ of Definition 26. A quasi-matrix for LET_F is finite, and
545 similarly to FDE (see Remark 8), it is intuitively clear that the valuation se-
546 mantics provides a decision procedure for LET_F . A detailed algorithm will be
547 presented elsewhere.

548 3.3 Some facts about LET_F

549 Fact 31 below shows how the operator \circ recovers classical logic in LET_F .

550 **Fact 31.** *Recovering classical logic in LET_F*
 551 *Suppose $\circ\lnot^{n_1}\alpha_1, \circ\lnot^{n_2}\alpha_2, \dots, \circ\lnot^{n_m}\alpha_m$ hold, for $n_i \geq 0$ (where, $\lnot^{n_i}, n_i \geq 0$, rep-*
 552 *resents n_i iterations of negations of the formula α_i). Then, for any formula β*
 553 *formed with $\alpha_1, \alpha_2, \dots, \alpha_m$ over $\{\wedge, \vee, \lnot\}$, β behaves classically.*

554 *Proof.*

555 First, we show that for any value of $n_i \geq 0$, $\circ\lnot^{n_i}\alpha_i \vdash \alpha_i \vee \lnot\alpha_i$ and $\circ\lnot^{n_i}\alpha_i, \alpha_i \wedge$
 556 $\lnot\alpha_i \vdash \gamma$, for any γ – i.e. excluded middle and explosion hold for α_i .

557 Suppose $\circ\lnot^{n_i}\alpha_i$ holds. So, $\lnot^{n_i}\alpha_i \vee \lnot\lnot^{n_i}\alpha_i$ and $\lnot^{n_i}\alpha_i \wedge \lnot\lnot^{n_i}\alpha_i \vdash \gamma$ hold. If n_i
 558 is even, $\lnot^{n_i}\alpha_i \dashv\vdash \alpha_i$, and if n_i is odd, $\lnot^{n_i}\alpha_i \dashv\vdash \lnot\alpha_i$. So, it is easily proved
 559 that $\lnot^{n_i}\alpha_i \vee \lnot\lnot^{n_i}\alpha_i \vdash \alpha_i \vee \lnot\alpha_i$. Since we have that $\circ\lnot^{n_i}\alpha_i \vdash \lnot^{n_i}\alpha_i \vee \lnot\lnot^{n_i}\alpha_i$, by
 560 transitivity, we get $\circ\lnot^{n_i}\alpha_i \vdash \alpha_i \vee \lnot\alpha_i$. In order to recover explosion, it can be
 561 easily proved that $\alpha_i \wedge \lnot\alpha_i \vdash \lnot^{n_i}\alpha_i \wedge \lnot\lnot^{n_i}\alpha_i$. Since we have that $\circ\lnot^{n_i}\alpha_i, \lnot^{n_i}\alpha_i \wedge$
 562 $\lnot\lnot^{n_i}\alpha_i \vdash \gamma$, by transitivity, we get $\circ\lnot^{n_i}\alpha_i, \alpha_i \wedge \lnot\alpha_i \vdash \gamma$.

563 Remember that full classical logic can be obtained by adding explosion and
 564 excluded middle to the introduction and elimination rules of \wedge and \vee , $\alpha_1 \rightarrow \alpha_2$
 565 being defined as $\lnot\alpha_1 \vee \alpha_2$. Now, in order to prove the result, it is enough
 566 to show that for any formula β formed with $\alpha_1, \alpha_2, \dots, \alpha_m$ over $\{\wedge, \vee, \lnot\}$, if
 567 $\circ\lnot^{n_1}\alpha_1, \circ\lnot^{n_2}\alpha_2, \dots, \circ\lnot^{n_m}\alpha_m$ hold, then $\vdash \beta \vee \lnot\beta$ and $\beta, \lnot\beta \vdash \gamma$ hold.

568 Let $\Gamma = \{\circ\lnot^{n_1}\alpha_1, \circ\lnot^{n_2}\alpha_2, \dots, \circ\lnot^{n_m}\alpha_m\}$.

569 If $\beta = \alpha_i$, it has been proved above. The remaining cases are proved by induction
 570 on the complexity of β .

571 Case 1. $\beta = \lnot\delta$. I.H. $\Gamma, \delta, \lnot\delta \vdash \gamma$ and $\Gamma \vdash \delta \vee \lnot\delta$. It can be easily proved that
 572 $\Gamma, \lnot\delta, \lnot\lnot\delta \vdash \gamma$ and $\Gamma \vdash \lnot\delta \vee \lnot\lnot\delta$.

573 Case 2. $\beta = \delta_1 \wedge \delta_2$. I.H. $\Gamma, \delta_1, \lnot\delta_1 \vdash \gamma$ and $\Gamma \vdash \delta_1 \vee \lnot\delta_1$, *mutatis mutandis* for
 574 δ_2 . It can be proved that $\Gamma, \delta_1 \wedge \delta_2, \lnot(\delta_1 \wedge \delta_2) \vdash \gamma$ and $\Gamma \vdash (\delta_1 \wedge \delta_2) \vee \lnot(\delta_1 \wedge \delta_2)$
 575 The remaining cases are left to the reader. \square

576 We have seen in Fact 9 that for an implication $\alpha \rightarrow \beta$ defined in FDE
 577 as $\lnot\alpha \vee \beta$, modus ponens and the deduction theorem do not hold. Both are
 578 recovered for the defined implication in LET_F for classical propositions.

579 **Fact 32.**

580 1. In LET_F , for classical propositions, modus ponens holds: $\circ\alpha, \alpha, \lnot\alpha \vee \beta \vdash \beta$.

Proof.

$$\frac{\lnot\alpha \vee \beta \quad \frac{\circ\alpha \quad \alpha \quad [-\alpha]^1}{\beta} \text{EXP}^\circ \quad [\beta]^1}{\beta} \text{1,}\vee\text{E}$$

581 \square

582 2. In LET_F , the following form of the deduction theorem holds: $\circ\alpha, \alpha \vdash \beta$
 583 implies $\circ\alpha \vdash \lnot\alpha \vee \beta$.

Proof.

$$\frac{\frac{\frac{\circ\alpha}{\alpha \vee \neg\alpha} \text{ PEM}^\circ \quad \frac{\frac{\frac{\circ\alpha, [\alpha]^1}{\vdots} \beta}{\neg\alpha \vee \beta} \vee I \quad \frac{[-\alpha]^1}{\neg\alpha \vee \beta} \vee I}{\neg\alpha \vee \beta} \text{ 1, \vee E}}{\neg\alpha \vee \beta}}{\neg\alpha \vee \beta}}$$

584

□

585 **Definition 33.** *Supplementing and complementing negations (Carnielli et al.,*
 586 *2007, pp. 12ff)*

- 587 1. We say that a unary connective $*$ in a logic \mathcal{L} is a supplementing nega-
 588 tion if: (i) for some formula α , $*\alpha$ is not a bottom particle, and (ii) for
 589 any Γ, α and β : $\Gamma, \alpha, *\alpha \vdash_{\mathcal{L}} \beta$.
- 590 2. We say that a unary connective $*$ in a logic \mathcal{L} is a complementing nega-
 591 tion if: (i) for some formula α , $*\alpha$ is not a top particle;
 592 (ii) for any Γ, α and β : $\Gamma, \alpha \vdash_{\mathcal{L}} \beta$ and $\Gamma, *\alpha \vdash_{\mathcal{L}} \beta$ implies $\Gamma \vdash_{\mathcal{L}} \beta$.

593 If $*$ is a complementing negation, for any α , at least one between α and $*\alpha$ hold,
 594 and if $*$ is a supplementing negation, it cannot be that both α and $*\alpha$ hold.
 595 Each one expresses one half of classical negation, the former excluded middle,
 596 the latter explosion. If a logic \mathcal{L} has a (primitive or defined) negation connective
 597 that is both supplementing and complementing, then \mathcal{L} has a classical negation.
 598 A complementing negation and a supplementing negation can be defined in
 599 LET_F .

600 **Definition 34.** *The following unary connectives can be defined in LET_F :*

- 601 1. The connective truth: $\oplus\alpha \stackrel{\text{def}}{=} \circ\alpha \wedge \alpha$;
- 602 2. The connective falsity: $\sim\alpha \stackrel{\text{def}}{=} \circ\alpha \wedge \neg\alpha$;
- 603 3. The connective falsity-excluding: $\ominus\alpha \stackrel{\text{def}}{=} \bullet\alpha \vee \alpha$;
- 604 4. The connective truth-excluding: $\approx\alpha \stackrel{\text{def}}{=} \bullet\alpha \vee \neg\alpha$.


605 *The tables are the following:*

α	0		1	
$\neg\alpha$	0	1	0	1
$\circ\alpha$	0	0	1	0
$\bullet\alpha$	1	1	0	0
$\oplus\alpha$	0	0	0	0
$\sim\alpha$	0	0	1	0
$\ominus\alpha$	1	1	1	1
$\approx\alpha$	1	1	1	0

606

607 These connectives have been named for the following reasons. According to the
608 proposed interpretation: (1) $\circ\alpha \wedge \alpha$ means that there is conclusive evidence for
609 α , and so α is true ($\oplus\alpha$); (2) $\circ\alpha \wedge \neg\alpha$ means that there is conclusive evidence
610 for the falsity of α , and so α is false ($\sim\alpha$); (3) $\bullet\alpha \vee \alpha$ means that there is no
611 conclusive evidence for α , or α holds, and so it excludes the falsity of α ($\ominus\alpha$); (4)
612 $\bullet\alpha \vee \neg\alpha$ means that there is no conclusive evidence for α , or $\neg\alpha$ holds, and so it
613 excludes the truth of α ($\approx\alpha$). It is also clear from the table above and Definition
614 33 that $\sim\alpha$ is a supplementing negation (if $v(\alpha) = 1$, $v(\sim\alpha) = 0$, they cannot
615 be both 1), while $\approx\alpha$ is a complementing negation (if $v(\alpha) = 0$, $v(\approx\alpha) = 1$, they
616 cannot be both 0).¹⁴ We conjecture that no classical negation can be defined in
617 LET_F .¹⁵

618 These four connectives enjoy some interesting logical relations w.r.t. each
619 other that can be displayed by a square of oppositions:



620

621 $\sim\alpha$ and $\ominus\alpha$ are contrary propositions (i.e., they can both be false, but they
622 cannot both be simultaneously true); $\approx\alpha$ and $\ominus\alpha$ are subcontrary propositions
623 (i.e., they can both be true, but they cannot both be simultaneously false); $\oplus\alpha$
624 (resp. $\sim\alpha$) is the dual of $\ominus\alpha$ (resp. $\approx\alpha$); $\oplus\alpha$ (resp. $\sim\alpha$) is the contradictory
625 of $\approx\alpha$ (resp. \ominus). Notice that in LET_F , \circ is the dual of \bullet , and \neg is the dual
626 of itself (on duality between non-deterministic connectives in Logics of Formal
627 Inconsistency and Undeterminedness, see Carnielli et al. (2019)).

628 **Fact 35.**

629 1. $\circ\alpha \wedge \alpha \wedge \neg\alpha$, $\circ\alpha \wedge \bullet\alpha$, $\oplus\alpha \wedge \sim\alpha$, $\oplus\alpha \wedge \approx\alpha$, and $\ominus\alpha \wedge \sim\alpha$ are bottom particles
630 in LET_F .

¹⁴Although \sim is explosive, it is not a classical negation, since $\alpha \vee \sim\alpha$ does not hold, which is shown by the valuation $v(\alpha) = v(\neg\alpha) = v(\circ\alpha) = 0$, and although $\alpha \vee \approx\alpha$ holds, \approx is also not a classical negation, since $\alpha, \approx\alpha \vdash \beta$ does not hold, which is shown by the valuation $v(\alpha) = v(\neg\alpha) = v(\bullet\alpha) = 1$.

¹⁵One possibility for proving that classical negation is not definable in LET_F is to adapt the methods of Lahav, Marcos, and Zohar (2016), although they are devoted to non-classical negations from a modal viewpoint. We have been unable however, to find a convincing argument in this direction.

631 2. \sim is a supplementing negation in LET_F .

632 3. \approx is a complementing negation in LET_F .

633 *Proof.* In a few steps from the rules *Cons*, *EXP^o* and $\bullet R2$. □

634 **Theorem 36.** *The following propositions are theorems of LET_F :*

635 1. $\circ\alpha \vee \bullet\alpha$

636 2. $\alpha \vee \neg\alpha \vee \bullet\alpha$

637 3. $(\bullet\alpha \wedge \alpha) \vee (\bullet\alpha \wedge \neg\alpha) \vee \bullet\alpha \vee (\bullet\alpha \wedge \alpha \wedge \neg\alpha) \vee (\circ\alpha \wedge \alpha) \vee (\circ\alpha \wedge \neg\alpha)$

638 4. $\alpha \vee \neg\alpha \vee \Theta\alpha$

639 5. $\alpha \vee \neg\alpha \vee \approx\alpha$

640 6. $\Theta\alpha \vee \sim\alpha \vee \bullet\alpha$

641 *Proof.* Items 1 and 2 follow from the rules *Comp* and *R2*. To prove 3, from
642 $\circ\alpha \vdash \alpha \vee \neg\alpha$, we obtain $\circ\alpha \vdash (\circ\alpha \wedge \alpha) \vee (\circ\alpha \wedge \neg\alpha)$, and so $\circ\alpha \vdash (\bullet\alpha \wedge \alpha) \vee$
643 $(\bullet\alpha \wedge \neg\alpha) \vee \bullet\alpha \vee (\bullet\alpha \wedge \alpha \wedge \neg\alpha) \vee (\circ\alpha \wedge \alpha) \vee (\circ\alpha \wedge \neg\alpha)$. On the other hand,
644 $\bullet\alpha \vdash (\bullet\alpha \wedge \alpha) \vee (\bullet\alpha \wedge \neg\alpha) \vee \bullet\alpha \vee (\bullet\alpha \wedge \alpha \wedge \neg\alpha) \vee (\circ\alpha \wedge \alpha) \vee (\circ\alpha \wedge \neg\alpha)$ holds. Now,
645 use 1 and $\vee E$. The proofs of 4, 5, and 6 are left to the reader. Notice that item
646 3 corresponds to the six scenarios presented in Section 3.1. □

647 4 Probabilistic semantics for LET_F

648 We now present a probabilistic semantics for LET_F and *FDE*.

649 **Definition 37.** *Given a logic \mathcal{L} , with a derivability relation \vdash and a language*
650 *L , a probability distribution for \mathcal{L} is a real-valued function $P : L \rightarrow \mathbb{R}$ satisfying*
651 *the following conditions:*

652 1. *Non-negativity:* $0 \leq P(\alpha) \leq 1$ for all $\alpha \in L$;

653 2. *Tautologicity:* If $\vdash \alpha$, then $P(\alpha) = 1$;

654 3. *Anti-Tautologicity:* If $\alpha = \perp$, then $P(\alpha) = 0$;

655 4. *Comparison:* If $\alpha \vdash \beta$, then $P(\alpha) \leq P(\beta)$;

656 5. *Finite additivity:* $P(\alpha \vee \beta) = P(\alpha) + P(\beta) - P(\alpha \wedge \beta)$.

657 The clauses above can be regarded as meta-axioms that define probability func-
658 tions for an appropriate logic \mathcal{L} just by taking \vdash as the derivability relation of
659 \mathcal{L} , and so the notion of probability can be regarded as logic-dependent. These
660 clauses define probability functions for both *FDE* and LET_F just by employing

661 respectively \vdash_{FDE} and \vdash_{LET_F} .¹⁶ From now on, we will concentrate on LET_F ,
 662 but it should be clear that the meta-axioms 1, 4 and 5 above define probability
 663 distributions for FDE as well.

664 **Definition 38.** *LET_F-probability distribution*
 665 *Let $\Sigma = \{\alpha_1, \dots, \alpha_n, \dots\}$ be a (finite or infinite) collection of propositions in the*
 666 *language L_2 of LET_F . A LET_F -probability distribution over Σ is an assign-*
 667 *ment of probability values P to the elements of Σ that can be extended to a full*
 668 *probability function $P : L_2 \rightarrow \mathbb{R}$ according to Definition 37.*

669 It is a common view that the classical truth-values *true* (1) and *false* (0)
 670 can be identified with the endpoints of probabilities in the unit interval $[0, 1]$.
 671 On the other hand, interpretations $v : L \rightarrow \{0, 1\}$ of a formal language L can
 672 be regarded as degenerate probability functions $P : L \rightarrow [0, 1]$. The class of
 673 logics that make possible such an identification can be seen as a special case of
 674 probability logic. The standard view, however, is rather the opposite: it claims
 675 that probability logic presupposes, and so it depends on, classical logic.¹⁷ But
 676 the connection between logic and probability theory is far from being restricted
 677 to classical logic. The fact that probability distributions can be defined based on
 678 a non-classical consequence relation, in our view, makes clear that the relation
 679 between logic and probability goes beyond the realm of classical logic.

680 4.1 Conditional probability

681 The notion of conditional probability of α given β is defined as usual, for $P(\beta) \neq 0$:

$$682 \quad P(\alpha/\beta) = \frac{P(\alpha \wedge \beta)}{P(\beta)}$$

683 In terms of evidence, a statement $P(\alpha/\beta)$ is to be read as a measure of how
 684 much the evidence available for β affects the evidence for α .

685
 686 Some useful theorems of conditional probability of LET_F -distributions are
 687 the following, with the caveat that $P(\beta) \neq 0$ in all cases where $P(\alpha/\beta)$ is men-
 688 tioned:

¹⁶Probability functions have been defined in this way for classical logic, for intuitionistic logic without implication in Weatherson (2003), and for the paraconsistent logics Ci and Cie in Bueno-Soler and Carnielli (2016, 2017).

¹⁷In a recent article, Demey et al. (2013) claim that “probability theory presupposes and extends classical logic”, and leave aside all the attempts to combine probability theory with non-classical logics. These attempts, however, not only do exist, but have also been successful in combining probability theory with non-classical approaches to logical consequence. We think Demey *et al.* are mistaken, not only because they ignore non-classical approaches to probability logic, but also because they underestimate the view according to which classical and some non-classical logics can be seen as special cases of probability logic. It is worth noting that attempts to put together probability theory and non-classical logics can be traced back to Łukasiewicz (1913) and Tarski (1935).

689 **Theorem 39.**

690 *The following properties hold when the probabilities in the denominators are*
 691 *different from 0.*

- 692 1. $P(\alpha_1 \wedge \dots \wedge \alpha_n) = P(\alpha_1/\alpha_2 \wedge \dots \wedge \alpha_n) \dots P(\alpha_{n-1}/\alpha_n)P(\alpha_n)$ (Chain Rule).
- 693 2. $P(\alpha/\beta \wedge \gamma) = \frac{P(\alpha/\gamma) \cdot P(\beta/\alpha \wedge \gamma)}{P(\beta/\gamma)}$.
- 694 3. $P(\alpha \wedge \beta/\gamma) = P(\alpha/\gamma) \cdot P(\beta/\alpha \wedge \gamma) = P(\beta/\gamma) \cdot P(\alpha/\beta \wedge \gamma)$.
- 695 4. $P(\alpha \vee \beta/\gamma) = P(\alpha/\gamma) + P(\beta/\gamma) - P(\alpha \wedge \beta/\gamma)$.
- 696 5. $P(\alpha \vee \beta/\gamma) = P(\alpha/\gamma) + P(\beta/\gamma)$ if α and β are logically incompatible, i.e.,
 697 $\alpha \wedge \beta$ act as $a \perp$ (see Section 4.2).
- 698 6. $P(\alpha/\beta) + P(\neg\alpha/\beta) - P(\bullet\alpha/\beta) \leq P(\alpha \vee \neg\alpha/\beta)$.
- 699 7. If $P(\circ\alpha) = 1$, or equivalently $P(\bullet\alpha) = 0$, then $P(\alpha \vee \neg\alpha) = 1$ and $P(\alpha \wedge$
 700 $\neg\alpha) = 0$.
- 701 8. $P(\alpha/\beta) + P(\neg\alpha/\beta) = 1$, if $P(\circ\alpha) = 1$.
- 702 9. $P(\beta/\circ\beta) + P(\neg\beta/\circ\beta) = 1$.

703 *Proof.*

704 Items 1 to 4 are quite elementary properties coming from the general
 705 definition of conditional probability: $P(\alpha/\beta) = \frac{P(\alpha \wedge \beta)}{P(\beta)}$, which gives the
 706 alternative product rule $P(\alpha \wedge \beta) = P(\alpha/\beta) \cdot P(\beta)$. The chain rule (item
 707 1) is derived by successive applications of product rule. Items 2 to 4 are
 708 easy consequences of the definition of conditional probability and clause 5
 709 of Definition 37.

710 Item 5: since $\alpha \wedge \beta$ is a bottom particle in this case, $P(\alpha \wedge \beta) = 0$, and the
 711 result follows from 4.

712 Item 6 is a consequence of Fact 25 ($\alpha \wedge \neg\alpha \vdash \bullet\alpha$), Comparison and ele-
 713 mentary inequalities, plus the definition of conditional probability.

714 Item 7: Easy consequence of Definition 22, R1 (Fact 24) and Comparison
 715 (Definition 35).

716 Item 8: If $P(\circ\alpha) = 1$, then by 9 $P(\alpha \vee \neg\alpha) = 1$, and by Lemma 40 (below)
 717 we have $P((\alpha \vee \neg\alpha) \wedge \beta) = P(\beta) = P((\alpha \wedge \beta) \vee (\neg\alpha \wedge \beta)) = P(\alpha \wedge \beta) +$
 718 $P(\neg\alpha \wedge \beta) - P(\alpha \wedge \neg\alpha \wedge \beta)$. Since $P(\alpha \wedge \neg\alpha \wedge \beta) = 0$ ($P(\circ\alpha) = 1$ implies
 719 $P(\alpha \wedge \neg\alpha) = 0$), we obtain $P(\beta) = P(\alpha \wedge \beta) + P(\neg\alpha \wedge \beta)$. Dividing both
 720 sides by $P(\beta)$ obtains the result, in view of the definition of conditional
 721 probability.

722 Item 9: In LET_F , $\vdash \circ\beta \leftrightarrow \circ\beta \wedge (\beta \vee \neg\beta) \leftrightarrow (\circ\beta \wedge \beta) \vee (\circ\beta \wedge \neg\beta)$ (proof left to
723 the reader). Thus $P(\circ\beta) = P((\circ\beta \wedge \beta) \vee (\circ\beta \wedge \neg\beta)) = P(\circ\beta \wedge \beta) + P(\circ\beta \wedge \neg\beta)$
724 by Finite Additivity, since $\circ\beta \wedge \beta \wedge \neg\beta$ is a bottom particle (Fact 33).
725 Dividing both sides by $P(\circ\beta)$ yields the result.

726 □

727 4.2 Independence and incompatibility

728 Intuitively, two propositions are independent if the fact that one holds does
729 not have any effect on whether or not the other holds, and vice-versa. Two
730 propositions α and β , are said to be independent w.r.t. a distribution P if
731 $P(\alpha \wedge \beta) = P(\alpha) \cdot P(\beta)$. Two propositions can be independent relative to
732 one probability distribution and dependent relative to another. Alternatively,
733 independence can be defined as follows: α is independent of β if $P(\alpha/\beta) = P(\alpha)$
734 (or equivalently, $P(\beta/\alpha) = P(\beta)$).¹⁸ Classically, α and $\neg\alpha$ are never independent
735 (unless one of them has probability zero). In view of item 4 of Theorem 42
736 below, $P(\alpha \wedge \neg\alpha) \leq P(\bullet\alpha)$, hence when $P(\alpha) \cdot P(\neg\alpha) > P(\bullet\alpha)$, α and $\neg\alpha$ are
737 not independent. In this way, $P(\bullet\alpha)$ can be regarded as a bound on the ‘degree
738 of independence’ between α and $\neg\alpha$.

739 Intuitively, two propositions α and β are logically incompatible if α cannot
740 hold when β holds, and vice-versa. Two propositions α and β , are said to
741 be logically incompatible if $\alpha, \beta \vdash \gamma$, for any γ , or equivalently, if $\alpha \wedge \beta$ is a
742 bottom particle. Logically incompatible propositions α and β with non-zero
743 probabilities are always *dependent* since $0 = P(\alpha \wedge \beta) \neq P(\alpha) \cdot P(\beta)$. Again, for
744 non-zero probabilities, classically α and $\neg\alpha$ are incompatible, and so dependent.
745 In LET_F , however, they are neither necessarily incompatible nor necessarily
746 dependent, when $P(\circ\alpha) < 1$. We saw in Fact 35 item 1 that $\alpha \wedge \neg\alpha \wedge \circ\alpha$ as well
747 as $\circ\alpha \wedge \bullet\alpha$ defines a bottom particle in LET_F . From clause 3 of Definition 37, it
748 follows that for any probability distribution P , $P(\alpha \wedge \neg\alpha \wedge \circ\alpha) = 0$ and $P(\circ\alpha \wedge$
749 $\bullet\alpha) = 0$. So, in LET_F α and $\sim\alpha$ are always logically incompatible and hence
750 dependent, while α and $\neg\alpha$ can be independent.

751 An interesting property concerning the behavior of probability measures in
752 LET_F , related to independence in ‘extreme cases’, occurs when $P(\alpha) = 1$. In
753 such cases α is independent from the probability measure of any other distinct
754 proposition β . This kind of property contributes to the dynamics of evidence, in
755 the sense of the interpretation of preservation of conclusive and non-conclusive
756 evidence in LET_F , in such a way that the increasing of conclusive evidence
757 tends to truth.

758 **Lemma 40.** *Independence of propositions with maximal probability*
759 *If $P(\alpha) = 1$ then $P(\alpha \wedge \beta) = P(\alpha) \cdot P(\beta)$, for $\beta \neq \alpha$*

¹⁸Although mathematically equivalent to the former, this characterization of independence by means of conditional probability is debatable, as shown in Fitelson and Hájek (2017), where it is argued that the more general Popperian theory of conditional probability should be adopted, leading to a revision of conventional insights about probabilistic independence. The traditional notions are employed here for mathematical convenience.

760 *Proof.* If $P(\alpha) = 1$ then $P(\alpha \vee \beta) = 1$ from Comparison, since $\alpha \vdash \alpha \vee \beta$. By
 761 Finite Additivity $1 = P(\alpha \vee \beta) = P(\alpha) + P(\beta) - P(\alpha \wedge \beta)$. As $P(\alpha) = 1$, it follows
 762 that $P(\alpha \wedge \beta) = P(\beta)$. \square

763 The restriction $\alpha \neq \beta$ in the above lemma intends to avoid the problematic cases
 764 of ‘self-independence’ of extreme events. As mentioned before, two events α
 765 and β are considered to be independent if $P(\alpha \wedge \beta) = P(\alpha) \cdot P(\beta)$, for some
 766 probability distribution P . This leads to a puzzling situation concerning events
 767 α such that $P(\alpha) = 0$ or $P(\alpha) = 1$. In such cases, $P(\alpha) = P(\alpha \wedge \alpha) = P(\alpha) \cdot P(\alpha)$
 768 in both cases. In this way, extreme probabilities can be regarded as independent
 769 of themselves, an uncomfortable situation, as recognized in Fitelson and Hájek
 770 (2017).

771 Lemma 40 leads immediately to the independence of consistent and incon-
 772 sistent propositions in extreme cases:

- 773 1. If $P(\circ\alpha) = 1$ then $P(\circ\alpha \wedge \beta) = P(\beta)$, for $\beta \neq \circ\alpha$
- 774 2. If $P(\bullet\alpha) = 1$ then $P(\bullet\alpha \wedge \beta) = P(\beta)$, for $\beta \neq \bullet\alpha$
- 775 3. If $P(\beta) = 1$ then $P(\circ\alpha \wedge \beta) = P(\circ\alpha)$, for $\beta \neq \circ\alpha$
- 776 4. If $P(\beta) = 1$ then $P(\bullet\alpha \wedge \beta) = P(\bullet\alpha)$, for $\beta \neq \bullet\alpha$

777 Evidence can be increasing or decreasing in an historical series, leading to a dy-
 778 namic of evidence. This can be expressed in mathematical terms by elementary
 779 series. Let $\lim_{i \rightarrow \infty} P_i(\alpha) = \lambda$ mean that the sequence of values $P_1(\alpha), P_2(\alpha), \dots, P_i(\alpha) \dots$
 780 is strictly monotonous and converges to $\lambda \in [0, 1]$.

781 **Lemma 41.** *The dynamics of evidence*

- 782 1. If $\lim_{i \rightarrow \infty} P_i(\circ\alpha) = 1$ or $\lim_{i \rightarrow \infty} P_i(\bullet\alpha) = 0$, then $\lim_{i \rightarrow \infty} P_i(\alpha \vee \neg\alpha) = 1$
 783 and $\lim_{i \rightarrow \infty} P_i(\alpha \wedge \neg\alpha) = 0$.
- 784 2. If $\lim_{i \rightarrow \infty} P_i(\circ\alpha) = 1$ or $\lim_{i \rightarrow \infty} P_i(\bullet\alpha) = 0$, then $\lim_{i \rightarrow \infty} (P_i(\alpha) + P_i(\neg\alpha)) =$
 785 1.

786 *Proof.* Suppose $\lim_{i \rightarrow \infty} P_i(\circ\alpha) = 1$; by *PEM*^o and Comparison, $P_i(\circ\alpha) \leq P_i(\alpha \vee$
 787 $\neg\alpha) \leq 1$. By the Squeeze Theorem of elementary calculus for series (aka the
 788 Sandwich Theorem) $\lim_{i \rightarrow \infty} P_i(\alpha \vee \neg\alpha) = 1$. All other limits are proved in similar
 789 ways. \square

790 The meaning of Lemma 41 is precisely that the values of $P_i(\circ\alpha)$ can be in-
 791 terpreted as degrees of classicality, in the sense that greater values of $P_i(\circ\alpha)$
 792 indicate that the situation is approaching classicality and, conversely, the val-
 793 ues of $P_i(\bullet\alpha)$ can be interpreted as degrees of anticlassicality, in the sense that
 794 smaller values of $P_i(\bullet\alpha)$ indicate that the situation is approaching classicality.

795
 796 Some useful (though almost all immediate) properties of *LET_F*-distributions
 797 are the following:

798 **Theorem 42.**

- 799 1. If $\alpha \dashv\vdash \beta$, then $P(\alpha) = P(\beta)$.
- 800 2. $P(\alpha \vee \beta) = P(\alpha) + P(\beta)$, if α and β are logically incompatible.
- 801 3. $P(\alpha \vee \beta \vee \gamma) = P(\alpha) + P(\beta) + P(\gamma) - P(\alpha \wedge \beta) - P(\alpha \wedge \gamma) - P(\beta \wedge \gamma) + P(\alpha \wedge \beta \wedge \gamma)$.
- 802 4. $P(\alpha \wedge \neg\alpha) \leq P(\bullet\alpha)$.
- 803 5. $P(\circ\alpha) \leq P(\alpha \vee \neg\alpha)$.
- 804 6. $P(\circ\alpha) = 1 - P(\bullet\alpha)$.
- 805 7. $P(\oplus\alpha \wedge \neg\alpha) = 0$, $P(\sim\alpha \wedge \alpha) = 0$.
- 806 8. $P(\circ\alpha \vee \bullet\alpha) = 1$, $P(\alpha \vee \neg\alpha \vee \bullet\alpha) = 1$
- 807 9. $P(\circ\alpha \vee (\alpha \wedge \neg\alpha)) \leq P(\alpha \vee \neg\alpha)$
- 808 10. $1 + P((\alpha \vee \neg\alpha) \wedge \bullet\alpha) = P(\alpha \vee \neg\alpha) + P(\bullet\alpha)$
- 809 11. $1 + P((\alpha \wedge \neg\alpha) \vee \circ\alpha) = P(\alpha \wedge \neg\alpha) + P(\circ\alpha)$
- 810 12. If $P(\circ\alpha) = 1$ (or equivalently $P(\bullet\alpha) = 0$), then $P(\neg\alpha) = 1 - P(\alpha)$
- 811 13. If $P(\circ\alpha) = 1$ (or equivalently $P(\bullet\alpha) = 0$), then $P(\alpha \vee \neg\alpha) = 1$ and $P(\alpha \wedge$
812 $\neg\alpha) = 0$.

813 *Proof.* Routine, from the axioms of probability and the derivability relation of
814 LET_F . We just sketch the proof of items 12 and 13. For 12, suppose $P(\circ\alpha) = 1$;
815 by items 4 and 5 above, PEM° , and Comparison, $1 = P(\circ\alpha) \leq P(\alpha \vee \neg\alpha)$, and
816 $P(\alpha \wedge \neg\alpha) \leq P(\bullet\alpha) = 0$, hence by Finite Additivity $P(\alpha \vee \neg\alpha) + P(\alpha \wedge \neg\alpha) =$
817 $1 + 0 = P(\alpha) + P(\neg\alpha)$. Hence $P(\neg\alpha) = 1 - P(\alpha)$. For 13 a similar reasoning as
818 of 12 is obtained. \square

819 Items 1 and 2 are usual results in probabilistic logic, and 3 is a particular
820 case of the Inclusion-Exclusion property for finite probability, easily adapted
821 for propositions, that hold for arbitrary finite disjunctions (see Grinstead and
822 Snell (1997)). Items 4 and 5 establish constraints on the values of $P(\circ\alpha)$,
823 $P(\alpha)$ and $P(\neg\alpha)$. Item 7 concerns bottom particles, and 8, theorems of LET_F
824 essential for proving total probability theorems (Section 4.3 below). Items 12
825 and 13 show the classical behavior of probabilities when $P(\circ\alpha) = 1$.

826 **4.3 Total probability theorems for LET_F**

In the classical approach to probability, total probability theorems compute the probability of an event β in a sample space partitioned into exclusive and exhaustive events. Typically, for a partition in two pieces, a total probability theorem that reflects excluded middle assumes the following form:

$$P(\beta) = P(\beta \wedge \alpha) + P(\beta \wedge \neg\alpha).$$

827 Here, however, we are not really talking about sample spaces, about events
 828 themselves, but rather about the information related to such events, that we call
 829 an *information space*. In the standard approach to probability theory, we start
 830 from a group of events, say, the two outcomes of tossing a coin, and attribute
 831 probabilities to these events, whose sum is always equal to 1. Let α express that
 832 the toss of a coin comes up heads. The sample space is thus divided into two
 833 parts, α and $\neg\alpha$, corresponding respectively to heads and tails (not heads). If
 834 the coin behaves as expected, their probability are the same.

835 On the other hand, we consider here a language that is able to express in-
 836 formation about some event – for example, the result of a referendum – that
 837 comes from different sources and may be unreliable. Such information is consti-
 838 tuted by evidence for ‘yes’ and for ‘no’ that can be non-conclusive, incomplete,
 839 contradictory, more reliable or less reliable, and perhaps even conclusive. Let
 840 α express the result ‘yes’, and $\neg\alpha$ the result ‘no’. In this case, the propositions
 841 we are concerned with are α , $\neg\alpha$, $\circ\alpha$, $\bullet\alpha$, as well as other propositions of the
 842 language of LET_F formed from them, for example, $\bullet\alpha \vee \alpha$, $\alpha \wedge \neg\alpha$, $\circ\alpha \wedge \alpha$, etc.
 843 A LET_F -probability distribution attributes values to these propositions. The
 844 information space is thus constituted by such propositions and the measures
 845 of probabilities attributed to them by a LET_F -probability distribution P . Note
 846 that, contrary to the classical case, $P(\alpha) + P(\neg\alpha)$ can be greater or less than
 847 1 precisely because α and $\neg\alpha$ do not establish a partition of the information
 848 space.

849 Now, the question is: since we cannot rely on the classical, mutually exclusive
 850 partitions of the sample space, how can total probability theorems be stated? In
 851 order to provide such theorems for LET_F , we have to rely on the connectives \circ ,
 852 \bullet , and on the connectives defined in Fact 34. We also need a bit of terminology.

853 **Definition 43.** (*Cleavage*)

854 *Let us call a cleavage a (finite) family of propositions $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$. A cleavage*
 855 *is said to be exhaustive if $\alpha_1 \vee \alpha_2 \vee \dots \vee \alpha_n$ is a tautology, and so it covers all the*
 856 *information space, possibly with intersections. A cleavage is said to be exclusive*
 857 *when $\alpha_1 \vee \alpha_2 \vee \dots \vee \alpha_n$ are pairwise logically incompatible. In this case, it does*
 858 *not yield intersection of information (in the sense that $\alpha_i \wedge \alpha_j$ for $i \neq j$ is a*
 859 *bottom particle), and possibly does not cover the whole space. An exhaustive*
 860 *and exclusive cleavage is a partition.*

861 Items 2 to 5 of Theorem 36 cleave the information space exhaustively but not
 862 exclusively. Items 1 and 6, on the other hand, cleave the information space
 863 in parts that are exhaustive and exclusive, and so they are partitions. Notice

864 that item 3 of Theorem 36 corresponds to the six scenarios of conclusive and
 865 non-conclusive evidence that we have seen in Section 3.1. These scenarios can
 866 be graphically represented as follows:

867	$\bullet\alpha \wedge \alpha$	$\bullet\alpha \wedge \neg\alpha$	$\bullet\alpha$	$\bullet\alpha \wedge (\alpha \wedge \neg\alpha)$	$\circ\alpha \wedge \alpha$	$\circ\alpha \wedge \neg\alpha$
	1	2	3	4	5	6

868 Item 1 of Theorem 36 above emphasizes the division between non-conclusive
 869 evidence (scenarios 1 to 4) and conclusive evidence (scenarios 5 and 6), while
 870 item 6, in addition, splits the conclusive evidence into truth (5) and falsity
 871 (6). These propositions can be understood as expressing different ways we can
 872 look at the information space. The following total probability theorems can be
 873 obtained depending upon certain cleavages, based on Theorem 36.

874 **Theorem 44.** *Total probability theorems*

- 875 1. $P(\beta) = P(\beta \wedge \circ\alpha) + P(\beta \wedge \bullet\alpha)$, *w.r.t. the cleavage* $\{\circ\alpha, \bullet\alpha\}$.
- 876 2. $P(\beta) = P(\beta \wedge \alpha) + P(\beta \wedge \neg\alpha) + P(\beta \wedge \bullet\alpha) - P(\beta \wedge \alpha \wedge \bullet\alpha) - P(\beta \wedge \neg\alpha \wedge \bullet\alpha)$,
 877 *w.r.t. the cleavage* $\{\alpha, \neg\alpha, \bullet\alpha\}$.
- 878 3. $P(\beta) = P(\beta \wedge \circ\alpha \wedge \alpha) + P(\beta \wedge \circ\alpha \wedge \neg\alpha) + P(\beta \wedge \bullet\alpha) - P(\beta \wedge \bullet\alpha \wedge \alpha \wedge \neg\alpha)$,
 879 *w.r.t. cleavage* $\{\bullet\alpha \wedge \alpha, \bullet\alpha \wedge \neg\alpha, \bullet\alpha, \bullet\alpha \wedge \alpha \wedge \neg\alpha, \circ\alpha \wedge \alpha, \circ\alpha \wedge \neg\alpha\}$.
- 880 4. $P(\beta) = P(\beta \wedge \alpha) + P(\beta \wedge \neg\alpha) + P(\beta \wedge \Theta\alpha) - P(\beta \wedge \alpha \wedge \Theta\alpha) - P(\beta \wedge \neg\alpha \wedge \Theta\alpha)$,
 881 *w.r.t. the cleavage* $\{\alpha, \neg\alpha, \Theta\alpha\}$.
- 882 5. $P(\beta) = P(\beta \wedge \alpha) + P(\beta \wedge \neg\alpha) + P(\beta \wedge \approx\alpha) - P(\beta \wedge \alpha \wedge \approx\alpha) - P(\beta \wedge \neg\alpha \wedge \approx\alpha)$,
 883 *w.r.t. the cleavage* $\{\alpha, \neg\alpha, \approx\alpha\}$.
- 884 6. $P(\beta) = P(\beta \wedge \oplus\alpha) + P(\beta \wedge \sim\alpha) + P(\beta \wedge \bullet\alpha)$, *w.r.t. the cleavage* $\{\oplus\alpha, \sim\alpha, \bullet\alpha\}$.

885 *Proof.* 1. $\beta \dashv\vdash (\beta \wedge \circ\alpha) \vee (\beta \wedge \bullet\alpha)$. So, $P(\beta) = P((\beta \wedge \circ\alpha) \vee (\beta \wedge \bullet\alpha)) =$
 886 $P(\beta \wedge \circ\alpha) + P(\beta \wedge \bullet\alpha) - P(\beta \wedge \circ\alpha \wedge \bullet\alpha) = P(\beta \wedge \circ\alpha) + P(\beta \wedge \bullet\alpha)$. The remaining
 887 proofs are left to the reader. In view of Definition 3.1 (connectives $\oplus, \sim, \Theta, \approx$),
 888 some of these cleavages are equivalent.

889 □

890 4.4 Bayes' rule

891 As is well-known, Bayes' rule, or Bayes' theorem, computes the probability of
 892 an event based on previous information related to that event. The standard
 893 Bayes' rule proves that, for $P(\beta) \neq 0$:

$$894 \quad P(\alpha/\beta) = \frac{P(\beta/\alpha) \cdot P(\alpha)}{P(\beta)}$$

895 In the equation above, interpreted in terms of measures of evidence rather than
 896 standard probabilities, $P(\alpha)$ denotes the evidence available for α without taking

897 into consideration any evidence for β . The latter is supposed to affect somehow
898 the evidence for α , and so $P(\alpha/\beta)$ is the measure of the evidence for α after
899 β is taken into account. $P(\beta/\alpha)$, usually called the ‘likelihood’ in probability
900 theory, is the evidence for β when α is considered as given, and $P(\beta)$, usually
901 called the ‘marginal likelihood’, is the total evidence available for β , that takes
902 into account all the possible cases where β may occur. In what follows, we
903 define some relevant versions of Bayes’ rule. Differently from the classical case,
904 these versions are not equivalent. They show how the notion of classicality can
905 modify Bayesian probability updating.

906 **Theorem 45.** *Bayes’ Conditionalization Rules*

907

1.

$$908 \quad P(\alpha/\beta) = \frac{P(\beta/\alpha) \cdot P(\alpha)}{P(\beta/\circ\alpha) \cdot P(\circ\alpha) + P(\beta/\bullet\alpha) \cdot P(\bullet\alpha)}$$

909 *for* $P(\beta) \neq 0$, $P(\circ\alpha) \neq 0$, *and* $P(\bullet\alpha) \neq 0$.

910 *Proof.* From the definition of conditional probability and Theorem 44,
911 item 1. □

912 2.

$$913 \quad P(\alpha/\beta) = \frac{P(\beta/\alpha) \cdot P(\alpha)}{P(\beta/\alpha) \cdot P(\alpha) + P(\beta/\neg\alpha) \cdot P(\neg\alpha) + P(\beta/\bullet\alpha) \cdot P(\bullet\alpha) - P(\beta/\alpha \wedge \bullet\alpha) \cdot P(\alpha \wedge \bullet\alpha) - P(\beta/\neg\alpha \wedge \bullet\alpha) \cdot P(\neg\alpha \wedge \bullet\alpha)}$$

914 *for* $P(\beta) \neq 0$, $P(\alpha \wedge \bullet\alpha) \neq 0$, *and* $P(\neg\alpha \wedge \bullet\alpha) \neq 0$.

915 *Proof.* From the definition of conditional probability and Theorem 44,
916 item 2. □

917 3.

$$918 \quad P(\alpha/\beta) = \frac{P(\beta/\alpha) \cdot P(\alpha)}{P(\beta/\oplus\alpha) \cdot P(\oplus\alpha) + P(\beta/\sim\alpha) \cdot P(\sim\alpha) + P(\beta/\bullet\alpha) \cdot P(\bullet\alpha)}$$

919 *for* $P(\beta) \neq 0$, $P(\bullet\alpha) \neq 0$, $P(\oplus\alpha) \neq 0$, *and* $P(\ominus\alpha) \neq 0$.

920 *Proof.* From the definition of conditional probability and Theorem 44,
921 item 6. □

922 It should be clear that the process of limit can be easily established for the above
923 formulations of Bayes’ rules. If $\lim P_i(\circ\alpha) = 1$ (or equivalently $\lim P_i(\bullet\alpha) = 0$)
924 then item 1 above reduces to $P(\circ\alpha/\beta) = 1$. Analogously, if $\lim P_i(\bullet\alpha) = 0$ (or
925 equivalently $\lim P_i(\circ\alpha) = 1$), then items 2 and 3 above reduce to the standard
926 form of Bayes’ rule.

927 **5 Final remarks**

928 This paper has been conceived to be a further development of the approach
 929 to paraconsistency as preservation of evidence presented in Carnielli and Ro-
 930 drrigues (2017, 2019), where an interpretation of contradictions in terms of non-
 931 conclusive evidence was proposed. The underlying assumption is that there are
 932 no true contradictions, but rather argumentative contexts in which conflicting
 933 evidence, as well as the absence of any evidence, may occur. The valuation
 934 semantics is able to express only that there is or there is not evidence for a
 935 proposition α , while the probabilistic semantics presented here intends to ex-
 936 press the degree of evidence enjoyed by a given proposition. The acceptance of
 937 scenarios in which $P(\alpha) + P(\neg\alpha) > 1$, however, does not mean that there may
 938 be something like ‘contradictory sample spaces’, or ‘contradictory probabilistic
 939 spaces’. The latter would be the probabilistic counterpart of contradictions in
 940 reality, a view on paraconsistency not endorsed by us. In our view, it is the
 941 information available about some collection of events that can be contradictory.
 942 So, instead of talking about sample spaces, the concept of an information space
 943 has been introduced here.

944 Both LET_J and LET_F are Logics of Formal Inconsistency and Undeter-
 945 minedness suitable for an intuitive interpretation in terms of preservation of
 946 evidence and truth. The intuition regarding \circ and \bullet as ‘classically contradic-
 947 tory’ w.r.t. each other had already been presented in Carnielli, Coniglio, and
 948 Rodrigues (2019, Section 4.4). LET_F , however, as far as we know, is the first
 949 formal system where these connectives are both primitive and have the deduc-
 950 tive behavior given by rules *Cons* and *Comp*, that are in some sense analogous
 951 to explosion and excluded middle. The connective \bullet , and the fact that $\circ\alpha \vee \bullet\alpha$
 952 and $\alpha \vee \neg\alpha \vee \bullet\alpha$ are theorems of LET_F , are essential for proving total probability
 953 theorems and Bayes’ rules (Theorems 44 and 45).

954 The probabilistic semantics of LET_F has been axiomatically stated in defi-
 955 nitions 37 and 38. Accordingly, $P(\alpha) + P(\neg\alpha)$ can be greater or less than 1, and
 956 this is interpreted as scenarios, respectively, of conflicting evidence, and little
 957 or no evidence. When $P(\circ\alpha) = 1$, the classical behavior of $P(\alpha)$ and $P(\neg\alpha)$ is
 958 restored, and this is interpreted as saying that the evidence available for α and
 959 $\neg\alpha$ is subjected to the laws of standard probability theory. But $P(\circ\alpha)$ may be
 960 less than 1, and in this case, according to the axioms, it expresses the degree to
 961 which $P(\alpha)$ and $P(\neg\alpha)$ are expected to behave classically (the value of $P(\circ\alpha)$
 962 establishes constraints on the values of $P(\alpha \vee \neg\alpha)$ and $P(\alpha \wedge \neg\alpha)$, cf. Lemma 41).
 963 Accordingly, $P(\circ\alpha) < 1$ can be intuitively interpreted as expressing the reliabil-
 964 ity of the available evidence for α and $\neg\alpha$: greater reliability corresponds to a
 965 greater degree of classicality.

966 Our treatment here does not intend to express degrees of belief by means
 967 of probability measures. The notion of evidence for α , as explained in Section
 968 2.2.1, does not imply belief in α . So, the degree of evidence for α measured
 969 by a statement $P(\alpha) = \epsilon$ is not a measure of the belief of an agent in α . How-
 970 ever, nothing a fortiori prevents the formal system proposed here, together with
 971 its probabilistic semantics, of being interpreted, or used, as a tool to measure

972 degrees of belief, uncertainty, or some other relation between agents and propo-
 973 sitions. Similar remarks apply to the connective \circ . In $P(\circ\alpha) = \epsilon$, the value
 974 of ϵ expresses the degree to which it is expected that $P(\alpha)$ behaves classically.
 975 Indeed, ϵ can also be interpreted as the degree of reliability of evidence for
 976 α , coherence with previous data or with a historical series of measures of evi-
 977 dence for α , or even with a subjective ingredient, for example, as the degree of
 978 trustfulness of the belief in α , or certainty/uncertainty of α .

979 The rules for \circ and \bullet , due to their dual character, show a symmetry that
 980 deserves to be further investigated from the proof-theoretic point of view. There
 981 are some extensions of LET_J and LET_F that also deserve to be studied. The
 982 operator \bullet and the rules *Cons* and *Comp* can be added to LET_J , obtaining
 983 a logic that differs from LET_F only in the implication for the non-classical
 984 propositions. Two intuitively appealing equivalences are the following:

985 1. $\circ\alpha \dashv\vdash \circ\circ\alpha$

986 2. $\circ\alpha \dashv\vdash \circ\neg\alpha$

987 It was shown in Carnielli and Rodrigues (2017, Fact 17) that LET_J has no
 988 theorems of the form $\circ\alpha$ (the same result also holds for LET_F), and it was
 989 argued that LET_J (and so LET_F) was conceived in such a way that \circ has to be
 990 introduced from outside the formal system. This is in line with the idea that
 991 information about conclusive evidence for a proposition α comes from outside
 992 the formal system. But it is also very reasonable to suppose that once the truth
 993 value of a proposition α has been established, and so $\circ\alpha$ holds and α has classical
 994 behavior, then $\circ\alpha$, $\circ\circ\alpha$, and so on, also have classical behavior. Conversely, it is
 995 also reasonable to conclude $\circ\alpha$ from $\circ\circ\alpha$, and so on. These ideas are expressed
 996 by 1 above. The equivalence 2 above makes explicit inside the system the first
 997 part of the result achieved by Fact 31 (to wit: once $\circ\neg^n\alpha$ is proved, and so it
 998 follows that $\neg^n\alpha$ is subjected to classical logic, for any formula $\neg^m\alpha$, $m \geq 0$,
 999 $\neg^m\alpha$ is also subjected to classical logic). Valuation semantics for these rules
 1000 are straightforward, and adding these rules would produce a decidable formal
 1001 system.

1002 We believe that the probabilistic semantic relation presented in Section 4
 1003 will succeed as a tool for dealing with real argumentative contexts, including in-
 1004 vestigative scenarios and databases concerned with different degrees of evidence
 1005 attributed to propositions. But this claim needs to be further investigated.

1006 References

- 1007 P. Achinstein. Concepts of evidence. In *Evidence, Explanation, and Realism*.
 1008 Oxford University Press, 2010a.
- 1009 P. Achinstein. Why philosophical theories of evidence are (and ought to be) ig-
 1010 nored by scientists. In *Evidence, Explanation, and Realism*. Oxford University
 1011 Press, 2010b.

- 1012 A. R. Anderson and N. Belnap. *Entailment: The Logic of Relevance and Ne-*
1013 *cessity (Vol. 1)*. Princeton University Press, 1975.
- 1014 A. R. Anderson and N. D. Belnap. First degree entailments. *Mathematische*
1015 *Annalen*, 149:302–319, 1963.
- 1016 A. R. Anderson, N. Belnap, and Dunn J. M. *Entailment: The Logic of Relevance*
1017 *and Necessity (Vol. 2)*. Princeton University Press, 1992.
- 1018 N. D. Belnap. How a computer should think. In *Contemporary Aspects of*
1019 *Philosophy (ed. G. Ryle)*. Oriel Press, 1977a.
- 1020 N. D. Belnap. A useful four-valued logic. In *Modern Uses of Multiple Valued*
1021 *Logics (eds. Epstein, G. and Dunn, J. M.)*. D. Reidel, Dordrecht, 1977b.
- 1022 J. Bueno-Soler and W. Carnielli. Paraconsistent probabilities: Consistency,
1023 contradictions and Bayes’ theorem. In J. Stern, editor, *Special Issue Statistical*
1024 *Significance and the Logic of Hypothesis Testing*. Entropy 18(9), 2016. Open
1025 access <http://www.mdpi.com/1099-4300/18/9/325/htm>.
- 1026 J. Bueno-Soler and W. Carnielli. Paraconsistent probabilities, their significance
1027 and their uses. In P. Gouveia, C. Caleiro, and F. Donisio, editors, *Logic and*
1028 *Computation: Essays in Honour of Amilcar Sernadas*, pages 197–230. College
1029 Publications, 2017.
- 1030 C. Caleiro, W. Carnielli, M. E. Coniglio, and J. Marcos. Two’s company: “The
1031 Humbug of Many Logical Values”. *Logica Universalis*, 2:175–194, 2005.
- 1032 W. Carnielli and A. Rodrigues. An epistemic approach to paraconsistency: a
1033 logic of evidence and truth. *Synthese*, 196:3789–3813, 2017. doi: 10.1007/
1034 s11229-017-1621-7.
- 1035 W. Carnielli and A. Rodrigues. On epistemic and ontological interpretations
1036 of intuitionistic and paraconsistent paradigms. *Logic Journal of the IGPL*,
1037 2019. doi: 10.1093/jigpal/jzz041.
- 1038 W. Carnielli, M. E. Coniglio, and J. Marcos. Logics of formal inconsistency. In
1039 Gabbay and Guenther, editors, *Handbook of Philosophical Logic*, volume 14.
1040 Springer, 2007.
- 1041 W. Carnielli, M. Coniglio, and A. Rodrigues. Recovery operators, paraconsis-
1042 tency and duality. *Logic Journal of the IGPL*, 2019. doi: 10.1093/jigpal/
1043 jzy054.
- 1044 N. C. A. da Costa and E. Alves. A semantical analysis of the calculi C_n . *Notre*
1045 *Dame Journal of Formal Logic*, 18(4):621–630, 1977.
- 1046 L. Demey, B. Kooi, and J. Sack. Logic and probability. *Stanford Encyclopedia of*
1047 *Philosophy*, 2013. URL [https://plato.stanford.edu/archives/sum2017/
1048 entries/logic-probability/](https://plato.stanford.edu/archives/sum2017/entries/logic-probability/).

- 1049 J. M. Dunn. Intuitive semantics for first-degree entailments and ‘coupled trees’.
1050 *Philosophical Studies*, 29(3):149–168, 1976.
- 1051 J. M. Dunn. Information in computer science. In P. Adriaans & J. van Benthem,
1052 editor, *Philosophy of Information. Volume 8 of Handbook of the Philosophy*
1053 *of Science*, pages 581–608. Elsevier, 2008.
- 1054 B. Fitelson and A. Hájek. Declarations of independence. *Synthese*, 194(10):
1055 3979–3995, 2017.
- 1056 M. Fitting. Paraconsistent logic, evidence, and justification. *Studia Logica*, 105
1057 (6):1149–1166, 2016a.
- 1058 M. Fitting. Justifications that might be wrong (abstract). In *Modalities, Con-*
1059 *ditionals, and Values: A Symposium on Philosophical Logic in Celebration of*
1060 *the Centenary of Georg Henrik von Wright. May 23-25, 2016, Helsinki, Fin-*
1061 *land*, 2016b. URL [https://www.cs.helsinki.fi/u/hakli/vw/abstracts.](https://www.cs.helsinki.fi/u/hakli/vw/abstracts.html)
1062 [html](https://www.cs.helsinki.fi/u/hakli/vw/abstracts.html).
- 1063 C. M. Grinstead and J. L. Snell. *Introduction to Probability*. American Mathe-
1064 matical Society, Providence, RI, 1997.
- 1065 T. Kelly. Evidence. *The Stanford Encyclopedia of Philosophy (Fall 2014, ed. E.*
1066 *Zalta)*, 2014.
- 1067 O. Lahav, J. Marcos, and Y. Zohar. It ain’t necessarily so: basic sequent systems
1068 for negative modalities. In L. Beklemishev, S. Demri, and A. Máté, editors,
1069 *Advances in Modal Logic*. College Publications, London, 2016.
- 1070 A. Loparic. A semantical study of some propositional calculi. *The Journal of*
1071 *Non-Classical Logic*, 3(1):73–95, 1986.
- 1072 A. Loparic. Valuation semantics for intuitionistic propositional calculus and
1073 some of its subcalculi. *Principia*, 14(1):125–133, 2010.
- 1074 A. Loparic and E. Alves. The semantics of the systems C_n of da Costa. In
1075 A. I. Arruda, N. C. A. da Costa, and A. M. Sette, editors, *Proceedings of the*
1076 *III Brazilian Conference on Mathematical Logic, Recife, 1979*, pages 161–172.
1077 Brazilian Logic Society, São Paulo, 1979.
- 1078 J. Łukasiewicz. *Die logischen Grundlagen der Wahrscheinlichkeitsrechnung*.
1079 Spólka Wydawnicza Polska, Krakow, 1913.
- 1080 J. Marcos. Nearly every normal modal logic is paranormal. *Logique et Analyse*,
1081 48:279–300, 2005.
- 1082 H. Omori and H. Wansing. 40 Years of FDE: An Introductory Overview. *Studia*
1083 *Logica, Special Issue “40 Years of FDE”*, 105:1021–1049, 2017. doi: 10.1007/
1084 s11225-017-9748-6.

- 1085 G. Shafer. *A Mathematical Theory of Evidence*. Princeton University Press,
1086 1976.
- 1087 A. Tarski. Wahrscheinlichkeitslehre und mehrwertige Logik. *Erkenntnis*, 5:
1088 174–175, 1935.
- 1089 D. van Dalen. *Logic and Structure*. Springer, 4th edition, 2008.
- 1090 B. Weatherson. From classical to intuitionistic probability. *Notre Dame Journal*
1091 *of Formal Logic*, 44:111–123, 2003.
- 1092 L. Zadeh. Review of Glenn Shafer, *A Mathematical Theory of Evidence*. *AI*
1093 *Magazine*, 5(3):81–83, 1984.