

Differentiation of Discrete Functions

Young W Lim

Nov 13, 2024

Copyright (c) 2024 Young W. Lim.

Permission is granted to copy, distribute and/or modify this document under the terms of the GNU Free Documentation License, Version 1.2 or any later version published by the Free Software Foundation; with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts.

This work is licensed under a Creative Commons
"Attribution-NonCommercial-ShareAlike 3.0 Unported"
license.



Based on
Introduction to Matrix Algebra, Autar Kaw
<https://ma.mathforcollege.com>

Outline

- 1 Approximations of a first derivative
 - Direct Fit Polynomials

Outline

- 1 Approximations of a first derivative
 - Direct Fit Polynomials

Direct Fit Polynomials

given $n+1$ data points

$$(x_0, y_0), (x_1, y_1), \dots, (x_{n-1}, y_{n-1}), (x_n, y_n)$$

one can fit a n^{th} order polynomial given by

$$P_n(x) = a_0 + a_1x + \dots + a_{n-1}x^{n-1} + a_nx^n$$

to find the first derivative

$$P'_n(x) = \frac{dP_n(x)}{dx} = a_1 + 2a_2x + \dots + (n-1)a_{n-1}x^{n-2} + na_nx^{n-1}$$

similarly other derivatives can be found

Direct Fit Polynomials

given $(n+1)$ data points

$$(x_0, y_0), (x_1, y_1), \dots, (x_{n-1}, y_{n-1}), (x_n, y_n)$$

one can fit a n^{th} order polynomial given by

$$P_n(x) = a_0 + a_1x + \dots + a_{n-1}x^{n-1} + a_nx^n$$

to find the first derivative

$$P'_n(x) = \frac{dP_n(x)}{dx} = a_1 + 2a_2x + \dots + (n-1)a_{n-1}x^{n-2} + na_nx^{n-1}$$

similarly other derivatives can be found

Lagrange Polynomials (1)

given $(n+1)$ data points

$$(x_0, y_0), (x_1, y_1), \dots, (x_{n-1}, y_{n-1}), (x_n, y_n)$$

one can fit a n^{th} order Lagrange polynomial given by

$$f_n(x) = \sum_{i=0}^n L_i(x) f(x_i)$$

where n in $f_n(x)$ stands for the n^{th} order polynomial that approximates the function $y = f(x)$

given at $(n+1)$ data points

as $(x_0, y_0), (x_1, y_1), \dots, (x_{n-1}, y_{n-1}), (x_n, y_n)$

Lagrange Polynomials (2-1)

one can fit a n^{th} order Lagrange polynomial given by

$$f_n(x) = \sum_{i=0}^n L_i(x) f(x_i)$$

where n in $f_n(x)$ stands for the n^{th} order polynomial for the function $y = f(x)$

$$f_n(x) = a_0 + a_1x + \cdots + a_{n-1}x^{n-1} + a_nx^n = \sum_{i=0}^n L_i(x) f(x_i)$$

given at $(n+1)$ data points as $(x_0, y_0), (x_1, y_1), \dots, (x_{n-1}, y_{n-1}), (x_n, y_n)$, and

$$L_i(x) = \prod_{j=0, j \neq i}^n \frac{x - x_j}{x_i - x_j}$$

Lagrange Polynomials (2-2)

given a set of $(n+1)$ nodes $\{x_0, x_1, \dots, x_n\}$,
 which must all be distinct, $x_j \neq x_i$ for indices $j \neq i$,

the **Lagrange basis** for polynomials of degree $\leq n$ for those nodes
 is the set of polynomials $\{L_0(x), L_1(x), \dots, L_n(x)\}$

$$L_i(x) = \prod_{j=0, j \neq i}^n \frac{x - x_j}{x_i - x_j}$$

$$= \frac{x - x_0}{x_i - x_0} \cdot \frac{x - x_1}{x_i - x_1} \cdots \frac{x - x_{i-1}}{x_i - x_{i-1}} \cdot 1 \cdot \frac{x - x_{i+1}}{x_i - x_{i+1}} \cdot \frac{x - x_{i+2}}{x_i - x_{i+2}} \cdots \frac{x - x_n}{x_i - x_n}$$

https://en.wikipedia.org/wiki/Lagrange_polynomial

Lagrange Polynomials (2-3)

each **Lagrange basis** of degree n take values

$L_i(x_j) = 0$ if $j \neq i$ and $L_i(x_i) = 1$.

Using the **Kronecker delta** this can be written $L_i(x_j) = \delta_{ij}$.

Each **basis polynomial** can be explicitly described by the product:

$$\begin{aligned}
 L_i(x) &= \prod_{j=0, j \neq i}^n \frac{x - x_j}{x_i - x_j} \\
 &= \frac{x - x_0}{x_i - x_0} \cdot \frac{x - x_1}{x_i - x_1} \cdots \frac{x - x_{i-1}}{x_i - x_{i-1}} \cdot 1 \cdot \frac{x - x_{i+1}}{x_i - x_{i+1}} \cdot \frac{x - x_{i+2}}{x_i - x_{i+2}} \cdots \frac{x - x_n}{x_i - x_n}
 \end{aligned}$$

https://en.wikipedia.org/wiki/Lagrange_polynomial

Lagrange Polynomials (2-4)

$$\begin{aligned}
 L_i(x_j) &= \prod_{k=0, k \neq i}^n \frac{x_j - x_k}{x_i - x_k} \\
 &= \frac{x_j - x_0}{x_i - x_0} \cdot \frac{x_j - x_1}{x_i - x_1} \cdots \frac{x_j - x_{i-1}}{x_i - x_{i-1}} \cdot 1 \cdot \frac{x_j - x_{i+1}}{x_i - x_{i+1}} \cdot \frac{x_j - x_{i+2}}{x_i - x_{i+2}} \cdots \frac{x_j - x_n}{x_i - x_n} \\
 &= 0 \quad \because \exists j = k \neq i
 \end{aligned}$$

$$\begin{aligned}
 L_i(x_i) &= \prod_{k=0, k \neq i}^n \frac{x_i - x_k}{x_i - x_k} \\
 &= \frac{x_i - x_0}{x_i - x_0} \cdot \frac{x_i - x_1}{x_i - x_1} \cdots \frac{x_i - x_{i-1}}{x_i - x_{i-1}} \cdot 1 \cdot \frac{x_i - x_{i+1}}{x_i - x_{i+1}} \cdot \frac{x_i - x_{i+2}}{x_i - x_{i+2}} \cdots \frac{x_i - x_n}{x_i - x_n} \\
 &= 1
 \end{aligned}$$

https://en.wikipedia.org/wiki/Lagrange_polynomial

Lagrange Polynomials (3)

hen to find the first derivative, one can differentiate $f_n(x)$ for other derivatives.

$$f_n(x) = \sum_{i=0}^n L_i(x) f(x_i)$$

For example, the second order Lagrange polynomial passing through $(x_0, y_0), (x_1, y_1), (x_2, y_2)$

$$f_2(x) = L_0(x)f(x_0) + L_1(x)f(x_1) + L_2(x)f(x_2)$$

$$L_0(x) = \prod_{j=0, j \neq i}^2 \frac{x - x_j}{x_0 - x_j}, \quad L_1(x) = \prod_{j=0, j \neq i}^2 \frac{x - x_j}{x_1 - x_j}, \quad L_2(x) = \prod_{j=0, j \neq i}^2 \frac{x - x_j}{x_2 - x_j}$$

Lagrange Polynomials (4)

the second order Lagrange polynomial passing through $(x_0, y_0), (x_1, y_1), (x_2, y_2)$

$$f_2(x) = L_0(x)f(x_0) + L_1(x)f(x_1) + L_2(x)f(x_2)$$

$$L_0(x) = \prod_{j=0, j \neq i}^n \frac{x - x_j}{x_0 - x_j}, \quad L_1(x) = \prod_{j=0, j \neq i}^n \frac{x - x_j}{x_1 - x_j}, \quad L_2(x) = \prod_{j=0, j \neq i}^n \frac{x - x_j}{x_2 - x_j}$$

$$L_0(x) = \frac{x - x_1}{x_0 - x_1} \cdot \frac{x - x_2}{x_0 - x_2}, \quad L_1(x) = \frac{x - x_0}{x_1 - x_0} \cdot \frac{x - x_2}{x_1 - x_2}, \quad L_2(x) = \frac{x - x_0}{x_2 - x_0} \cdot \frac{x - x_1}{x_2 - x_1}$$

$$f_2(x) = \frac{x - x_1}{x_0 - x_1} \cdot \frac{x - x_2}{x_0 - x_2} f(x_0) + \frac{x - x_0}{x_1 - x_0} \cdot \frac{x - x_2}{x_1 - x_2} f(x_1) + \frac{x - x_0}{x_2 - x_0} \cdot \frac{x - x_1}{x_2 - x_1} f(x_2)$$

Lagrange Polynomials (5)

the second order Lagrange polynomial passing through $(x_0, y_0), (x_1, y_1), (x_2, y_2)$

$$f_2(x) = L_0(x)f(x_0) + L_1(x)f(x_1) + L_2(x)f(x_2)$$

$$f_2(x) = \frac{x-x_1}{x_0-x_1} \cdot \frac{x-x_2}{x_0-x_2} f(x_0) + \frac{x-x_0}{x_1-x_0} \cdot \frac{x-x_2}{x_1-x_2} f(x_1) + \frac{x-x_0}{x_2-x_0} \cdot \frac{x-x_1}{x_2-x_1} f(x_2)$$

$$\frac{d}{dx}(x-x_1)(x-x_2) = (x-x_2) + (x-x_1) = 2x - (x_1 + x_2)$$

$$\frac{d}{dx} f_2(x) = \frac{2x - (x_1 + x_2)}{(x_0 - x_1)(x_0 - x_2)} f(x_0) + \frac{2x - (x_0 + x_2)}{(x_1 - x_0)(x_1 - x_2)} f(x_1) + \frac{2x - (x_0 + x_1)}{(x_2 - x_0)(x_2 - x_1)} f(x_2)$$

$$\frac{d^2}{dx^2} f_2(x) = \frac{2f(x_0)}{(x_0 - x_1)(x_0 - x_2)} + \frac{2f(x_1)}{(x_1 - x_0)(x_1 - x_2)} + \frac{2f(x_2)}{(x_2 - x_0)(x_2 - x_1)}$$

Tangent Lines

- as $h \rightarrow 0$, $Q \rightarrow P$
and the **secant line** \rightarrow the **tangent line**
- the slope of the **tangent line**

$$\begin{aligned}m_{\text{tangent}} &= \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{(a+h) - a} \\ &= \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}\end{aligned}$$

