

RESOLUTION 2.12

GUIDELINES FOR THE USE OF ACOUSTIC DETERRENT DEVICES

The Meeting of the Parties to the Agreement on the Conservation of Cetaceans of the Black Sea, Mediterranean Sea and contiguous Atlantic area:

On recommendation of the ACCOBAMS Scientific Committee,

Aware of the fact that cetaceans are particularly vulnerable to disturbance at certain stages of their life cycle,

Particularly conscious of the interaction of Cetaceans with some coastal and artisanal fisheries and the resulting conflicts,

Convinced that the importance of coastal and artisanal fisheries in the sustainable development of the ACCOBAMS range States implies the integration of conservation activities with socio-economic development,

Noting that acoustic deterrents represent a new technique for the remediation of by-catches in fishing gears effects of which on nature and on different components of biodiversity cannot be fully assessed or predicted at present as they may produce significant noise pollution and possibly exclude cetaceans from certain areas,

Noting also that widespread use of acoustic deterrents in fishery and aquaculture operations in the Mediterranean indicates that these could raise conservation concerns,

Recalling that:

- Article II.3 in which Parties shall apply the conservation, research and management measures, which shall address inter alia the assessment and management of human-cetacean interactions and in implementing this measures, they shall apply the precautionary principle,
- The Conservation Plan, which is fully part of the Agreement, requires the Parties to establish and implement legislative, regulatory or administrative measures:
 - To minimize adverse effects of fisheries on the conservation status of cetaceans,
 - For impact assessments to be carried out in order to provide a basis for either allowing or prohibiting the continuation or the future development of activities that may affect cetaceans or their habitat in the Agreement area as well as establishing the conditions under which such activities may be conducted,

Recalling also Resolution 2.16 on Man-made noise,

Taking into account the advice of the Scientific Committee on the ineffectiveness and harmful potential of AHDs¹ to alleviate conflicts between cetaceans and fisheries or mariculture operations in the Agreement and considering pingers² to be less invasive than AHD and their use is, in principle, oriented towards cetacean conservation,

Aware of the scarcity of controlled studies of the efficacy of pingers in reducing by-catches in the Agreement area,

¹ Acoustic devices used to avoid dolphin interaction with fishing activities and aquaculture have the potential to adversely impact cetacean and other animal populations. They may damage the hearing of cetaceans and, if used extensively, exclude them from significant portions of their habitat. These undesired effects are well documented for "acoustic harassment devices" (AHD) which are used, for example, to prevent animals from approaching aquaculture facilities; AHD produce high source levels (>185dB re 1 P at 1m) and operate primarily in the mid to high frequency range (c.5-30kHz).

² "Pingers" are used primarily to alert cetaceans to existence of fishing gear and thus avoid entanglement. These are low-intensity (generally <150dB re 1 P at 1m) sources that operate in the mid to high frequencies between about 2.5-109kHz, with harmonics to much higher frequencies (Reeves *et al.*,2001). These devices are considered to be less invasive than AHD and their use is, in principle, oriented towards cetacean conservation.

1. *Adopts* the “Guidelines for technical measures to minimize cetacean-fishery conflicts in the Mediterranean and Black Seas” in Annex 1;
2. *Urges* the Parties:
 - To strictly regulate the use of AHDs to alleviate conflicts between cetaceans and fisheries or mariculture operations in the Agreement area;
 - To strongly recommend that the use of pingers, where authorized and appropriate, only be conducted with controlled studies to ensure that they are an effective mitigation measure;
 - To link any use of pingers with an observer scheme designed to monitor their effectiveness over time;
3. *Charges* the Scientific Committee to update technological informations on this topic, with a view to developing common sets of guidelines for use in the ACCOBAMS area;
4. *Charges* the Secretariat, in relation the regional fisheries bodies and all financing agencies and donor Countries to consider helping developing Countries acquire and use relevant technology, and with appropriate education and training of fishermen.

ANNEX 1

GUIDELINES FOR TECHNICAL MEASURES TO MINIMISE CETACEAN-FISHERY CONFLICTS IN THE MEDITERRANEAN AND BLACK SEAS

Edited by Simon Northridge, Caterina Fortuna and Andrew Read

Administrative introduction

These Guidelines have been developed in response to requests made by member states to the ACCOBAMS Secretariat for advice on how to minimize conflicts between small cetaceans and fisheries in the Mediterranean and Black Seas. Implicit in the Agreement between member states is the assertion that culling is an inappropriate and usually ineffective means of addressing such conflicts with unacceptable consequences for the conservation of small cetaceans.

There is still much uncertainty over many aspects of the mitigation tools that have been used in attempts to minimise cetacean fishery conflicts. In some cases the efficacy of the methods used is still questionable. These Guidelines have been compiled with the knowledge that there are no certain solutions to any of these problems, and that much scientific work remains to be done to understand how they can be resolved in the long term. Governments are urged to support research efforts in this area.

Terminology

Conflicts between fisheries and cetaceans generally take one or both of two forms. These are: the accidental capture of cetaceans in fishing operations (**bycatch**) and the **depredation** of fishing gear by cetaceans, leading to loss of catch and damage to fishing gear. In many cases these two problems occur in the same fisheries, and resolving the latter problem may help to resolve the former.

The 2001 ICRAM workshop (Reeves *et al* 2001) recognized a variety of potential mitigation methods to deal with cetacean bycatch and depredation of fish catches in static net fisheries in the Mediterranean. Perhaps the most widely-used methods involve acoustic devices of one form or another. The ICRAM workshop recognized two major categories of acoustic mitigation devices: Acoustic Harassment Devices (AHDs) and Acoustic Deterrent Devices (ADD), including pingers.

Pingers are relatively low-intensity (generally <150dB re 1 μ P at 1m) battery-powered sound generators that operate in the mid to high sound frequencies (between about 10 kHz to around 100 kHz). Pingers are usually designed to prevent small cetaceans from becoming entangled in gill nets, however a new generation of such devices has been designed to mitigate the depredation. At the other extreme, AHDs are designed to work by causing pain, discomfort or irritation to potential predators, and have been developed primarily with the aim of discouraging seals from approaching caged fish. Pingers are usually small (hand-sized) devices that run for weeks, months or years on small batteries. AHDs, in contrast, have relatively high sound source levels (typically >185dB re 1 μ P at 1m) and operate primarily in the low to mid frequency range (c. 5-30 kHz). They are typically bulky pieces of equipment powered from mains electricity or large lead-acid vehicle batteries. As they have primarily been designed with seals in mind, AHDs produce sound within pinniped hearing sensitivities, which are typically lower than those of small odontocetes. Not all acoustic devices necessarily fall into one category or another and the difference between the two types of device, especially in terms of their acoustic output, is qualitative.

How do acoustic devices work?

It remains unclear how most of these devices work and a range of possible mechanisms has been postulated. These include: in the case of AHDs discomfort; scaring; deterring; masking of the animals' acoustic detection senses; or simple confusion. However, in most cases the exact behavioural mechanism by which AHDs work is unclear.

In some cases, it appears that ADDs function in an aversive manner. For example, several studies have shown that harbour porpoises (*Phocoena phocoena*) and, to a lesser extent, bottlenose dolphins (*Tursiops truncatus*) avoid pingers

(Koschinski & Culik 1997, Kastelein *et al.* 2000, Culik *et al.* 2001, Laake *et al.* 1998, Cox *et al.* 2003, Goodson *et al.* 1994, Anonymous 2003b). Further details of this research are available on the ACCOBAMS website. Nevertheless, the scope of this research is limited. The response of small cetaceans to any acoustic stimulus is likely to be context-dependent and our understanding of their reaction to any such sound is limited at best.

Do they work?

Both practical experience and several experimental studies have shown that pingers are able to significantly reduce the bycatch of harbour porpoises in gillnets (Kraus *et al.* 1997, Gearin *et al.* 2000, SMRU *et al.* 2000, Larsen *et al.* 2000). Several other studies have shown a similar effect with other small cetacean species including the striped dolphin (*Stenella coeruleoalba*), common dolphin (*Delphinus delphis*) and franciscana (*Pontoporia blainvillei*) (Barlow and Cameron 1999, Imbert *et al.* 2001, Imbert *et al.* 2002, Bordino *et al.* 2002, Bordino *et al.* 2004). The exact reduction in by-catch depends on many factors including the behavioural response of the species in question and the degree to which devices are properly used and maintained.

Early types of AHD were shown to be ineffective in the medium to long term in several experimental studies in North America. Pinnipeds habituated to these devices and sometimes came to regard them as a dinner bell, resulting in increased depredation at salmon capture sites (Mate and Harvey 1980). Since these early studies, a new generation of AHDs has been designed for the salmon aquaculture industry. Unfortunately, there have been very few experimental studies to show whether or not these new generation AHDs are effective in reducing depredation. One study in Sweden, in which one model of 'seal scarer' AHD was used close to a salmon netting station was shown to be effective over a short period of several weeks (Westerberg *et al.* 1999).

Several studies in the Mediterranean have tested the effectiveness of acoustic deterrents in reducing damage to gear and depredation caused by bottlenose dolphins. The results of these studies, while promising in some cases, do not present a clear and straightforward answer to the question. Studies to date are summarised in Box 1 below.

Concerns about the use of acoustic devices

Several concerns have been raised about the use of acoustic devices. Louder devices, such as AHDs designed to keep pinnipeds away from fish farm sites, have been shown to exclude cetaceans from large areas (Olesiuk *et al.* 2002, Morton and Symonds 2002, Johnston 2002). Concerns have, therefore, been raised that the widespread use of such devices may significantly reduce the habitat available for cetaceans in an area. This concern has also been expressed with respect to the large-scale use of pingers, although the spatial scale of such exclusion is likely to be much smaller for each individual device. Small-scale exclusion has been reported for harbor porpoises around active pingers (Culik *et al.* 2001, Berggren *et al.* 2002), but intensive use of such devices over a large area may be a cause for concern if small cetaceans are likewise excluded from significant parts of their habitat. The potential exclusion effect of pingers may be ameliorated to some extent by the finding that continued exposure to such devices may lead to a diminution (though not a disappearance) of the behavioral response and, thus, the area of exclusion (Cox *et al.* 2001).

The possibility has also been raised that some of the AHDs in use around aquaculture sites may cause physical damage to animals nearby. It might be assumed that animals would choose to remain at a comfortable distance from a very loud sound source, but in situations in which aversive signals are only emitted sporadically it is possible that a cetacean or seal might get close enough to a sound source to suffer auditory damage if the device was activated. Theoretical studies suggest that auditory damage would be possible for cetaceans within 10m of a sound source. Pinnipeds, with less sensitive hearing, are less likely to be damaged unless they were even closer (Gordon and Northridge, 2002; Taylor *et al.* 1997).

In the Mediterranean, where small populations of the highly endangered Mediterranean monk seal still survive, there are important concerns about the possibility of both habitat exclusion and hearing damage to seals as a result of the use of AHDs (Reeves *et al.* 2001).

Depredation – approaches to minimizing the problem

There are numerous accounts of dolphins depredating fisheries in the Mediterranean, and more details of these can be found on the ACCOBAMS website (http://www.accobams.org/index_science.htm). Fisheries involved include hook and line fisheries, purse seine or *lampara* fisheries and gillnet fisheries. While not the only species involved, bottlenose dolphins appear to be the most frequently implicated.

Member States in the ACCOBAMS area have committed themselves to protecting cetaceans, and thus have a duty to assist fishermen in finding appropriate means of minimizing these conflicts. Experience in many areas shows that if fishermen are not given appropriate assistance and guidance that they may resort to inappropriate measures to deal with the problem. Appropriate mitigation measures should therefore be sought and encouraged by Member States.

At present there does not appear to be any one simple panacea that will solve the problem of depredation. It is likely that solutions will be case-specific, and the national authorities of member states will need to determine which the most likely routes to resolve the problem. These guidelines are intended to summarize information at present and assist national or regional authorities to find the most promising avenues. It should be stressed that at present there has been no demonstration of long term effectiveness of any solution.

Acoustic mitigation measures represent a potential avenue that may lead to a solution, but many other appropriate ideas should also be explored, including changes in fishing practices and behavioural conditioning of animals (Reeves *et al.* 2001). Member states should be encouraged to explore such ideas.

Several acoustic deterrents are currently being marketed for use in the ACCOBAMS region to minimise dolphin depredation. It is important to note that no study of such devices has yet shown anything more than a short-term effect. Further trials are urgently required, particularly as there are concerns that animals may habituate to acoustic deterrent signals over time and resume depredation. A summary of the trials conducted so far is given in Box 1. At the present time, no acoustic device has been shown effective at reducing depredation over the medium to long-term.

The acoustic devices marketed to reduce depredation are all relatively quiet, none approaching the sound source levels achieved in the AHDs used at aquaculture sites. This is largely because AHDs are very expensive and require significant power inputs, whereas most of the lower power devices are less expensive and run on standard alkaline or lithium cells. Box 2 lists some of the available devices.

Not all trials done so far have involved battery-powered sound sources, and some have relied on physical sound production using bells, tubes or clangers (see Box 1). Although these sounds may reduce depredation over the very short term, their effects are not long-lasting.

As some of these devices may effectively limit cetacean habitat availability, member states should be aware of where and how they are being used, and should consider ways to monitor their use. If certain devices are shown to be effective at reducing depredation over the long-term, it may be advisable to certify them for use as mitigation tools. Member States should determine the number of users, the number and type of devices, their output levels, the exposure schedule, the gear type on which they are being used, the area and season of use and the number of 'target' and 'non target' species present (notably monk seals). ACCOBAMS can provide a central registry to maintain these data. Further details of the number of units that have been sold to certain areas could usefully be obtained from manufacturers.

The main species involved in depredation is the bottlenose dolphin *Tursiops truncatus*. This species, like other cetaceans, may show an obvious **startle** reaction to novel stimuli that could lead to excessively optimistic expectations by the fishermen. In fact, this species learns rapidly, is extremely adaptable and likely to habituate in the long run to almost any noise. Therefore, alternative mitigation strategies or "combined approaches" - such as changes in fishing practices or behavioural conditioning should be favoured.

Overall, acoustic tools to minimise dolphin predation should be used only in an experimental manner. Government agencies should continue to learn how and if they work, and in what circumstances, and also the nature and extent of

any ill-effects that they might have, including habituation to the signal. With adequate co-operation and transfer of experience, much may be learned with little expenditure.

Member states should also be aware that other approaches, such as changes in fishing practice or behavioural conditioning, may also prove useful avenues for further research.

Bycatch –unintentional capture in fishing operations

There are numerous records of bycatch of cetaceans in the ACCOBAMS area. Almost all species of cetaceans that are present in any number in the ACCOBAMS area have been recorded taken in some fishing operation or other. In the Black Sea the largest number of animals taken are harbour porpoises. In the Mediterranean and Contiguous Atlantic areas common and striped dolphins are the species most often recorded. A summary of information on bycatches is presented in Box 3.

European Council regulation 812/2004 will require the use of pingers in many northern European gill and entangling net fisheries from 2005 – 2006. The intention of this regulation is primarily to minimise bycatch of harbour porpoises in EU waters. As noted above, pingers have been shown to be effective in reducing porpoise bycatch in a number of fisheries in Europe and North America, and there is no evidence yet that their effectiveness is diminished through time. It should be noted that there have been at least two studies in which bycatch of delphinid species in driftnets has been demonstrably reduced through the use of pingers. Box 2 summarises the types of pinger that are currently available to reduce bycatch, and the tests that have been carried out to show that they work.

It must also be recognised that bycatch of cetaceans cannot ever be completely eliminated by the use of acoustic devices. Pingers have been shown to reduce porpoise bycatch by 90% or more in carefully controlled field experiments. Similar studies have shown a reduction of dolphin bycatch by 80% or more.

Where pinger use has been mandated in other areas, including northern Europe, accompanying observer/monitoring programmes have been mandated to ensure that the efficacy of these devices is maintained. This is even more important where delphinids are concerned, as they may be less easily deterred from entanglement than porpoises.

Any intention to deploy pingers should be preceded by a practicability trial in which selected vessels are equipped with the devices so that deployment issues can be addressed. Experience elsewhere shows that while one pinger may work in one fishery, unexpected problems may arise in another fishery. Issues of concern include how the devices are attached to the net, how they affect fishing efficiency and whether they lead to net fouling. Specific expertise to address these issues can be made available through the ACCOBAMS Secretariat.

Other issues, including spacing, costs, battery replacement, and enforcement (where this is needed) need to be considered in advance of any deployment programme. Again, expertise in these areas is available and can be contacted through the ACCOBAMS secretariat.

As with measures to reduce depredation, acoustic approaches are not the only possible solution. Other approaches may include, on a case by case basis, time or area closures for fisheries, or switching to other gear types.

Final remarks

The possible adverse impacts of acoustic devices on cetaceans, at both individual and population level, remain poorly known. Furthermore, their effectiveness in reducing depredation is still in the process of being assessed. There is scientific evidence that pingers may reduce the by-catch of harbour porpoises and other small cetaceans in some fisheries. It is still too early to say whether acoustic devices will be effective in reducing depredation over the long term. More focused, long-term research on these topics is urgently needed.

Further information can also be accessed at the following website:

ACCOBAMS

www.accobams.org

Cetaceans Bycatch Resource Center

www.cetaceanbycatch.org

International Dolphin Conservation Programme

<http://europa.eu.int/scadplus/printversion/en/lvb/l28083.htm>

Summary of current legislation for the conservation of cetaceans

http://europa.eu.int/comm/fisheries/doc_et_publ/liste_publi/studies/bycatch/07_10legislation.htm

National Marine Fisheries Service

<http://www.nmfs.noaa.gov/bycatch.htm>

Other information

http://europa.eu.int/comm/fisheries/doc_et_publ/liste_publi/studies/bycatch/contents.htm

BOX 1: Studies examining effectiveness of acoustic deterrents

Species	Type of interaction	Fishery	Author	Country	Device/Manufacturer
Harbour porpoise (<i>Phocoena phocoena</i>)	Bycatch	Bottom set nets	Larsen 1999, Larsen <i>et al.</i> 2002	Denmark	Pinger/AQUAtec Sub Sea Ltd.
Harbour porpoise	Bycatch	Bottom set nets	Kraus <i>et al.</i> 1997, Trippel <i>et al.</i> 1999, Gearin <i>et al.</i> 2000	Canada, and USA	Pinger/Dukane Corporation
Common dolphin (<i>Delphinus delphis</i>)	Bycatch	Drift nets	Barlow and Cameron 2003	USA	Pinger/Dukane Corporation
Striped dolphins (<i>Stenella coeruleoalba</i>)	Bycatch	Drift nets	Imbert <i>et al.</i> 2002	France	Pinger/AQUAtec Sub Sea Ltd.
Bottlenose dolphins (<i>Tursiops truncatus</i>)	Depredation	Set nets	Goodson <i>et al.</i> 2001	Italy	Pinger/AQUAtec Sub Sea Ltd.
Bottlenose dolphins	Depredation	Set nets	Gazo <i>et al.</i> 2002 also as IWC paper in Shimonoseki	Spain	Pinger/AQUAtec Sub Sea Ltd.
Bottlenose dolphins	Depredation	Set nets	Northridge <i>et al.</i> 2003, Vernicos <i>et al.</i> 2003	Greece	Pinger/SaveWave BV
Bottlenose dolphins	Depredation	Set nets	Anonymous 2003a	Italy	Pinger/STM Dolphin Deterrent Device
Bottlenose dolphins	Depredation	Set nets, Purse seine	Ben Naceur 1994, Zahri <i>et al.</i> 2004	Morocco, Tunisia	Dolphin scaring tube/handmade
Franciscana (<i>Pontoporia blainvillei</i>)	Bycatch	Set nets	Bordino 2003 and Bordino <i>et al.</i> 2004	Argentina	Pinger/AIRMAR

BOX 2: Available deterrent devices

Produced by	Dukane (dismissed)	Aquatec			Savewave		Airmar	Fumunda	STM
Model	Netmark 1000	Aquamark 100 Porpoise deterrent	Aquamark 200 Acoustic Cetacean Deterrent	Aquamark 300 Pinger	Endurance	White Saver & Black Saver	Gillnet pinger	FMDP2000	DDD Dolphin Dissuasive Device
Gear	Gillnets and drifnets	Gillnets	Gill, drift and trammel nets	Gillnets	Gill and trammel nets	Gill, trammel and trawling nets	Gillnets	Driftnet	Trammel nets
Mitigation use		bycatch	Depredation and bycatch	bycatch	depredation	depredation	bycatch	bycatch	depredation
Frequency (kHz)	10	20-160	5-160	10 (tonal)		5-90 & 30-160	10	10	1-500
Source level (dB re 1µPa at 1m)	130	145	145	132	155	155	132	130-134	NA
High-frequency harmonics	Yes	Yes	Yes	Yes	Yes	Yes	Yes	No	Yes
Pulse duration (ms)	3	2-3		3		2-9	3	3	NA
Inter-pulse period (s)	4	4-30	4-30	4		4-16	4	4	NA
Wet switch	No	Yes	Yes	Yes	Yes	Yes	No	Yes	Yes
Battery	4 Alkaline AA cells	1 D-Cell Alkaline	1 D-Cell Alkaline	1 D-Cell Alkaline	Sealed 9 v unit		1 D-Cell Alkaline	1 lithium	4 alkaline 1,5V
Life	800 hours	1.5-2 years	1.5-2 years	1.5-4 years	8000 hours	2000 hours	> 1 year	15 months	3 months
Battery change	Yes	No	No	No	Yes	Yes	Yes	Yes	?
# of emitters	1	1	1	1	1	1	1		NA
Maximum depth	200	200	200	200	200	200	275	200	300
Distance between pingers	100	200	200	200				100	200
Dimensions	168 x 55mm (Ø)	164mm x 58mm (Ø)	164mm x 58mm (Ø)	164mm x 58mm (Ø)	200mm x 60mm		156mm x 53mm (Ø)	152mm x 46mm (Ø)	185mm x 61mm (Ø)
Weight (g)	400	410	410	370	400	Trawls: 6 units per net	408	230	740
Price (Euro)	Discontinued	100	100	100	55-70	55-70	44.72	74.80	223

Web site	NA	www.netPinger.net	www.savewave.net	www.airmair.com	www.fumunda.com	www.stm-products.com
----------	----	--	--	--	--	--

Box 3: Bycatch summary of information for the ACCOBAMS area

Gear Type	Nation	Season	Location	Target species	Bycatch species	Known or suspected	Monitored/ Estimated
Drift nets ("spadara" and other types) (mesh size 18 to 42 cm)	Morocco, Turkey, France, Italy, a few vessels are also present in Albania, Algeria, Greece, Monaco	April-August	Mediterranean	<i>Xiphias gladius</i> , <i>T. alalunga</i>	<i>S. coeruleoalba</i> , <i>Ziphius cavirostris</i> (<i>Globicephala</i> spp.), <i>D. delphis</i> , <i>Grampus griseus</i> , <i>Physeter macrocephalus</i> , <i>Balaenoptera physalus</i> , <i>B. acutorostrata</i>	Known	Monitored and extrapolated: Di Natale <i>et al.</i> , 1999; Di Natale <i>et al.</i> , 1992; Silvani <i>et al.</i> 1999; Di Natale <i>et al.</i> 1993
Drift nets ("Thonaille") (mesh size 18 to 24 cm)	France, Monaco	May-September	Mediterranean	<i>T. thynnus</i>	<i>S. coeruleoalba</i>	Known	Monitored and extrapolated: Imbert <i>et al.</i> 2001, 2002
Drift nets (mesh size 8 to 16 cm)	Italy	Spring-Autumn	Mediterranean	<i>Sarda sarda</i> , <i>Auxis rochei</i> , other small tuna species.	<i>T. truncatus</i> , <i>Grampus griseus</i>	Known	Estimated total: Di Natale & Notarbartolo di Sciara, 1994
Drift nets (mesh size 4 to 7 cm)	Many coastal areas	Spring	Mediterranean	<i>Scomber</i> spp., <i>Boops boops</i> , and other small pelagic species	<i>S. coeruleoalba</i> , <i>Tursiops truncatus</i>	Suspected: many interactions with fishing gear	
Bottom set gillnets (including coastal trammels)	Many coastal areas	All	Mediterranean	<i>Mullus</i> spp., <i>Sepia</i> spp. Sparidae, <i>Scorpaena</i> spp. other demersal species	<i>Ziphius cavirostris</i> , <i>D. delphis</i> <i>S. coeruleoalba</i> , <i>Grampus griseus</i> , <i>T. truncatus</i> , <i>Physeter macrocephalus</i>	Known: also high level of gear interaction	Di Natale, 1989; Di Natale & Notarbartolo, 1994; Bradai, 2000; Centro Studi Cetacei, 1987-2000; Lauriano <i>et al.</i> , 2001.
Bottom set gillnets	Many deep coastal areas	All	Mediterranean	<i>Palinurus elephas</i> , <i>Merluccius merluccius</i>	<i>T. truncatus</i>	Gear interactions known	CORISA, 1992
Bottom set gillnets for turbot and dogfish	All range Countries	April-June	Black Sea	<i>P. maeotica</i> , <i>Sualus acanthias</i>	<i>Phocoena phocoena</i> , <i>T. truncatus</i>	Known: high impact	Birkun 2002
Bottom set gillnets for sturgeon	All range Countries	April-June	Black Sea	<i>Acipenser</i> spp., <i>Huso huso</i>	<i>Phocoena phocoena</i> , <i>T. truncatus</i> , <i>D. delphis</i>	Known: low impact	Birkun 2002
Bottom set gillnets for turbot	Turkey	April-June	Black Sea	<i>P. maeotica</i> , <i>Sualus acanthias</i>	<i>Phocoena phocoena</i>	Known: high impact	Birkun 2002

Bottom set gillnets for turbot	Turkey	April-June	Black Sea	<i>P. maeotica, Sualus acanthias</i>	<i>T. truncatus</i>	Known: very low impact	Birkun 2002
Middle-water set gillnets	Many coastal areas	All	Mediterranean	<i>Boops boops, Oblada melanura, Trachurus sp., Spicara spp.</i>	<i>T. truncatus</i>	Known	Di Natale pers comm.
Set gillnets for sprat and anchovy	Romania	March-May	Black Sea	<i>S.s. phalaericus, E .e. ponticus</i>	<i>Phocoena phocoena</i>	Known	Birkun 2002
Set gillnets for scad	Romania	July-September	Black Sea	<i>Trachurus spp.</i>	<i>D. delphis</i>	Known	Birkun 2002
Trap nets	Bulgaria, Georgia, Ukraine	May-June	Black Sea		<i>T. truncatus</i>	Very low impact	Birkun 2002
Purse seine	All	All	Mediterranean	<i>Sardina pilchardus, Engraulis encrasiculus, other small pelagic species</i>	<i>T. truncatus</i>	Known: occasional plus many gear interatciouns	Bradai, 2000
Purse seine (mullet and anchovy)	Kerch Strait, Crimea	November-December	Black Sea	<i>M. soiuy, E .e. ponticus</i>	<i>T. truncatus</i>	Low impact	Birkun 2002
Tuna purse seine	Spain, France, Italy, Greece, Tunisia, Turkey, Croatia, Algeria, Morocco	March-October	Mediterranean	<i>Thunnus thynnus</i>	<i>S. coeruleoalba.</i>	Known: rare	Magnaghi & Podesta, 1987; Podestà & Magnaghi, 1989
Tuna traps	Spain, Italy, Tunisia, Libya, Morocco, Croatia	April-July	Mediterranean	<i>Thunnus thynnus</i>	<i>T. truncatus B. acutorostrata, Orcinus orca</i>	Known: Interactions are sporadic	Di Natale, 1992; Bradai, 2000; Di Natale & Mangano, 1983
Bottom trawl	All areas	All	Mediterranean	A large range of demersal species	<i>T. truncatus.</i> A very high number of interactions is reported	Known.	Silvani et al., 1992
Harpoons	Italy, Turkey	April-August	Mediterranean	<i>Xiphias gladius, Thunnus thynnus, Tetrapturus belone</i>	<i>S. coeruleoalba, Grampus griseus, Physeter macrocephalus, Ziphius cavirostris, D. delphis.</i>	Known: reports of deliberate harpooning in the 1980s, no recent cases recorded;	Di Natale, 1992
Drifting long lines	Spain, Italy, Greece, Albania, Turkey, Cyprus, Lebanon, Egypt, Libya, Tunisia,	March-December	Mediterranean	<i>Xiphias gladius, Thunnus thynnus</i>	<i>Stenella coeruleoalba, Grampus griseus, T. truncatus, Pseudorca crassidens, Globicephala melas, Ziphius cavirostris,</i>	Known: probably low level	Duguy et al. 1983; Di Natale & Mangano, 1983; Di Natale, 1992 Di Natale et al., 1993

	Algeria, Morocco, Malta				<i>Physeter macrocephalus</i> , <i>Balaenoptera physalus</i>		
Drifting long lines	Spain, Italy, Greece, Albania	Spring-Autumn	Mediterranean	<i>Thunnus alalunga</i> and other small tunas	<i>S. coeruleoalba</i> , <i>T. truncatus</i> ..	Frequent interactions are already reported	Di Natale <i>et al.</i> , 1992
Pelagic pair trawl	Italy	All	Mediterranean	Pelagic schooling species	<i>T. truncatus</i>	Known	Vallini, pers.com
Pelagic trawl	France, Italy	All	Mediterranean	Demersal species	Delphinids	Suspected, by analogy	No
Pelagic trawl	Georgia, Ukraine	November-December	Black Sea	<i>E .e. ponticus</i>	<i>D. delphis</i>	Known	Birkun 2002
Encircling gillnets	Spain, Italy, Greece	Spring-Summer	Mediterranean	<i>Boops boops</i> , <i>Oblada melanura</i> , <i>Belone belone</i> , <i>Spicara spp.</i> other small and medium size pelagic species	<i>Tursiops truncatus</i>	Suspected	Goodson <i>et al.</i> , 2001
Bottom long lines	Spain, Italy, Greece, Albania,	All	Mediterranean	<i>Merluccius merluccius</i> , Sparidae spp., <i>Lepidopus caudatus</i>		Suspected: fishermen report sporadic interactions	
Rod and reel	Spain, France, Italy	Spring-Summer	Mediterranean	<i>Thunnus thynnus</i>		Suspected: fishermen report sporadic interactions	
Hand-line	Spain, Italy, Greece	Spring-Summer-Autumn	Mediterranean	<i>Thunnus thynnus</i>		Suspected: fishermen have reported a few interactions	
Jigging line	Spain, Italy, Greece	May-September	Mediterranean	<i>Todarodes sagittatus</i> , <i>Illex sp.</i>		Suspected: Very frequent interactions are reported by fishermen	

Based on:

Anonymous (2002). Report of the Second Meeting of the Subgroup on Fishery and Environment (SGFEN) of the Scientific, Technical and Economic Committee for Fisheries (STECF): Incidental catches of small cetaceans. Commission of the European Communities, Brussels, 11-14 June 2002.

Birkun, A., Jr. 2002. Interactions between cetaceans and fisheries in the Black Sea. In: G. Notarbartolo di Sciara (ed.). Cetaceans of the Mediterranean and Black Seas: State of Knowledge and Conservation Strategies. A report to the ACCOBAMS Secretariat, Monaco, February 2002. Section 10, 11 pp.

References

- Anonymous (2003)a. Electroacoustic prototype for controlling the behaviour of marine mammals, IRMA-CNR: 55+ Figures and Annexes.
- Anonymous (2003)b. Trial of acoustic deterrents ('porpoise pingers') for prevention of porpoise (*Phocoena phocoena*) bycatch - Phase 1 - Deployment Trial. Financial Instrument for Fisheries Guidance (FIG) Project; SeaFish Report No CR201, Sea Fish Industry Authority.
- Barlow, J., Cameron, G.A. 2003. Field experiments show tha acoustic pingers reduce marine mammals bycatch in the California drift gill net fishery. Marine Mammal Science 19(2):265-283.
- Berggren, P., J. Carlstrom, et al, 2002. "Mitigation of small cetacean by-catch; evaluation of acoustic alarms (MISNET)." Final Report to the European Commission.
- Bordino, P., Kraus, S., Albareda, D., Fazio, A., Palmerio, A., Mendez, M. and Botta, S. 2003. Reducing incidental mortality of Franciscana dolphin (*Pontoporia blainvillei*) with acoustic warning devices attached to fishing nets. Mar. Mammal Sci. 18(4):833-42.
- Bordino, P., S. Kraus, et al. (2004). Acoustic devices help to reduce incidental mortality of the Franciscana dolphin (*Pontoporia blainvillei*) in coastal gillnets. Presented to the Scientific Committee of the International Whaling Commission, Sorrento, June 2004.
- Carlstrom, J., Berggren P., Dinnetz, F., and Borjesson, P. 2002. A field experiment using acoustic alarms (pingers) to reduce harbour porpoise by-catch in bottom-set gillnets. ICES Journal of Marine Science, 59: 816–824.
- Cox, T. M., A. J. Read, et al. (2001). "Will harbour porpoises (*Phocoena phocoena*) habituate to pingers?" Journal of Cetacean Research and Management 3(1): 81-86.
- Culik, B.M., Koschinski, S., Tregenza, N., Ellis, G.M., 2001. Reactions of harbor porpoises *Phocoena phocoena* and herring *Clupea harengus* to acoustic alarms. Marine Ecology Progress Series 211,255–260.
- Gazo, M., Brotons, JM., Aguilar, A. 2002. Testing low-intensity transponders to mitigate bottlenose dolphin depredation on trammel nets. European Cetacean Society, 16th Annual Conference – Liege, Belgium. Abstract.
- Gearin, P.J., Gosh, M.E., Laake, J., Cooke, L., Delong, R.L. and Hughes, K.M. 2000. Experimental testing of acoustic alarms (pingers) to reduce bycatch of harbour porpoise, *Phocoena phocoena*, in the state of Washington. J. Cetacean Res. Manage. 2(1):1-10.
- Goodson, A. D., R. H. Mayo, et al. (1994). "Field testing passive acoustic devices designed to reduce the entanglement of small cetaceans in fishing gear." Rep. Int. Whal. Commn (Special Issue 15): 597-605.
- Goodson, A.D., Datta, S., Dremiere, P.-Y. and Di Natale, A. (2001). EC contract DGXIV 98/019 - Project ADEPTs, Final Report to the European Commission.
- Gordon, J. & Northridge, S. (2002) Potential impacts of Acoustic Deterrent Devices on Scottish Marine Wildlife. Scottish Natural Heritage Commissioned Report F01AA404.
- Imbert, G., J.-C. Gaertner, et al. (2001). Expertise Scientifique de la Thonaille Medterranee; Rapport d'etape Suivi Scientifique. Marseille, Centre d'Océanologie de Marseilles, Universite de la Mediterranee: 90.
- Imbert, G., J.-C. Gaertner, et al. (2002). Effet des répulsifs acoustiques sur la capture de dauphins dans les thonailles. Marseille, Universite de la Méditerranée Centre D'Océanologie de Marseille: 36.
- Johnston DW, 2002. The effect of acoustic harassment devices on harbour porpoises (*Phocoena phocoena*) in the Bay of Fundy, Canada. Biological Conservation 108, 113-118.
- Kastelein, R.A., Rippe, H.T., Vaughan, N., Schooneman, N.M., Verboom, W.C., de Haan, D., 2000b. The effects of acoustic alarms on the behavior of harbor porpoises (*Phocoena phocoena*) in a floating pen. Marine Mammal Science 16, 46–64.
- Koschinski, S. & Culik, B. 1997. Detering harbour porpoises (*Phocoena phocoena*) from gillnets: Observed reactions to passive reflectors and pingers. Rep. Int. Whal. Commn. 47: 659-668.
- Kraus, S.D., Read, A.J., Solow, A., Baldwin, K., Spradlin, T., Anderson, E., Williamson, J., 1997. Acoustic alarms reduce porpoise mortality. Nature 388, 525.
- Laake, J., Rugh, D. & Baraff, L. 1998. Observations of harbor porpoise in the vicinity of acoustic alarms on a set gill net. NOAA Tech Memo. NMFA-AFSC-84. 40pp.
- Larsen, F. 1999. The effect of acoustic alarms on the by-catch of harbour porpoises in the Danish North Sea gill net fishery. Paper SC/51/SM41 presented to the IWC Scientific Committee Meeting, May 1999, (unpublished), 8pp.

- Larsen, F., Eigaard, O.R. & Tougaard, J. 2002. Reduction of harbour porpoise bycatch in the North Sea by high-density gillnets. Paper SC/54/SM30 presented to the IWC Scientific Committee Meeting, May 2002, (unpublished), 13pp.
- Morton AB & Symonds HK, 2002. Displacement of *Orcinus orca* (L.) by high amplitude sound in British Columbia, Canada. ICES Journal of Marine Science 59(1), 71-80.
- Northridge S., Vernicos D. and Raitzos-Exarchopolous D. 2003. Net depredation by bottlenose dolphins in the Aegean: First attempts to quantify and to minimise the problem. Paper SC/55./SM25 presented to the IWC Scientific Committee Meeting 2003, (unpublished).
- Olesiuk P.F., Nichol L.M., Sowden P.J. & Ford J.K.B. 2002. Effect of the sound generated by an acoustic harassment device on the relative abundance and distribution of harbour porpoises (*Phocoena phocoena*) in Retreat Passage, British Columbia. Marine Mammal Science 18(4) 843-863.
- Reeves, R.R., Hofman, R.J., Silber, G.K. and Wilkinson, D. 1996. Acoustic deterrence of harmful marine mammal-fishery interactions: proceedings of a workshop held in Seattle, Washington, 20-22 March 1996. NOAA Tech. Memo. NMFS-OPR-10. 70pp.
- Reeves, R.R., Read, A.J., Notarbartolo di Sciara, G., 2001. Report of the Workshop on Interactions between Dolphins and Fisheries in the Mediterranean: Evaluation of Mitigation Alternatives. Istituto Centrale per la Ricerca Scientifica e Tecnologica Applicata al Mare, Rome, Italy.
- SMRU, UCC, CFPO and ISWFO (2001) Reduction Of Porpoise Bycatch In Bottom Set Gillnet Fisheries, Final report to European Commission, Study Contract 97/095
- Taylor, V.J, Johnston, D.W. & Verboom, W.C. (1997) Acoustic Harassment Device (AHD) use in the aquaculture industry and implications for marine mammals. In: Proceeding Symposium on Bio-sonar and Bioacoustics, Loughborough University U.K**
- Trippel, E.A, Strong, M.B., Terhune, J.M., Conway, J.D. 1999. Mitigation of harbour porpoise (*Phocoena phocoena*) bycatch in the gillnet fishery in the lower Bay of Fundy. Can. J. Fish Aquat. Sci. 56: 113-123.
- Vernicos, D., Raitzos, D.E. and Northridge, S. (2003). An experimental approach to limit dolphin damage to nets in a trammel net fishery in Greece. p70. Abstract. European Cetacean Society, 17th Conference – Las Palmas de Gran Canaria, 9-13 March 2003.
- Westerberg, H., A. Fjalling, et al. (1999). Evaluation of an acoustic seal-scarer at salmon trapnets in the Baltic. International Conference on Baltic Seals. Parnu.
- Zahri, Y., Abid, N., Elouamari, N., Abdellaoui, B. (2004). Etude de l'interaction entre le grand dauphin et le pêche a la senne coulissante en méditerranée marocaine, Institut National de Recherche Halieutique.