



Minutes of the FRAC OSBPI Working Group Meeting

7 February 2024 – 10:00 to 16:00

Corteva lab, Eschbach, Germany

Participants

Corteva Jean-Luc Genet (Chair)

Przemek Szubstarski

Mamadou Mboup

Syngenta

Stefano Torriani

David Ranner

Irina Metaeva

Bayer

Jürgen Derpmann

Andreas Mehl

Christian Zupanc

Disclaimer

The technical information contained in the global guidelines/the website/the publication/the minutes is provided to CropLife International/RAC members, non-members, the scientific community and a broader public audience.

While CropLife International and the RACs make every effort to present accurate and reliable information in the guidelines, CropLife International and the RACs do not guarantee the accuracy, completeness, efficacy, timeliness, or correct sequencing of such information. CropLife International and the RACs assume no responsibility for consequences resulting from the use of their information, or in any respect for the content of such information, including but not limited to errors or omissions, the accuracy or reasonableness of factual or scientific assumptions, studies, or conclusions.

Inclusion of active ingredients and products on the RAC Code Lists is based on scientific evaluation of their modes of action; it does not provide any kind of testimonial for the use of a product or a judgment on efficacy. CropLife International and the RACs are not responsible for, and expressly disclaim all liability for, damages of any kind arising out of use, reference to, or reliance on information provided in the guidelines.

Listing of chemical classes or modes of action in any of the CropLife International/RAC recommendations must not be interpreted as approval for use of a compound in a given country. Prior to implementation, each user must determine the current registration status in the country of use and strictly adhere to the uses and instructions approved in that country.

Introduction

A FRAC OSBPI Working Group was formed in 2015 to generate common resistance management recommendations for the fungicides oxathiapiprolin and fluoxapiprolin. OSBP fungicides are active against oomycete fungi and used for the control of Phytophthora and downy mildews of numerous crops. OSBPIs inhibit an oxysterol binding protein (OSBP) homologue. Oxysterol binding proteins are implicated in the movement of lipids between membranes, among other processes. Inhibiting OSBP may disrupt other processes in the fungal cell, such as signaling, maintaining cell membranes, and the formation of more complex lipids that are essential for the cell to survive.

Oxathiapiprolin and fluoxapiprolin are cross-resistant.

OSBPIs have been classified under the FRAC Code 49. The resistance risk is medium to high.

FRAC Code	Target site and code	Group name	Chemical group	Common name	Comments
49	F9 lipid homeostasis and transfer/storage	OSBPI oxysterol binding protein homologue inhibition	piperidinyl-thiazole-isoxazolines	Oxathiapiprolin Fluoxapiprolin	Resistance risk assumed to be medium to high (single site inhibitor). Resistance management required.

OSBPI Minutes of the 2024 discussions

Review of sensitivity monitoring 2023

Grape downy mildew (*Plasmopara viticola*)

Data presented by Bayer, Corteva and Syngenta

In 2023, sensitivity data have been generated for samples originating from Austria, Brazil, Bulgaria, China, Croatia, France, Germany, Hungary, Italy, Portugal, Romania, Switzerland, Slovakia, and Spain.

Most 2023 samples in monitored areas were sensitive.

In France, extensive monitoring shows a largely sensitive situation. Again, less sensitive populations were detected in commercial vineyards of Armagnac. A single population with reduced sensitivity was also found in Vallée du Rhône.

In Italy, there was an increase in the detection of less sensitive populations/isolates in Veneto (Treviso), Emilia Romagna (Ravenna) and Friuli-Venezia-Giulia.

Single cases of reduced sensitivity were also found in a small number of trial sites in Austria (Steiermark), south Italy, Croatia, north Portugal, and Spain (Galicia).

Data generated in 2023 data was also presented for grape downy mildew populations from Brazil and China which were all characterized as sensitive.

Molecular characterization of less sensitive isolates collected in 2023 mostly revealed the presence of target site mutations N752I and I792F (homolog to N837I and I877F, based on *P. infestans* homology numbering)

As in previous years, a few samples/isolates carried target site mutations G685R, N752Y and L778W (homolog to 770, 837 and 863 positions based on *P. infestans* homology numbering).

Potato/tomato late blight (*Phytophthora infestans*)

Data presented by Bayer, Corteva and Syngenta

In 2023, sensitivity data have been generated for samples originating from potato and tomato crops in Belgium, Brazil, China, Croatia, Colombia, Denmark, France, Germany, Greece, Hungary, Ireland, Italy, Latvia, Lithuania, Netherlands, Norway,

Poland, Portugal, Romania, Slovakia, Spain, Sweden, United Kingdom, and Vietnam.

All samples collected from tomato crops in Europe were sensitive.

In potato, in 2023, high frequencies of resistance were found in Belgium, West-Germany and the Netherlands. In these countries, reduced efficacy of OSBPI-based products has been observed.

Single resistant samples were found in France, Latvia, Lithuania, Poland, Norway, and the UK with no reports of reduced product performance.

Resistance was also found in potato crops of the Antiochia region of Colombia and in south Vietnam in tomato.

All 2023 samples from potato crops in Brazil, China, Croatia, Italy, Greece, Hungary, Romania, Sweden, Denmark, Ireland, Portugal, and Slovakia Croatia were fully sensitive.

Molecular studies conducted in 2023 revealed the presence of target site mutations G770V and N837I/F/L. Strains with double mutations N837F + G770V were also reported and are also associated with high resistance factors. Mutation I816M was detected for the first time.

Genotypes of *Phytophthora infestans* are classified by Euroblight. These genotypes could express variable levels of aggressiveness, but are not linked to fungicide resistance, which is genetically independent. OSBPI resistance has been found in multiple Euroblight genotypes.

Cucurbit downy mildew (*Pseudoperonospora cubensis*)

Data presented by Corteva and Syngenta

In 2023, sensitivity data have been generated for samples originating from China and Vietnam.

2022 and 2023 molecular studies conducted with samples from Vietnam did not reveal any target site mutations.

In 2023, populations from China with higher EC₅₀ values were detected in Henan, Shandong, Liaoning, Hebei, Guangxi, Guandong, Sichuan, and Yunnan provinces.

Lettuce downy mildew (*Bremia lactucae*)

No 2023 data was presented.

In 2021, sensitivity data have been generated for samples originating from Belgium, Croatia, Germany and Greece. All the samples analyzed were sensitive.

Onion downy mildew (*Peronospora destructor*)

No 2023 data was presented.

In 2020, populations from Bulgaria, Croatia, Italy, Germany, Spain, Greece, Slovenia, Poland, Lithuania, Netherlands and Hungary were analyzed at Syngenta by sequencing of the OSBP gene. None of the target site mutations known to cause reduced sensitivity to OSBPI fungicides in other pathogens were detected.

Sunflower downy mildew (*Plasmopara halstedii*)

Data presented by Corteva

In 2023, sensitivity data have been generated for samples originating from Bulgaria, Greece, France, Romania, and Turkey. As in previous years, all isolates were characterized as sensitive.

Citrus Phytophthora root rot (*Phytophthora palmivora* and *P. nicotianae*)

No 2023 data was presented.

In 2020, samples of *Phytophthora palmivora* and *P. nicotianae* from the USA (Florida) were found to be fully sensitive.

Recommendations for 2024

OSBPI – General Use Recommendations

- Fungicide programs must deliver effective disease management. Apply OSBPIs at effective rates and intervals according to manufacturers' recommendations. Effective disease management throughout the season is a critical component to delay the build-up and spread of resistant pathogen populations.
- Apply OSBPIs only preventatively and in mixtures with effective fungicides from different cross-resistance groups.
- The mixture partner should give effective control of the target disease(s) at the rate and interval selected.
- Foliar exposure to OSBPI products should not exceed thirty-three percent (33%) of the total period of protection needed per crop.

The number of foliar applications of OSBPI products within a total disease management program must be limited as follows:

OSBPI – Grapes

- Make no consecutive applications of OSBPI fungicides.
- Make no more than two (2) applications per season.
- If the total number of applications targeting downy mildew is 4 or less, apply only one (1) application of OSBPI.

OSBPI – Potato

- Make no consecutive applications of OSBPI fungicides.
- Make no more than three (3) applications per season.
- If the total number of applications targeting late blight is 6-10, make no more than two (2) applications per season.
- If the total number of applications targeting late blight is 5 or less, make only one (1) application per season.
- In countries where OSBPI resistance has developed, make no more than two (2) applications per season.

OSBPI – All other crops

- Make no more than four (4) applications or maximum 33% of the total period of protection needed per crop, whichever is more restrictive.
- Where the total number of fungicide applications targeting oomycetes is less than three (3), apply no more than one (1) application of an OSBPI product.
- There should be no more than two (2) foliar applications of any OSBPI product per crop for the control of soil-borne pathogens.
- Applications of OSBPI-containing products are to be made no more than two (2) times in sequence before applying a fungicide with a different mode of action. In areas where the agronomic risk is very high (e.g., continuous cucurbit cropping) and resistance has already been reported, further restrictions to the number of consecutive applications and alternation are recommended.

OSBPI – Seed/soil treatments

- Where an OSBPI fungicide is used as a seed/soil treatment* then, as a precautionary measure, it should be counted as one of the permitted foliar OSBPI applications per crop.
- The first foliar fungicide applied after an OSBPI seed/soil treatment* must not contain an active substance with the OSBPI MoA and must also provide satisfactory disease control against the fungal pathogen(s) targeted by the OSBPI seed/soil treatment*.
- If needed, continue foliar application with alternating fungicidal MoAs for the remaining growing period.

* directed stem sprays are interpreted as foliar not soil application

OSBPI – Multiple crops

- In case of non-cucurbit multiple crops, do not make more than six (6) foliar applications of OSBPI product per year on the same acreage or greenhouse, targeting the same pathogen.
- Specifically, in the case of cucurbit crops, do not make more than four (4) applications per year on the same acreage or greenhouse, targeting *P. cubensis*.

OSBPI – Nursery crops

- OSBPI products must not be used in nursery production of transplanted agricultural crops.

Next meeting:

26 February 2025 (Bayer site in Monheim, Germany).