

**Fergusonite-(Ce)****(Ce, La, Y)NbO<sub>4</sub>**

©2001-2005 Mineral Data Publishing, version 1

**Crystal Data:** [Tetragonal.] *Point Group:* [4/m] [by analogy to fergusonite-(Y)].  
As prismatic dipyrarnidal crystals, to 1 mm, and as irregular poikiloblasts, to 1 cm.

**Physical Properties:** *Fracture:* Conchoidal. *Hardness* = [5.5–6.5] *VHN* = 720–800  
D(meas.) = 5.45–5.48 D(calc.) = n.d.

**Optical Properties:** Semitransparent. *Color:* Dark red to black; reddish in transmitted light.  
*Luster:* Adamantine.  
*Optical Class:* Uniaxial (-).  $\omega = 2.30$   $\epsilon = 2.20$

**Cell Data:** *Space Group:* [I4<sub>1</sub>/a.] *a* = 5.17 *c* = 5.30 *Z* = [2]

**X-ray Powder Pattern:** Novopoltavskii massif, Russia.  
3.1 (10), 1.882 (9), 1.631 (5), 1.566 (4), 2.60 (3), 3.5 (2), 2.76 (2)

**Chemistry:**

	(1)
U <sub>3</sub> O <sub>8</sub>	1.11
Nb <sub>2</sub> O <sub>5</sub>	42.01
Ta <sub>2</sub> O <sub>5</sub>	1.00
TiO <sub>2</sub>	0.05
ThO <sub>2</sub>	0.88
RE <sub>2</sub> O <sub>3</sub>	52.26
Fe <sub>2</sub> O <sub>3</sub>	0.13
CaO	0.81
H <sub>2</sub> O	1.79
Total	[100.04]

(1) Novopoltavskii massif, Russia; original total given as 99.94%; RE = La 12%, Ce 34%, Pr 4%, Nd 21%, Sm 3%, Eu 0.5%, Gd 5%, Tb 0.2%, Dy 1.7%, Ho 0.3%, Er 2%, Tm 0.1%, Yb 1.6%, Y 11.6%; corresponds to [(Ce, La, Y)<sub>0.96</sub>Ca<sub>0.04</sub>Th<sub>0.01</sub>]<sub>Σ=1.01</sub>(Nb<sub>0.98</sub>Ta<sub>0.02</sub>)<sub>Σ=1.00</sub>O<sub>4.00</sub> • 0.30H<sub>2</sub>O.

**Polymorphism & Series:** Dimorphous with fergusonite-beta-(Ce).

**Occurrence:** In a carbonatite (Novopoltavsk massif, Russia); in magnesium-rich skarn of the outer granite contact zone of a carbonatite-derived dolomitic marble (Bayan Obo, China).

**Association:** Magnetite, phlogopite, graphite, chlorite, quartz, pyrite, columbite, allanite, bastnäsite, parisite, apatite, uranpyrochlore, aeschynite, fersmite, monazite, calcite, dolomite (Novopoltavsk massif, Russia).

**Distribution:** In the Chernigovskaya zone, Novopoltavskii massif, Azov Sea area, Ukraine. From the eastern Bayan Obo Fe–Nb–RE deposit, 130 km north of Baotou, Inner Mongolia, China. On Granite Mountain, near Little Rock, Pulaski Co., Arkansas, USA.

**Name:** For its relation to *fergusonite*-(Y), and its dominant *cerium* content.

**Type Material:** A.E. Fersman Mineralogical Museum, Academy of Sciences, Moscow, Russia, 79595.

**References:** (1) Kapustin, Y.L. (1976) New occurrence of accessory cerium fergusonite in carbonatites. *Trudy Mineral. Muzeya Akad. Nauk SSSR*, 25, 166–172 (in Russian). (2) (1977) *Chem. Abs.*, 87, 152 (abs. ref. 1). (3) Kapustin, Y.L. (1986) Rare-earth mineralization in late carbonatites of the Novopoltavsk Massif. *Novye Dannye Mineral.*, 33, 43–52 (in Russian). (4) (1989) *Amer. Mineral.*, 74, 946 (abs. ref. 3). (5) Peishan Zhang and Pejje Tao (1987) Characteristics of the fergusonite- and aeschynite-group minerals in China. *Zhongguo Xitu Xuebao*, 5(1), 1–7 (in Chinese). (6) (1987) *Chem. Abs.*, 107, 241 (abs. ref. 5).

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.