

Original Research

Effective Strategies for Reducing CH₄ and N₂O Emissions from Cow Manure Using Rice Straw as Basal Feed

Hutwan Syarifuddin^{1,2*}, Muhammad Afdal¹, Yun Alwi¹, Ahmad Yani¹, Sri Novianti¹,
Yudha Gusti Wibowo³

¹Animal Science Study Program, Faculty of Animal Husbandry, Jambi University, Jalan Jambi – Muara Bulian KM.15, Mendalo Darat, Jambi Luar Kota District, Muaro Jambi Regency, Jambi, Indonesia

²Environmental Science Study Program, Postgraduate Program, Jambi University, Jalan Arif Rahman Hakim, Telanaipura District, Jambi City, Jambi, Indonesia

³Sustainable Mining and Environmental Research Group, Department of Mining Engineering, Institut Teknologi Sumatera, Lampung, Indonesia

Received: 26 April 2024

Accepted: 3 August 2024

Abstract

The increasing demand for livestock products has led to a significant rise in manure quantities, which negatively impacts the environment by contributing to climate change and increasing greenhouse gas (GHG) emissions. Livestock production primarily generates methane (CH₄) and nitrous oxide (N₂O), which are major emissions from these systems. Various studies have explored effective strategies to reduce CH₄ and N₂O emissions produced from enteric fermentation and manure management. This study aims to develop an effective strategy to reduce CH₄ and N₂O emissions from cattle fed with rice straw. The study was conducted by measuring CH₄, N₂O, temperature, and humidity from cow manure in 50 cattle sheds. Data analysis was performed using the IPCC guidelines, and the analysis of feed ingredients and livestock manure was conducted using proximate analysis. The Analytical Hierarchy Process (AHP) method was employed to determine effective strategies for reducing CH₄ and N₂O emissions. The results indicated that the highest emissions of CH₄ and N₂O were 171.321 ppm and 51,053.539 ppm, respectively, with an average temperature of 29.51°C and humidity of 79.21% RH. Three alternative strategies were identified as effective in decision-making based on their weight order: releasing methane gas into the atmosphere, storing methane gas in the soil, and utilizing methane gas as green energy (biogas).

Keywords: greenhouse gases emission, animals manure emission, climate change

*e-mail: hutwan_syarifuddin@unja.ac.id

Tel.: +627-807-950

Introduction

Rice is one of the mainstay commodities in Indonesia, serving as a staple food and playing a crucial role in the economy by providing income, creating jobs, alleviating poverty, and enhancing food security. In 2023, Indonesia's rice production was approximately 53.63 million tons of dry milled rice (DMR), with every ton of DMR generating 1.5 tons of rice straw. In Jambi Province, the rice harvest area is estimated to reach 61.38 thousand hectares, producing around 274.56 thousand tons of DMR [1]. Typically, farmers burn the rice straw on agricultural land, with only a small portion used as animal feed. The burning of rice straw releases particles (PM_{2.5}, PM₁₀) and greenhouse gases (CO, SO_x, NO_x) into the atmosphere, adversely affecting the environment and human health. This issue requires more in-depth study, especially considering that rice cultivation and biomass burning contribute 97% and 92% to the greenhouse gas (GHG) effect, respectively, with methane emissions from rice fields estimated at over 170 Tg per year [2–4].

Numerous studies have explored the valorization of rice straw for composting, ruminant feed, biofuel production, as an adsorbent, and biochar production. Rice straw can serve as feed for ruminant cattle, such as cows [5]. Cows have fiber-degrading microorganisms in their digestive tracts, allowing them to utilize rice straw, which can reduce feed costs, as feed constitutes nearly 70% of livestock production expenses, providing economic benefits to cattle farmers. In Jambi Province, particularly in Muara Jambi Regency, the average farmer raises 2 to 4 cows per family as a side business.

Rice straw is a cheap, abundant, and practical feed source, with a dry matter content of 84.22%, crude protein of 4.60%, crude fiber of 28.86%, crude fat of 1.52%, and nitrogen-free extract materials of 50.80% [6, 7]. According to [3], rice straw can feed 2–3 beef cattle per hectare per year. However, its high lignin and silica content, low crude protein, poor digestibility, and low palatability limit its use as feed. The type of feed affects the emissions of CH₄, N₂O, and CO₂ from livestock, which are produced by microbial fermentation in the digestive system and manure management [8, 9]. Livestock contributes 9% of CO₂ emissions, 37% of CH₄, and 65% of anthropogenic N₂O globally, totaling 7.1 gigatons of CO₂-e per year, or 14.5% of all anthropogenic GHG emissions [10]. Locally estimating the emissions from different livestock types is essential to understanding their GHG contributions. Research indicates that CH₄ emissions from livestock manure depend on feed type, physiological status, and manure management, with adult cows producing the highest CH₄ and dairy cows the highest N₂O [11, 12]. For example, in West Java, livestock emissions in 2022 were reported as 2.49 Gg tons of CO₂-e CH₄ from enteric fermentation, 0.22 Gg tons of CO₂-e CH₄ from manure management, and 0.87 Gg tons of CO₂-e direct N₂O from fertilizer management [10]. Another study reported CH₄ emissions from manure management at 0.0239 Gg CH₄ per year and direct N₂O emissions at 465.85 kg N₂O per year [13].

Several studies have measured CH₄ and N₂O emissions from cattle using Tier 1 methods, but these are limited to livestock populations and emission factors without a specific focus on Balinese cattle fed with rice straw. This study aims to fill this gap by developing an effective strategy to reduce CH₄ and N₂O emissions from cattle in Muara Jambi Regency, Jambi Province, using rice straw as feed. This approach could increase farmers' income while reducing GHG emissions, representing a novel aspect of this research. Previous research has explored CH₄ reduction using feed additives like 3-nitrooxypropanol (3NOP) and halogenated CH₄ analogs, achieving reductions of 76% and 98%, respectively [11]. Given the significant contribution of livestock to GHG emissions, this study employs the Analytical Hierarchy Process (AHP) method to develop effective strategies for reducing CH₄ and N₂O emissions, a gap not addressed in earlier studies [14]. This research aims to develop an effective strategy to reduce CH₄ and N₂O gas emissions from cattle using rice straw as feed and provide valuable insights for government and policymakers on using rice straw as animal feed and developing effective strategies for emission reduction.

Study Area

The research was conducted from May 2023 to December 2023 in the cattle breeding area of Puduk Village, Kumpeh Ulu District, covering an area of 16.54 km² [1]. Kumpeh Ulu District is located at 1°33'31.77" S and 103° 40'44.24" E, with the research site situated at an elevation of 8–13 meters above sea level. The site was deliberately selected based on the following considerations: 1) it has a cattle population of approximately 110 heads; 2) it includes around 600 hectares of paddy fields; 3) it supports 50 cattle farmers. The map of the research location is shown in Fig. 1.

Data Source

This study was conducted using a descriptive survey approach with 50 farmers who raise cattle traditionally as respondents. The study utilized both primary and secondary data. Primary data was obtained through direct surveys of the respondents. Measurements were taken for temperature and humidity, CH₄ and N₂O gas emissions, dry matter (DM), crude protein (CP), crude fat, crude fiber (CF), ash, acid detergent fiber (ADF), neutral detergent fiber (NDF), and gross energy (GE) from kumpai grass, rice straw, tofu dregs, and manure. Furthermore, a focus group discussion (FGD) was conducted with respondents to develop an effective strategy for reducing CH₄ and N₂O emissions from cattle fed with rice straw basal feed. Consultations were also held with key informants to compile priority sequences and pairwise comparisons between indicators in the AHP method, and to determine criteria, sub-criteria, and alternatives in decision-making.




Fig. 1. Map of the study area.

Data Analysis

The determinants of GHG emissions from the livestock sector are influenced by several factors, including livestock population, type of livestock, housing, type of animal feed, manure/waste management, and farmer behavior in cattle cultivation. The formula for calculating greenhouse gas emissions follows the Tier-1 method [15]. To calculate CH₄ emissions from the feed fermentation process in the rumen (enteric fermentation), the formula issued in the IPCC guidebook [15, 16] is used:

$$\text{Enteric CH}_4 \text{ emission (CO}_2\text{-e } \frac{\text{ton}}{\text{hrad}}) = \text{livestock population} \times FFe \times \frac{21}{1000} \quad \text{Eq. 1}$$

$$\text{Manure CH}_4 \text{ emission (CO}_2\text{-e } \frac{\text{ton}}{\text{hrad}}) = \text{livestock population} \times FFe \times \frac{21}{1000} \quad \text{Eq. 2}$$

$$\text{N}_2\text{O (CO}_2\text{e } \frac{\text{tons}}{\text{years}}) = \text{population} \times (0.05 \times FEn) / \left(\frac{1000}{BB}\right) \times 365 \times 44/28 \times 293/100 \quad \text{Eq. 3}$$

Where Fee = Enteric emission factor (kg CH₄/head/day), Fem = Emission factor from manure (kg CH₄/head/day), 21/1000 = conversion for CH₄ to CO₂ and from kg to tons, Fen = N₂O emission factor from manure (kg N₂O/kg manure/day), 293/1000 = conversion for N₂O to CO₂ and from kg to tons, 44/28 = Convert from N₂O-N to N₂O; 0.05 = Average excretion N (kg N/head/year).

To calculate GHG emissions using the Tier-1 method, the IPCC provides default emission factors (FE) for various types of livestock. For Indonesia, the FE values for CH₄ emissions from enteric fermentation and manure management are used as the default factors for the entire Asian region [10, 15, 17]. The GHG emissions from the livestock sector are calculated using the IPCC Tier-1 method in three parts: CH₄ emissions from enteric fermentation, CH₄ emissions from manure management, and N₂O emissions from manure management. By combining these three calculations, the total GHG emissions from the livestock sector can be determined. To analyze the concentrations of CH₄ and N₂O in samples, gas chromatography is used, equipped with a Flame Ionization Detector (FID) for CH₄ and an Electron Capture Detector (ECD) for N₂O. The emissions of CH₄ and N₂O gases are then calculated using established equations [18]:

$$E = \frac{dc}{dt} \times \frac{Vch}{Ach} \times \frac{mW}{mV} \times \frac{273.2}{273.2 + T} \quad \text{Eq. 4}$$

Table 1. Scale to compare factors in AHP paired comparison.

Factor i Compared to Factor j	Quantitative Value
Equally important	1
More important	3
Much more important	5
Quite more important	7
Absolutely more important	9
Intermediate values	2, 4, 6, 8

Table 2. Standardized Random Index (RI) values of mean random consistency index.

Hierarchy Matrix	1	2	3	4	5	6	7	8	9	10
RI	0	0	0,58	0,90	1,12	1,12	1,32	1,41	1,45	1,49

In the provided equations, E represents the emissions of CH₄ and N₂O gases, measured in milligrams per square meter per day. The term dc/dt denotes the difference in CH₄ and N₂O concentration over time, expressed in parts per million per minute. V_{ch} stands for the volume of the chamber box, measured in cubic meters, while A_{ch} represents the area of the chamber box, measured in square meters. The variables mW signify the molecular weight of CH₄, measured in grams, with specific values provided for CH₄ and N₂O at 0°C. The term mV indicates the molecular volume of CH₄, measured in liters. Finally, T represents the average temperature during gas sampling, measured in degrees Celsius.

The Analytical Hierarchy Process (AHP) method is used for developing effective strategies in decision-making [19]. This method involves several stages:

Stage 1: Determining critical factors. The AHP method helps establish the sequence of criteria to be considered and make the most reasonable decisions [20, 21]. According [22] to the process, the comparative value of factors is determined based on a priority comparison scale. On this scale (Table 1), pairwise comparisons between equally important factors have a quantitative value of 1. A more important factor has a value of 3, a much more important factor has a value of 5, a quite more important factor has a value of 7, and an absolutely more important factor has a value of 9. Intermediate values of 2, 4, 6, and 8 are used for factors that fall between these levels of importance.

Stage 2: Normalizing the matrix. To normalize the criterion importance matrix, the value of each cell in a column is divided by the total value of that column. The weighted average (W_i) is then calculated by summing the weights of the X_i factor relative to X_j after normalization and dividing by n. To ensure the reliability of the weight (W_i), the consistency index CR (consistency

ratio) is calculated. A CR value of less than 0.1 indicates a consistent result. The formula for calculating CR is provided in the corresponding equations.

$$CR = \frac{CI}{RI} \quad \text{Eq. 5}$$

Where:

$$CI = \frac{\lambda - 1}{n - 1}$$

$$RI = \frac{CI1 + CI2 + \dots + CIn}{n} \quad \text{Eq. 6}$$

$$\lambda_{max} = \frac{1}{n} \left(\frac{\sum_{n+1}^n W_{1n}}{W_{11}} + \frac{\sum_{n+1}^n W_{2n}}{W_{12}} + \frac{\sum_{n+1}^n W_{nn}}{W_{1n}} \right) \quad \text{Eq. 7}$$

The Random Index (RI) is used as a reference value, as shown in Table 2. The eigenvalue of the matrix is denoted as λ_{max} . To calculate the total S value, use the following formula:

$$s_i = \sum_{j=1}^n (\text{weight of } X_i \text{ relative to } X_j) \quad \text{Eq. 8}$$

Stage 3 involves creating a hierarchy of the total S values. This is achieved by applying reclassification methods and regression algorithms to classify the total S values according to their respective ranges, depending on the study's content. The hierarchical structure for decision-making, aimed at effectively reducing CH₄ and N₂O emissions from cattle fed with rice straw, is presented in Fig. 2.

Results and Discussion

Cattle contribute significantly to global warming by producing methane (CH₄) and nitrous oxide (N₂O)




Fig. 2. Analytical hierarchical structure of effective strategies for reducing CH₄, N₂O gas emissions from cattle.

gases. The greenhouse gas (GHG) effects from livestock are influenced by factors such as livestock type, population, feed management, and manure management. The concentrations of CH₄ and N₂O gases from manure in 50 cattle sheds were measured using the MQ4 sensor method (Table 3).

Methane is produced during the decomposition of organic matter under oxygen-deficient conditions, particularly in enteric fermentation processes, livestock manure, and rice fields [18, 23, 24]. N₂O is generated from microbial transformations in soil and livestock manure, with increased

Table 3. Average Gas Concentrations of CH₄, CO₂, N₂O from the measurement results using MQ4 sensors in 50 cattle sheds.

No	CH ₄ (ppm)	CO ₂ (ppm)	N ₂ O (ppm)	Temperature (°C)	Moisture (Rh)
1	98.612	2,465.302	29,386.394	30.66	76.60
2	102.621	2,565.515	30,580.939	30.97	78.96
3	100.601	2,515.015	29,978.979	30.85	76.15
4	110.971	2,774.265	33,069.239	29.57	76.89
5	113.131	2,828.265	33,712.919	29.57	76.89
6	133.911	3,347.765	39,905.359	29.52	79.21
7	159.971	3,999.265	47,671.239	29.51	79.21
8	168.431	4,210.765	50,192.319	29.51	79.21
9	131.481	3,287.015	39,181.219	29.54	79.21
10	149.151	3,728.765	44,446.879	29.53	79.21
11	165.581	4,139.515	49,343.019	29.52	79.21
12	171.321	4,283.015	51,053.539	29.51	79.21
13	154.501	3,862.515	46,041.179	29.52	79.21
14	143.941	3,598.515	42,894.299	29.51	79.21
15	117.541	2,938.515	35,027.099	29.53	77.62
16	124.371	3,109.265	37,062.439	29.36	81.67
17	165.581	4,139.515	49,343.019	29.35	82.56
18	165.581	4,139.515	49,343.019	29.46	82.43
19	171.321	4,283.015	51,053.539	29.13	83.12
20	171.321	4,283.015	51,053.539	29.32	83.16
21	149.151	3,728.765	44,446.879	30.97	78.96
22	143.941	3,598.515	42,894.299	30.85	76.15
23	119.791	2,994.765	35,697.599	29.57	76.89
24	133.911	3,347.765	39,905.359	29.57	76.89
25	171.321	4,283.015	51,053.539	29.52	79.21
26	165.581	4,139.515	49,343.019	29.53	79.21
27	138.861	3,471.515	41,380.459	29.52	79.22
28	129.081	3,227.015	38,466.019	29.51	79.22
29	119.791	2,994.765	35,697.599	29.52	79.21
30	124.371	3,109.265	37,062.439	29.51	79.25
31	129.081	3,227.015	38,466.019	29.53	77.62
32	133.911	3,347.765	39,905.359	31.20	82.17
33	133.911	3,347.765	39,905.359	32.13	82.65
34	92.822	2,320.552	27,660.974	30.15	80.16
35	124.371	3,109.265	37,062.439	29.57	80.01
36	138.861	3,471.515	41,380.459	29.86	79.82
37	129.081	3,227.015	38,466.019	29.55	79.65
38	92.802	2,320.052	27,655.014	29.67	78.89
39	94.712	2,367.802	28,224.194	30.14	81.13



No	CH ₄ (ppm)	CO ₂ (ppm)	N ₂ O (ppm)	Temperature (°C)	Moisture (Rh)
40	83.662	2,091.552	24,931.294	31.22	82.53
41	87.242	2,181.052	25,998.134	31.51	82.45
42	90.922	2,273.052	27,094.774	29.51	79.23
43	87.242	2,181.052	25,998.134	29.54	79.33
44	110.971	2,774.265	33,069.239	29.53	79.26
45	98.612	2,465.302	29,386.394	29.52	79.43
46	102.621	2,565.515	30,580.939	29.51	79.27
47	102.621	2,565.515	30,580.939	29.52	79.29
48	98.612	2,465.302	29,386.394	29.51	79.31
49	94.712	2,367.802	28,224.194	29.53	77.64
50	75.162	1,879.052	22,398.294	29.36	81.77

Note: par per million (ppm)

Table 4. Results of proximate analysis of feed ingredients and manure during the study.

Feed ingredients and manure	DM (%)	Crude Protein (%)	Crude Fat (%)	Crude Fiber (%)	Abu (%)	ADF (%)	NDF (%)	GE (cal/g)
Tofu Dregs	91.755	20.151	3.055	12.936	3.826	21.963	43.277	2.768
Kumpai grass	89.452	8.335	5.595	28.859	11.329	48.466	64.868	3.310
Paddy raw	96.133	4.803	3.713	27.646	14.720	58.045	98.004	3.553
Solid	95.734	14.900	11.835	25.230	21.283	53.392	57.360	2.652
Manure	97.657	12.246	2.908	32.819	9.947	36.311	52.820	2.388

Note: Dry Matter (DM), Acid detergent fiber (ADF), Neutral detergent fiber (NDF), Gross Energy (GE)

emissions occurring when nitrogen availability exceeds plant needs, especially in wet conditions [15].

The emissions of CH₄ and N₂O gases produced by cattle are also affected by the type of feed consumed, such as forage and concentrates. Common forage includes kumpai grass and rice straw, with additional feed often comprising tofu dregs and solids. CH₄ concentrations in manure reached a maximum of 171.321 ppm, while N₂O concentrations peaked at 51,053.539 ppm. The production of these gases is influenced by the crude fiber content of the feed, as high crude fiber levels increase methanogenesis activity during digestion [25].

According to studies, CH₄ emissions from livestock manure management in Jambi Province were as follows: 2869.38 CO₂-e tons per head in 2014, 3060.96 CO₂-e tons per head in 2015, 3217.60 CO₂-e tons per head in 2016, 3350.47 CO₂-e tons per head in 2017, and 3342.95 CO₂-e tons per head in 2018 [13]. The proximate analysis of feed ingredients and manure during the study is presented in Table 4.

Table 4 shows that the crude fiber content of kumpai grass and rice straw is high, at 28.859% and 27.646%, respectively. High crude fiber content affects the activity of rumen microbes in degrading fiber, as reflected in the Acid Detergent Fiber (ADF) and Neutral Detergent Fiber (NDF) values. Previous studies report crude fiber content in kumpai grass and rice straw at 32.85% and 32.99%–36.97%, respectively [26–28]. High crude fiber content leads to increased CH₄ gas production from enteric fermentation and in manure.

The addition of fatty acids and feed additives can impact methane gas production in ruminants [29–31]. According to research [32], providing fermented tofu dregs significantly reduces CH₄ gas flux, as fermented tofu pulp has a higher crude protein content (25.87%) compared to unfermented tofu pulp (21.08%). Emission reduction strategies in small ruminant farming can include improving feed efficiency, providing feed supplements, and utilizing food waste or agro-industrial by-products [33–35].




Fig. 3. Criteria weighting.

Table 5. Inconsistency and consistency ratio (CR) values by criteria.

Criteria	Inconsistency	Consistency (CR)
Government	0.09	0.91
Private	0.09	0.91
Farmer Rancher	0.09	0.91

Effective Strategies to Reduce CH₄ and N₂O Gases from Manure that Gets Rice Straw Feed

In strategizing the management of CH₄ and N₂O gas emissions from livestock manure, three key variables are identified: the government, the private sector, and livestock farmers. The government plays a crucial role in decision-making, while the private sector uses livestock products, and livestock farmers manage the cattle. The inconsistency value is 0.09, and the consistency ratio (CR) is 0.91. A high consistency value indicates the validity of the research. The weights of the three variables are illustrated in Fig. 3. Fig. 3 shows that the government has the largest role, with a weight of 0.687, followed by livestock farmers and the private sector. This significant role of the government is due to its involvement in formulating policies, regulations, and decisions for cattle farming and manure management. Manure contains organic matter such as proteins, carbohydrates, and fats, which can be used as a source of feed and energy for anaerobic bacteria growth. Livestock manure is composed of fatty acids, proteins, and carbohydrates, which are easily decomposed [36]. According to Indonesia's policy, the country aims for an unconditional target of a 29% reduction in emissions and a conditional target of up to 41% compared to

business-as-usual by 2030 [10]. The inconsistency and consistency ratio values are presented in Table 5.

Based on the criteria, nine sub-criteria synergize in managing CH₄ gas derived from cow manure. The inconsistency and consistency ratio values are shown in Table 6, and the weight of the sub-criteria is displayed in Fig. 4. Fig. 4 indicates that the number of cattle raised has the highest weight of 0.748, making it the dominant factor in producing CH₄ and N₂O gas emissions from manure. Livestock production is influenced by host-rumen microbiome-environment linkages, which ultimately benefit the animal's adaptation to the environment [23, 37, 38]. Current technology for managing livestock waste into energy at the research site is minimal. Alternatives for managing CH₄ gas are not optimal, as indicated by the pairwise comparison in Fig. 5. Fig. 5 shows that the weight of CH₄ gas released directly into the atmosphere is the highest at 5.426, followed by gas stored in the soil (1.962), and the weight for using CH₄ as green energy in the form of biogas is only 1.611.

Methane is estimated to contribute about 18% of total expected global warming in the next 50 years [39], with livestock accounting for approximately 9% of total global emissions [15]. Domestic animals contribute about 94% of total global animal emissions [36]. Although emissions

Table 6. Inconsistency value and consistency ratio of sub-criteria.

No	Sub Criteria	Inconsistency	Consistency Ratio (CR)
1	Number of cattle livestock	0.01	0.99
2	Land Availability	0.09	0.91
3	Capital in the cultivation of cattle	0.05	0.95
4	Farmer knowledge in methane gas management	0.09	0.91
5	Diseases and death of cattle	0.05	0.95
6	Demand for cattle production	0.05	0.95
7	Climate change	0.05	0.95
8	Cattle rearing system	0.05	0.95
9	Technology Management of livestock waste into energy	0.05	0.95




Fig. 4. Sub criteria weights.




Fig. 5. Alternative weights.

per unit of animal product have decreased, total emissions have increased due to the growth in livestock populations worldwide [40]. By 2050, total CH₄ emissions from ruminants are expected to increase significantly due to the rising demand for milk and meat [41].

To reduce CH₄ and N₂O gas emissions from manure with rice straw basal feed, several actions can be taken. These include increasing the government's role in promoting and using environmentally friendly energy sources like methane gas, increasing the number of cattle raised, enhancing farmer knowledge, utilizing available land through livestock integration with agriculture and plantations, reducing livestock mortality and disease rates, increasing the use of appropriate technology, assisting in livestock cultivation (e.g., CSR initiatives from private companies), enhancing livestock adaptation to climate changes, and increasing demand for livestock production to meet animal protein needs. Other strategies to reduce CH₄ emissions include using CH₄ inhibitors, tanniferous forages, electron sinks, oils and fats, and oilseeds, which can decrease daily methane emissions by an average of 21% [42]. The alternative approach is to reduce CH₄ gas released directly into the atmosphere by utilizing biogas as a source of green energy. Additionally, cattle feces and urine can be used as organic fertilizers [43]. Cow manure provides essential nutrients for plants, and cow urine can be processed into biourine for plant fertilization or sale [13].

Conclusions

The highest recorded emissions of CH₄ and N₂O gases were 171.321 ppm and 51.053.539 ppm, respectively, observed under conditions of 29.51°C temperature and 79.21% relative humidity. Effective strategies to reduce these emissions rely significantly on government regulations and policies. Three alternative approaches that have proven effective include reducing the amount of methane released into the atmosphere, storing methane in the soil, and utilizing methane as a source of green energy. Therefore, policymakers should focus on preventing the burning of rice straw and promoting the development of renewable energy sources, such as biogas derived from livestock manure, to mitigate the impact of these harmful emissions.

Acknowledgements

This work funded by Kementerian Pendidikan, Kebudayaan, Riset dan Teknologi through the research scheme of "Master of Environmental Science, Jambi University" under contract number 397/UN 21.11/PT.01.05/SPK/2023, April 17, 2023.

Conflict of Interest

The authors declare no conflict of interest.

References

1. BPS-STATISTICS MUARO JAMBI REGENCY. Muaro Jambi Regency In Figures. Jambi Province. BPS-Statistics Muaro Jambi Regency, pp. 123 , 2024.
2. SURMAINI E., RUNTUNUWU E., LAS I. Efforts of agricultural sector in dealing with climate change. *Jurnal Penelitian dan Pengembangan Pertanian*, **30** (1), 1, 2011.
3. ROMASANTA R.R., SANDER B.O., GAIHRE Y.K., ALBERTO M.A.C., GUMMERT M., QUILTY J., NGUYEN V.H., CASTALONE A.G., BALINGBING C., SANDRO J., CORREA T.J.R., WASSMANN R. How does burning of rice straw affect CH₄ and N₂O emissions? A comparative experiment of different on-field straw management practices. *Agriculture, Ecosystems & Environment*, **239**, 143, 2017.
4. MUJTAHID A.L.H.M., ABDULLAH Y.A.L.H.M., MEZBAUL B.M., ASLINA B.N., JAWAD I.A.M., MOFIJUL I.S.M. Enhancing bioelectricity generation and mitigating methane emissions in paddy fields: A novel approach using activated biochar in plant microbial fuel cells. *Energy Conversion and Management*, **307**, 118327, 2024.
5. NGUYEN T.T., NGO H.T., HA Q.Q., NGUYEN T.Q., LE T.Q., NGUYEN S.H., PHAM C.T., ZIEGLER T., VAN SCHINGEN-KHAN M., LE M.D. Molecular phylogenetic analyses and ecological niche modeling provide new insights into threats to the endangered Crocodile Lizard (*Shinisaurus crocodilurus*). *Frontiers of Biogeography*, **14** (1), 1, 2022.
6. NURLIZA., DOLOROSA E., HAMID A., YUSRA A. Rice Farming Performance for Sustainable Agriculture and Food Security in West Kalimantan. *AGRARIS. Journal of Agribusiness and Rural Development Research*, **3** (2), 84, 2017.
7. HAN X., SUN X., WANG C., WU M., DONG D., ZHONG T., THIES J.E., WU W. Mitigating methane emission from paddy soil with rice-straw biochar amendment under projected climate change. *Scientific Reports*, **6** (1), 1, 2016.
8. DONADIA A.B., TORRES R.N.S., SILVA H.M.D., SOARES S.R., HOSHIDE A.K., OLIVEIRA A.S.D. Factors Affecting Enteric Emission Methane and Predictive Models for Dairy Cows. *Animals*, **13** (11), 1857, 2023.
9. TSETEN T., SANJORJO R.A., KWON M., KIM S.W. Strategies to Mitigate Enteric Methane Emissions from Ruminant Animals. *Journal of Microbiology and Biotechnology*, **32** (3), 269, 2022.
10. SHIDDIEQY M.I., WIDIWATI Y., ROHAENI E.S., SASONGKO W.T., HADIATRY M.C., RIYANTI S. Emissions and trends of livestock greenhouse gases in West Java, Indonesia, *AIP Conference Proceedings*, **2957**, 070027, 2024.
11. KELLY L., KEBREAB E. Recent advances in feed additives with the potential to mitigate enteric methane emissions from ruminant livestock. *Journal of Soil and Water Conservation*, **78** (2), 111, 2023.
12. ALDRIAN E., PUSPOWARDOYO S., HARYANTO B. Emisi gas rumah kaca dari peternakan di Indonesia dengan metode TIER 2 IPCC. Jakarta: LIPI Press, pp. 153, 2019.
13. SYARIFUDDIN H., RAHMAN A.S.Y., DEVITRIANO D. Inventarisasi Emisi Gas Rumah Kaca (CH₄ dan N₂O) Dari Sektor Peternakan Sapi Dengan Metode Tier-1 IPCC di Kabupaten Muaro Jambi. *Jurnal Ilmiah Ilmu-Ilmu Peternakan*, **22** (2), 84, 2019.

14. DEHAGHI I.M., SALMANMAHINY A., KARIMI S., SHABANI A.A. Multi-criteria evaluation and simulated annealing for delimiting high priority habitats of *Alectoris chukar* and *Phasianus colchicus* in Iran. *Animal Biodiversity and Conservation*, **41** (1), 185, **2018**.
15. YONA L., CASHORE B., JACKSON R.B., OMETTO J., BRADFORD M.A. Refining national greenhouse gas inventories. *Ambio*, **49** (10), 1581, **2020**.
16. TEDESCHI L.O., ABDALLA A.L., ÁLVAREZ C., ANUGA S.W., ARANGO J., BEAUCHEMIN K.A., BECQUET P., BERNDT A., BURNS R., DE CAMILLIS C., CHARÁ J., ECHAZARRETA J.M., HASSOUNA M.C.D., SIGN L., KENNY D., MATHOT M., MAURICIO R.M., MCCLELLAND S.C., NIU M., ONYANGO A.A., KEBREAB E. Quantification of methane emitted by ruminants: a review of methods. *Journal of Animal Science*, **100** (7), 1, **2022**.
17. KOUAZOUNDE J.B., GBENOU J.D., BABATOUNDE S., SRIVASTAVA N., EGGLESTON S.H., ANTWI C., BAAH J., MCALLISTER T.A. Development of methane emission factors for enteric fermentation in cattle from Benin using IPCC Tier 2 methodology. *Animal*, **9** (3), 526, **2015**.
18. PRAMONO A., ADRIANY T.A., SUSILAWATI H.L. Mitigation scenario for reducing greenhouse gas emission from rice field by water management and rice cultivars. *Journal of Tropical Soils*, **25** (2), 53, **2020**.
19. NGUYEN S.H., NGUYEN D.N., THU N.N., PHAM H.H., PHAN H.A., DAO C.D. Current Soil Degradation Assessment in the Thua Thien Hue Province, Vietnam, by Multi-Criteria Analysis and GIS Technology. *Sustainability*, **15**, 14276, 1, **2023**.
20. ZANGMENE F.L., NGAPNA M.N., ATEBA M.C.B., MBOUDOU G.M.M., DEFO P.L.W., KOUO R.T., DONGMO A.K., OWONA S. Landslide susceptibility zonation using the analytical hierarchy process (AHP) in the Bafoussam-Dschang region (West Cameroon). *Advances in Space Research*, **71** (12), t5282, **2023**.
21. SREENIVASAN A., SURESH M., NEDUNGADI P.R.R.R. Mapping analytical hierarchy process research to sustainable development goals: Bibliometric and social network analysis. *Heliyon*, **9** (8), e19077, **2023**.
22. AKTER S., HANI U., DWIVEDI Y.K., SHARMA A. The future of marketing analytics in the sharing economy. *Industrial Marketing Management*, **104**, 85, **2022**.
23. NGUYEN T.T.M., BADHAN A.K., REID I.D., RIBEIRO G., GRUNINGER R., TSANG A., GUAN L.L., MCALLISTER T. Comparative analysis of functional diversity of rumen microbiome in bison and beef heifers. *Applied and Environmental Microbiology*, **89** (12), 1, **2023**.
24. ROHAENI E.S., MAILENA L., LESMAYATI S., ERMUNA S.S. Sustainability analysis for rice and duck farming in swampy land, Hulu Sungai Utara Regency, South Kalimantan. *IOP Conference Series: Earth and Environmental Science*, **648** (1), 012134, **2021**.
25. PATRA A., PARK T., KIM M., YU Z. Rumen methanogens and mitigation of methane emission by anti-methanogenic compounds and substances. *Journal of Animal Science and Biotechnology*, **26** (8), 13, **2017**.
26. MARJUKI M., SUNTORN W. Precision feeding management: New approach for better and more sustainable livestock production. *Journal of Science and Agricultural Technology*, **2** (2), 16, **2021**.
27. RISWANDI R., HAMZAH B., WIJAYA A., ABRAR A. Bali Heifers Performance on Cassava Leaves, Palm Oil Sludge and Yeast Supplementation in a Ration Based on Kumpai Grass (*Hymenachne amplexicaulis* (Rudge) Nees). *Advances in Animal and Veterinary Sciences*, **8** (8), 1, **2020**.
28. DHILLON A.K., SHARMA N., DOSANJH N.K., GOYAL M., MAHAJAN G. Variation in the nutritional quality of rice straw and grain in response to different nitrogen levels. *Journal of Plant Nutrition*, **41** (15), 1946, **2018**.
29. NDARU P.H., HUDA A.N., MASHUDI M. Pengaruh Penambahan Asam Lemak Pada Pakan Ternak Ruminansia Terhadap Kandungan Nutrisi Pakan. *TERNAK TROPIKA Journal of Tropical Animal Production*, **22** (1), 12, **2021**.
30. PALANGI V., LACKNER M. Management of Enteric Methane Emissions in Ruminants Using Feed Additives: A Review. *Animals*, **12** (24), 3452, **2022**.
31. UNGERFELD E. Opportunities and Hurdles to the Adoption and Enhanced Efficacy of Feed Additives Towards Pronounced Mitigation of Enteric Methane Emissions from Ruminant Livestock. *Methane*, **1**, 262, **2022**.
32. SYARIFUDDIN H., RAHMAN A.S.Y., DEVITRIANO D. CH₄ Gas Mitigation Strategy with the Use of Interpretative Structural Modeling Method. *Proceedings of the 3rd Green Development International Conference (GDIC 2020)*, Atlantis Press, pp. 474, **2021**.
33. GIAMOURI E., ZISIS F., MITSIOPOULOU C., CHRISTODOULOU C., PAPPAS A.C., SIMITZIS P.E., KAMILARIS C., GALLIOU F., MANIOS T., MAVROMMATIS A., TSIPLAKOU E. Sustainable Strategies for Greenhouse Gas Emission Reduction in Small Ruminants Farming. *Sustainability*, **15** (5), 4118, **2023**.
34. HONAN M., FENG X., TRICARICO J.M., KEBREAB E. Feed additives as a strategic approach to reduce enteric methane production in cattle: modes of action, effectiveness and safety. *Animal Production Science*, **62** (14), 1303, **2021**.
35. MERCADANTE M.E.Z., DE MELO CALIMAN A.P., CANESIN R.C., BONILHA S.F.M., BERNDT A., FRIGHETTO R.T.S., MAGNANI E., BRANCO R.H. Relationship between residual feed intake and enteric methane emission in Nelore cattle. *Revista Brasileira de Zootecnia*, **44** (7), 255, **2015**.
36. ARIFAN F., ABDULLAH, SUMARDIYONO S. Methane gas production from a mixture of cow manure, chicken manure, cabbage waste, and liquid tofu waste using the anaerobic digestion method. *IOP Conference Series: Earth and Environmental Science*, **623** (1), 012036, **2021**.
37. WANG W., DONG Y., GUO W., ZHANG X., DEGEN A.A., BI S., DING L., CHEN X., LONG R. Linkages between rumen microbiome, host, and environment in yaks, and their implications for understanding animal production and management. *Frontiers in Microbiology*, **15**, **2024**.
38. KAUR H., KAUR G., GUPTA T., MITTAL D., ALI S.A. Integrating Omics Technologies for a Comprehensive Understanding of the Microbiome and Its Impact on Cattle Production. *Biology*, **12** (9), 1200, 1, **2023**.
39. WILKES A., DIJK S.V. Tier 2 inventory approaches in the livestock sector: A collection of agricultural greenhouse gas inventory practices. *Global Research Alliance on Agricultural Greenhouse Gases CGIAR Climate Change Agriculture and Food Security, UNIQUE Forestry and Land Use*, **1** (10), 30, **2018**.
40. OPIO C., GERBER P., MOTTET A., FALCUCCI A., TEMPIO G., MACLEOD M., VELLINGA T., HENDERSON B., STEINFELD H. Greenhouse gas emissions from ruminant supply chains — a global life cycle assessment. Rome, Italy: Food and Agriculture Organization of the United Nations (FAO), **2013**.
41. GERBER P.J., STEINFELD H., HENDERSON B., MOTTET A., OPIO C., DIJKMAN J., FALCUCCI A.,

- TEMPIO G. Tackling climate change through livestock a global assessment of emissions and mitigation opportunities. Rome, Italy: Food and Agriculture Organization of the United Nations (FAO), **2013**.
42. ARNDT C., HRISTOV A.N., PRICE W.J., MCCLELLAND S.C., PELAEZ A.M., CUEVA S.F., OH J., DIJKSTRA J., BANNINK A., BAYAT A.R., CROMPTON L.A., EUGENE M.A., ENAHORO D., KEBREAB E., KREUZER M., MCGEE M., ECILE MARTIN C., NEWBOLD C.J., REYNOLDS C.K., SCHWARM A., SHINGFIELD K.J., VENEMAN J.B., YÁÑEZ-RUIZ D.R., YU Z. Full adoption of the most effective strategies to mitigate methane emissions by ruminants can help meet the 1.5 °C target by 2030 but not 2050. *Proceedings of the National Academy of Sciences of the United States of America (PNAS)*, **119** (20), e2111294119, **2023**.
43. VAGHAR SEYEDIN S.M., ZEIDI A., CHAMANEHPOUR E., NASRI M.H.F., VARGAS-BELLO-PÉREZ E. Methane Emission: Strategies to Reduce Global Warming in Relation to Animal Husbandry Units with Emphasis on Ruminants. *Sustainability*, **14** (24), 16897, **2022**.